

The Continuing Search for a Succinylcholine Replacement

THE introduction of succinylcholine into clinical practice in 1951 was a seminal development in the history of anesthesia. Since that time, anesthesiologists have had access to a neuromuscular blocking drug (muscle relaxant) with a very rapid onset and a duration of action of approximately 10 min. The clinical utility of succinylcholine has been counterbalanced by its many undesirable effects, ranging from the relatively benign (fasciculations) to life-threatening events (hyperkalemia or malignant hyperthermia).¹ The adverse effects of succinylcholine are almost all related to its depolarizing mechanism of action. Consequently, there has been a continual search for a nondepolarizing muscle relaxant that can replicate both the rapid onset and short duration of succinylcholine. In this issue of the Journal, three articles report on preclinical and preliminary clinical investigations of such a drug, GW280430A.²⁻⁴

Structurally, GW280430A is an asymmetric mixed-tetrahydroisoquinolinium chlorofumarate.⁵ Of the drugs in clinical practice, the one to which it bears the closest structural resemblance is mivacurium. The exact mechanism underlying its short duration is not fully elucidated; it appears to undergo rapid degradation in the plasma by chemical (nonenzymatic) mechanisms.⁶ GW280430A has been studied in a variety of animal species—cat, dog, and rhesus monkey—with promising results for onset and duration and a good safety profile with regard to histamine-releasing potential.^{3,4} In assessing the possible clinical utility of this drug, its apparent strengths must be weighed against its potential for adverse effects.

In humans, the onset of GW280430A over the range of 1.8–4.0xED₉₅ doses is 1.5–1.7 min and it has a very narrow range (1.1–2.0 min).² This narrow range of onset

times is remarkable for a nondepolarizing relaxant and resembles more the tight distribution of onset time seen with succinylcholine.⁷ However, this result may be an artifact due to the small number and physiologic homogeneity of the volunteer subjects studied; it is unlikely to represent the true distribution in patients. Regardless, the onset, although perhaps not as rapid as that of succinylcholine, may be sufficient to facilitate rapid-sequence intubation in most clinical situations.⁸ Of some interest is that onset time did not diminish with doses greater than 1.9xED₉₅. This may again be due to the small number of subjects, or it might be an example of the diminishing return observed as onset times approach the limit determined by circulation time and diffusion of drug into the neuromuscular junction.

In addition to its rapid onset, GW280430A has a very short duration of action. In a dose of 0.4 mg/kg (2.2xED₉₅), recovery to a train-of-four ratio of 0.9 took only 14 min. This may be regarded as essentially complete recovery for a nondepolarizing muscle relaxant.^{9,10} An equivalent degree of recovery for succinylcholine would be represented by the single twitch recovering to between 90% and 100% of predrug level. For a standard intubating dose of 1.0 mg/kg, this time interval is between 8 and 12 min.^{7,11} It appears that in respect to its duration of action, GW280430A resembles succinylcholine more closely than does any previous nondepolarizing muscle relaxant. Of particular interest, pharmacologic antagonism with edrophonium 0.5 mg/kg can decrease significantly the already short recovery time of GW280430A.² The possibility exists, therefore, that edrophonium-accelerated recovery from GW280430A might have a time course that replicates that of succinylcholine.

Increasing the dose of GW280430A does not seem to carry a great penalty in respect to the increase in duration of action.² The numbers in each dose group are too small to make definitive conclusions, but, for example, increasing the dose from 0.36 to 0.72 mg/kg increases recovery time to a train-of-four ratio of 0.9, from 11.9 to only 15.1 min. The other notable aspect of the drug is a lack of cumulative effect. The recovery slopes, as represented by the 5–95% or 25–75% intervals, are unaffected by the dose or duration of administration.² In summary, the time course of action of GW280430A more closely resembles succinylcholine than does any other nondepolarizing muscle relaxant to date.

In addition to its positive characteristics, there are some potential problems with GW280430A.² The structural group to which GW280430A belongs has the pro-

This Editorial View accompanies the following articles: Belmont MR, Lien CA, Tjan J, Bradley E, Stein B, Patel SS, Savarese JJ: Clinical pharmacology of GW280430A in humans. *ANESTHESIOLOGY* 2004; 100:768–73; Savarese JJ, Belmont MR, Hashim MA, Mook RA Jr, Boros EE, Samano V, Patel SS, Feldman PL, Schultz J-AI, McNulty M, Spitzer T, Cohn DL, Morgan P, Wastila WB: Preclinical pharmacology of GW280430A (AV430A) in the rhesus monkey and in the cat: A comparison with mivacurium. *ANESTHESIOLOGY* 2004; 100:835–45; Heerd PM, Kang R, The' A, Hashim M, Mook RJ Jr, Savarese JJ: Cardiopulmonary effects of the novel neuromuscular blocking drug GW280430A (AV430A) in dogs. *ANESTHESIOLOGY* 2004; 100:846–51.

Accepted for publication July 22, 2003. The author is not supported by, nor maintains any financial interest in, any commercial activity that may be associated with the topic of this editorial.

pensity to release histamine,¹² and GW280430A appears to share this. In doses of 3xED95 and above, histamine release with consequent hypotension, tachycardia, and flushing were noted. What does this mean for the clinical potential of the drug? In all likelihood, clinicians would accept a *low* risk of histamine-related effects if the drug could truly replace succinylcholine. It appears however, that histamine release may be an actual clinical problem.

One of the principal uses for a drug such as GW280430A is to facilitate rapid tracheal intubation. The onset of GW280430A is slightly slower than that of succinylcholine, so clinicians will tend to increase doses to promote good conditions for tracheal intubation. This phenomenon has been documented with mivacurium,^{13,14} rocuronium,¹³ and cisatracurium,¹⁵ in which doses in the range of 2.5–5.0xED95 are used to speed onset of paralysis prior to tracheal intubation. This experience suggests that clinicians push doses of muscle relaxants upward to improve conditions for tracheal intubation. It is likely that should GW280430A enter clinical use, doses of 3xED95 or greater would be administered. In this dose range, it seems that significant histamine release with consequent adverse effects would occur.

The recent experience with rapacurium is relevant to consideration of GW280430A and its clinical potential.¹⁶ To summarize, rapacurium is a steroidal muscle relaxant with a rapid onset and short duration.¹⁷ In clinical trials, there was an incidence of bronchospasm of approximately 9%.¹⁸ This was not thought to be of sufficient clinical importance to prevent the general release of the drug. Under the realities of widespread clinical use, there occurred several cases of life-threatening bronchospasm with rapacurium, particularly in children.¹⁶ As a result, 19 months after its release rapacurium was withdrawn by the manufacturer. Is there evidence of such a potential problem with GW280430A? The answer, unfortunately, is *yes*. GW280430A clearly can stimulate release of histamine. In the study of Belmont *et al.*, clinically significant histamine release occurred in one of four subjects who received 0.54 mg/kg (3.0xED95) and in three of four volunteers receiving 0.72 mg/kg (4xED95). These doses might conceivably be used clinically.²

What can we conclude from these articles? They demonstrate that developing a nondepolarizing replacement for succinylcholine is a realistic possibility. GW280430A has a time course of action very close to that of succinylcholine. There is, however, a large question mark

over its histamine-releasing potential. GW280340A may never be released into clinical practice, but it is quite conceivable that a drug closely related to it will be.

James E. Caldwell, M.B.Ch.B. Department of Anesthesia and Perioperative Care, University of California, San Francisco, San Francisco, California. caldwell@anesthesia.ucsf.edu

References

1. Savarese JJ, Caldwell JE, Lien CA, Miller RD: Pharmacology of muscle relaxants and their antagonists. *Anesthesia*, 5th edition. Edited by Miller RD. New York, Churchill Livingstone, 2000, pp 412–90
2. Belmont MR, Lien CA, Tjan J, Bradley E, Stein B, Patel SS, Savarese JJ: Clinical pharmacology of GW280430A in humans. *ANESTHESIOLOGY* 2004; 100:768–73
3. Heerdt PM, Kang R, The' A, Hashim M, Mook RJ Jr, Savarese JJ: Cardiopulmonary effects of the novel neuromuscular blocking drug GW280430A (AV430A) in dogs. *ANESTHESIOLOGY* 2004; 100:846–51
4. Savarese JJ, Belmont MR, Hashim MA, Mook RA Jr, Boros EE, Samano V, Patel SS, Feldman PL, Schultz J-AI, McNulty M, Spitzer T, Cohn DL, Morgan P, Wastila WB: Preclinical pharmacology of GW280430A (AV430A) in the rhesus monkey and in the cat: A comparison with mivacurium. *ANESTHESIOLOGY* 2004; 100:835–45
5. Boros EE, Samano V, Ray JA, Thompson JB, Jung DK, Kaldor I, Koble CS, Martin MT, Styles VL, Mook RA Jr, Feldman PL, Savarese JJ, Belmont MR, Bigham EC, Boswell GE, Hashim MA, Patel SS, Wisowaty JC, Bowers GD, Moseley CL, Walsh JS, Reese MJ, Rutkowski RD, Seifer AM, Spitzer TD: Neuromuscular blocking activity and therapeutic potential of mixed-tetrahydroisoquinolinium halofumarates and halosuccinates in rhesus monkeys. *J Med Chem* 2003; 46:2502–15
6. McNulty M, Brown A, Johnson T, Spitzer TD, Savarese J: The ultra-short acting nondepolarizing relaxant GW280430A undergoes rapid degradation by chemical mechanisms. *ANESTHESIOLOGY* 2000; 93:A-1037
7. Wright PM, Caldwell JE, Miller RD: Onset and duration of rocuronium and succinylcholine at the adductor pollicis and laryngeal adductor muscles in anesthetized humans. *ANESTHESIOLOGY* 1994; 81:1110–5
8. Kopman AF, Klewicka MM, Neuman GG: Reexamined: The recommended endotracheal intubating dose for nondepolarizing neuromuscular blockers of rapid onset. *Anesth Analg* 2001; 93:954–9
9. Kopman AF, Yee PS, Neuman GG: Relationship of the train-of-four fade ratio to clinical signs and symptoms of residual paralysis in awake volunteers. *ANESTHESIOLOGY* 1997; 86:765–71
10. Eriksson LI, Sundman E, Olsson R, Nilsson L, Witt H, Ekberg O, Kuylenstierna R: Functional assessment of the pharynx at rest and during swallowing in partially paralyzed humans: Simultaneous videomanometry and mechanomyography of awake human volunteers. *ANESTHESIOLOGY* 1997; 87:1035–43
11. Dhonneur G, Kirov K, Slavov V, Duvaldestin P: Effects of an intubating dose of succinylcholine and rocuronium on the larynx and diaphragm: An electromyographic study in humans. *ANESTHESIOLOGY* 1999; 90:951–5
12. Savarese JJ, Wastila WB: The future of the benzylisoquinolinium relaxants. *Acta Anaesthesiol Scand Suppl* 1995; 106:91–3
13. Pino RM, Ali HH, Denman WT, Barrett PS, Schwartz A: A comparison of the intubation conditions between mivacurium and rocuronium during balanced anesthesia. *ANESTHESIOLOGY* 1998; 88:673–8
14. Motamed C, Donati F: Intubating conditions and blockade after mivacurium, rocuronium and their combination in young and elderly adults. *Can J Anaesth* 2000; 47:225–31
15. Doenicke AW, Czeslick E, Moss J, Hoernicke R: Onset time, endotracheal intubating conditions, and plasma histamine after cisatracurium and vecuronium administration. *Anesth Analg* 1998; 87:434–8
16. Rajchert DM, Pasquariello CA, Watcha MF, Schreiner MS: Rapacurium and the risk of bronchospasm in pediatric patients. *Anesth Analg* 2002; 94:488–93
17. Kahwaji R, Bevan DR, Bikhazi G, Shanks CA, Fragen RJ, Dyck JB, Angst MS, Matteo R: Dose-ranging study in younger adult and elderly patients of ORG 9487, a new, rapid-onset, short-duration muscle relaxant. *Anesth Analg* 1997; 84:1011–8
18. Moore EW, Hunter JM: The new neuromuscular blocking agents: Do they offer any advantages? *Br J Anaesth* 2001; 87:912–25

Did Experimenter Bias Conceal the Efficacy of Spinal Opioids in Previous Studies with the Spinal Nerve Ligation Model of Neuropathic Pain?

THE current issue of ANESTHESIOLOGY contains a report of a laboratory study¹ that addresses the question, "Is it appropriate to treat neuropathic pain with spinal opioids?" This editorial examines the data that address this question, and the possibility that the differences between laboratory studies reporting positive and negative effects may be attributed, at least in part, to experimenter bias. We recognize that efficacy with opioids, and perhaps other analgesics, may be lost over time with chronic treatment, and that the results of opioids may also differ according to patient population, laboratory model, and route of administration. Therefore, we focus this discussion on whether spinal delivery of opioids has analgesic effects, regardless of the duration of action, in the commonly used spinal nerve ligation model of neuropathic pain.²

Initial clinical reports of chronic intrathecal morphine treatment, published 20 yr ago, showed both acute and sustained efficacy in patients with chronic pain, including neuropathic pain, although dose escalation occurred with time.^{3,4} At the same time, others reported little or no analgesic effects of spinal morphine in neuropathic pain patients.⁵ Later reports suggested that spinal opioids produce partial pain relief in neuropathic pain patients,^{6,7} and other studies reported that intrathecal opioids produce substantial or good analgesic effects in patients with neuropathic pain, even after long-term administration.^{8,9} Thus, the clinical literature regarding efficacy of spinal opioids for neuropathic pain has evolved from initial good efficacy, to poor efficacy, and, more recently, back to good efficacy in many patients.

A laboratory model of neuropathic pain in rats in which ligation of low lumbar spinal nerves results in reduced withdrawal threshold to tactile stimulation of the paw, akin to mechanical allodynia in patients with neuropathic pain, was first described over a decade ago.² Initial studies using this model showed it to be sympathetic-dependent, with hypersensitivity resolving after chemical or surgical sympathectomy.¹⁰ Several laboratories, including ours, showed that the spinal nerve liga-

tion model was also resistant to treatment with intrathecal morphine.¹¹⁻¹³ These were considered important validation studies because neuropathic pain was at that time thought to frequently be sympathetic-dependent and, as noted above, resistant to treatment with intrathecal morphine.

Neuropathic pain is now considered to be only infrequently sympathetic-dependent and often responds to spinal opioids, and these changes in understanding of clinical neuropathic pain have been mirrored by changes in results obtained in this spinal nerve ligation model. For example, the current authors more recently noted that the effect of sympathectomy was smaller than originally described and more variable, depending on rat strain,¹⁴ and we¹⁵ and others¹⁶ failed to observe an effect of sympathectomy on hypersensitivity using this model. In addition, the current report¹ demonstrates full efficacy of intrathecal morphine to reduce hypersensitivity to mechanical stimulation in rats with spinal nerve ligation or spared nerve injury, another model of neuropathic pain. The dose of intrathecal morphine found to be effective was small, and, as noted by the authors, was similar to that needed to treat nonneuropathic acute and chronic pain in these species.

It is somewhat reassuring that the data from the spinal nerve ligation model now seem to be consistent with the growing consensus that neuropathic pain is not very sympathetic-dependent and that spinal opioids are often effective in patients with neuropathic pain. However, these findings also raise the concern that the results obtained in studies with the spinal nerve ligation model of neuropathic pain may be affected by the expectations of the experimenters at the time the studies are conducted. In fact, Zhao *et al.*¹ were careful to use blinding procedures to prevent experimenter bias from affecting their results, and they suggest that experimenter bias could account for the negative results of earlier studies. Results of studies conducted in our laboratory support this suggestion. We previously reported that intrathecal morphine was ineffective after spinal nerve ligation,¹² but in a recent, rigorously blinded replication of our initial nonblinded study, we detected analgesic effects of intrathecal opioids (unpublished data) at doses similar to those reported by Zhao *et al.*¹

In addition to the differences in blinding procedures, many other factors could also contribute to the differences between the results of these studies. For example, even seasonal changes in the source of the protein included in commercial rodent chow—with no change in

This Editorial View accompanies the following article: Zhao C, Tall JM, Meyer RA, Raja SN: Antiallodynic effects of systemic and intrathecal morphine in the spared nerve injury model of neuropathic pain in rats. ANESTHESIOLOGY 2004; 100:905-11.

Accepted for publication January 17, 2004. Supported in part by grants NS39426 and NS41386 from the National Institutes of Health, Bethesda, Maryland.

the total protein or calorie content—can significantly affect the results of studies with these laboratory pain models.¹⁷ It is possible that such subtle differences in diet or some other unidentified and uncontrolled factor could also contribute to the differences in the results between these studies, so it is not appropriate to conclude that all of the differences in the results of the studies with the spinal nerve ligation model discussed above can be attributed to experimenter bias in the nonblinded studies.

However, experimenter bias is a real phenomenon that has been clearly demonstrated in clinical research. For example, in a clinical trial for multiple sclerosis, patient evaluations performed by nonblinded clinicians detected statistically significant therapeutic effects, but patient evaluations performed in the same study by blinded clinicians revealed that the treatment did not produce beneficial effects.¹⁸ Apparent therapeutic effects are often observed in initial, small, nonblinded clinical trials, but not in larger, blinded trials. For example, in a series of clinical studies conducted to test the therapeutic potential of a monoclonal antibody for rheumatoid arthritis, all of the initial, small, nonblinded trials reported therapeutic effects of the antibody.⁸ Subsequently, three large, blinded experiments, some of them conducted by the same investigators who had conducted some of the initial nonblinded studies, all failed to detect any beneficial effect of the treatment.¹⁹

This scenario is so common in clinical trials that it is not considered at all surprising. In these cases, the beneficial results of the treatment reported in the nonblinded trials are typically attributed to experimenter bias. For this reason, decisions about whether clinical treatments are truly effective are often based only on the most well controlled studies. For example, systemic opioids are now considered one of the first-line treatments for neuropathic pain, but that decision was based on only five recent studies, all of which were large, randomized, double-blind, placebo-controlled clinical trials (for review see Dworkin *et al.*²⁰). Clearly, there is a need for strict controls to prevent experimenter bias, which we recognize when testing the potential efficacy of treatments in humans.

Classic studies published more than 40 yr ago demonstrated that experimenter bias could also affect the results of laboratory studies. For example, in a conditioning study, planaria were given a 3-s light cue followed by a 1-s shock. Observers recorded whether the planaria made contractions or head turns during the light cue, which were the dependent measures used to determine if the planaria were learning to associate the light with the shock. Increasing numbers of contractions and head turns in response to the light would be evidence of classic conditioning. Observers who had been told to expect rapid conditioning recorded significantly increased numbers of anticipatory movements, compared

with observers who had been told to expect little or no evidence of conditioning.²¹ In a learning study in rats, experimenters trained rats to run to the dark arm of a T-maze for a food reward. Rats were given 10 trials each day for 5 days. Experimenters who were told to expect their rats to demonstrate the task quickly did in fact record faster acquisition and better performance in their rats than experimenters who had been told to expect that their rats would be poor learners.²² In another learning study, rats were trained to perform a series of tasks for a food reward in an operant or “Skinner” box. If the experimenter had been told that the rats had been bred for good performance in these tasks, better performance was recorded for the rats than if the experimenter had been told that the rats had been bred for poor performance in these tasks.²³ Anyone who has conducted tail flick or formalin testing in rodents recognizes that the expectations of the investigator can affect the results of laboratory pain tests as well. In the case of paw withdrawal in the relatively unrestrained animal, which often raises the paw to walk, groom itself, or make postural adjustments (*e.g.*, in the Hargreave or von Frey test), the influence of investigator bias can be even greater.

Experimenter bias is not limited to the time when the behavioral observations are recorded; it can also affect nonbehavioral measurements. For example, standard laboratory procedures used to count blood cells were shown to require a degree of consistency that was not possible, given the equipment and procedures used, yet laboratory results typically conformed to these unrealistic requirements for consistency; this could only have occurred as a result of the bias of laboratory technicians.²⁴ In addition to affecting initial measurements, experimenter bias can also play a role in the way the data are managed and analyzed. Experimenters can include or exclude data based in part on experimenter expectations, and they can perform additional analyses, analyzing the data as raw scores or differences from baselines, and dividing subjects into any number of subgroups based on a range of different criteria.^{25–27} All of these procedures serve to drastically inflate the probability of detecting an expected effect by increasing the probability of producing a false-positive result.

We suggest that experimenter bias should receive more attention in laboratory research, and that blinding procedures to guard against experimenter bias should be a more common practice in laboratory investigations. Blinding procedures should be used when recording any measurements, not just in behavioral testing, and all decisions about inclusion/exclusion of data should be made before the blind is broken. We would also caution that some discussion or plan should be considered as to how the data will be analyzed before studies are completed, to reduce the risk that the rate of false-positive results will be inflated through the use of numerous

unplanned analyses. Furthermore, we propose that the precise procedures used to prevent experimenter bias should be an essential component of the methods described in laboratory publications. Investigators should explicitly report the details of the procedures they used. A general statement that "the experimenter was blinded" is not sufficient. Finally, we also support the suggestion that laboratory studies should be evaluated in the same way that clinical trials are, with more emphasis on the difference between blinded and nonblinded studies and more weight given to studies that have carefully controlled for potential bias.²⁸

In summary, recent blinded experiments detected clear analgesic effects of spinal opioids in the spinal nerve ligation model. Previous, nonblinded experiments (one of them from our own laboratory) reported that spinal opioids were not effective for neuropathic pain, which was the expected result at that time. The differences in the results between these studies may have been attributed to experimenter bias in the earlier nonblinded experiments, a suggestion that is supported by the results of carefully blinded experiments conducted recently in our own laboratory. Experimenter bias is a well-recognized phenomenon in clinical trials, and although it does not receive as much attention in laboratory research, there should be no doubt that laboratory studies are just as vulnerable to experimenter bias as clinical studies. Thus, the very narrow focus of this editorial, on studies of spinal opioids in the spinal nerve ligation model of neuropathic pain, leads to consideration of a much more general issue—experimenter bias in laboratory research—and the suggestion that the use of rigorous blinding procedures may be just as important in laboratory experiments as they are in clinical trials.

James C. Eisenach, M.D.* Mark D. Lindner, Ph.D.† * Department of Anesthesiology and the Center for the Study of Pharmacological Plasticity in the Presence of Pain, Wake Forest University School of Medicine, Winston-Salem, North Carolina. † Pharmaceutical Research Institute, Bristol-Myers Squibb, Wallingford, Connecticut. eisenach@wfubmc.edu

References

- Zhao C, Tall JM, Meyer RA, Raja SN: Antiallodynic effects of systemic and intrathecal morphine in the spared nerve injury model of neuropathic pain in rats. *ANESTHESIOLOGY* 2004; 100:905–11
- Kim SH, Chung JM: An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat. *Pain* 1992; 50:355–63
- Coombs DW, Maurer LH, Saunders RL, Gaylor M: Outcomes and complications of continuous intraspinal narcotic analgesia for cancer pain control. *J Clin Oncol* 1984; 2:1414–20
- Penn RD, Paice JA: Chronic intrathecal morphine for intractable pain. *J Neurosurg* 1987; 67:182–6
- Arner S, Arner B: Differential effects of epidural morphine in the treatment of cancer-related pain. *Acta Anaesthesiol Scand* 1985; 29:32–6
- Krames ES: Intraspinal opioid therapy for chronic nonmalignant pain: Current practice and clinical guidelines. *J Pain Symptom Manage* 1996; 11:333–52
- Samuelsson H, Malmberg F, Eriksson M, Hedner T: Outcomes of epidural morphine treatment in cancer pain: Nine years of clinical experience. *J Pain Symptom Manage* 1995; 10:105–12
- Winkelmuller M, Winkelmuller W: Long-term effects of continuous intrathecal opioid treatment in chronic pain of nonmalignant etiology. *J Neurosurg* 1996; 85:458–67
- Hassenbusch SJ, Stanton-Hicks M, Covington EC, Walsh JG, Guthrey DS: Long-term intraspinal infusions of opioids in the treatment of neuropathic pain. *J Pain Symptom Manage* 1995; 10:527–43
- Kim SH, Chung JM: Sympathectomy alleviates mechanical allodynia in an experimental animal model for neuropathy in the rat. *Neurosci Lett* 1991; 134:131–4
- Bian D, Nichols ML, Ossipov MH, Lai J, Porreca F: Characterization of the antiallodynic efficacy of morphine in a model of neuropathic pain in rats. *NeuroReport* 1995; 6:1981–4
- Lavand'homme PM, Eisenach JC: Exogenous and endogenous adenosine enhance the spinal antiallodynic effects of morphine in a rat model of neuropathic pain. *Pain* 1999; 80:31–6
- Lee YW, Chaplan SR, Yaksh TL: Systemic and supraspinal, but not spinal, opioids suppress allodynia in a rat neuropathic pain model. *Neurosci Lett* 1995; 199:111–4
- Xie J, Park SK, Chung K, Chung JM: The effect of lumbar sympathectomy in the spinal nerve ligation model of neuropathic pain. *J Pain* 2001; 2:270–8
- Lavand'homme P, Pan HL, Eisenach JC: Intrathecal neostigmine, but not sympathectomy, relieves mechanical allodynia in a rat model of neuropathic pain. *ANESTHESIOLOGY* 1998; 89:493–9
- Ringkamp M, Eschenfelder S, Grehl EJ, Habler HJ, Meyer RA, Janig W, Raja SN: Lumbar sympathectomy failed to reverse mechanical allodynia- and hyperalgesia-like behavior in rats with L5 spinal nerve injury. *Pain* 1999; 79:143–53
- Shir Y, Ratner A, Raja SN, Campbell JN, Seltzer Z: Neuropathic pain following partial nerve injury in rats is suppressed by dietary soy. *Neurosci Lett* 1998; 240(2):73–6
- Noseworthy JH, Ebers GC, Vandervoort MK, Farquhar RE, Yetsis E, Roberts R: The impact of blinding on the results of a randomized, placebo-controlled multiple sclerosis clinical trial. *Neurology* 1994; 44:16–20
- Epstein WV: Expectation bias in rheumatoid arthritis clinical trials: The anti-CD4 monoclonal antibody experience. *Arthritis Rheum* 1996; 39:1773–80
- Dworkin RH, Backonja M, Rowbotham MC, Allen RR, Argoff CR, Bennett GJ, Bushnell MC, Farrar JT, Galer BS, Haythornthwaite JA, Hewitt DJ, Loeser JD, Max MB, Saltarelli M, Schmader KE, Stein C, Thompson D, Turk DC, Wallace MS, Watkins LR, Weinstein SM: Advances in neuropathic pain: Diagnosis, mechanisms, and treatment recommendations. *Arch Neurol* 2003; 60:1524–34
- Cordaro L, Ison JR: Psychology of the scientist: X. Observer bias in classical conditioning of the planarian. *Psychol Rep* 1963; 13:787–9
- Rosenthal R, Fode KL: The effect of experimenter bias on the performance of the albino rat. *Behav Sci* 1963; 8:183–9
- Rosenthal R, Lawson R: A longitudinal study of the effects of experimenter bias on the operant learning of laboratory rats. *J Psychiatr Res* 1963; 2:61–72
- Berkson J, Magath TB, Hurn M: The error of estimate of the blood cell count as made with the hemocytometer. *Am J Physiol* 1940; 128:309–23
- Assmann SF, Pocock SJ, Enos LE, Kasten LE: Subgroup analysis and other (mis)uses of baseline data in clinical trials. *Lancet* 2000; 355:1064–9
- Pocock SJ, Hughes MD, Lee RJ: Statistical problems in the reporting of clinical trials: A survey of three medical journals. *N Engl J Med* 1987; 317:426–32
- Pocock SJ, Assmann SE, Enos LE, Kasten LE: Subgroup analysis, covariate adjustment and baseline comparisons in clinical trial reporting: Current practice and problems. *Stat Med* 2002; 21:2917–30
- Lindner MD, Frydel BR, Francis JM, Cain CK: Analgesic effects of adrenal chromaffin allografts: Contingent on special procedures or due to experimenter bias? *J Pain* 2003; 4:64–73