

Low Doses of Fentanyl Block Central Sensitization in the Rat Spinal Cord In Vivo

Justus Benrath, M.D.,* Christina Brechtel,† Eike Martin, M.D., Ph.D.,‡ Jürgen Sandkühler, M.D., Ph.D.§

Background: μ -Opioid receptor agonists are strong analgesics. However, their usefulness for preemptive analgesia is controversial. The authors tested antinociceptive and preemptive properties of fentanyl as a μ -opioid receptor agonist in a model of spinal nociception *in vivo*.

Methods: C fiber-evoked potentials were recorded in the superficial laminae I-II of the rat lumbar spinal cord with glass microelectrodes in response to electrical stimulation of the sciatic nerve. High-frequency stimulation was applied on the sciatic nerve to induce long-term potentiation of C fiber-evoked field potentials, a form of central sensitization. To test the effect of fentanyl on acute nociception, fentanyl was infused intravenously at increasing doses ($6\text{--}192\ \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$). One hour after start of infusion, high-frequency stimulation was applied to evaluate effects of fentanyl on the induction of long-term potentiation.

Results: In the absence of fentanyl, high-frequency stimulation potentiated C fiber-evoked field potentials to $149 \pm 12\%$ of controls (mean \pm SEM; $n = 6$) for at least 1 h. Increasing doses of fentanyl led to a significant reduction of C fiber-evoked potentials in a dose-dependent manner. The induction of long-term potentiation was blocked by low doses of fentanyl (infusion $12\text{--}48\ \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$). At high doses, fentanyl did not block the induction of long-term potentiation (infusion $96\text{--}192\ \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$).

Conclusions: Low doses of fentanyl block the synaptic form of central sensitization in the rat spinal cord *in vivo*, but higher doses do not have this effect.

CHRONIC pain after surgery has been a major problem because 20% of patients attending chronic pain clinics implicated surgery as one of the causes of their chronic pain and, in approximately half of these cases, surgery was the sole cause.¹ Surgical damage is able to induce hyperalgesia (abnormal intense pain elicited by noxious stimulation) and allodynia (pain induced by normally nonpainful stimuli). These enhanced responses to noxious or nonnoxious stimuli result from peripheral² or central sensitization.³⁻⁵ The sensitization of peripheral nociceptors is typically limited in time to the period of primary injury. Central sensitization may, however, out-

last the period of noxious input to the central nervous system by hours to weeks, and in some patients, the period of abnormal pain sensitivity may last even longer. Recent evidence suggests that synaptic long-term potentiation (LTP) can be specifically induced at synapses between C-fiber afferents and neurons in superficial spinal dorsal horn that mediate hyperalgesia.⁶ This is considered a cellular mechanism of central sensitization and hyperalgesia (see Sandkühler⁴ for review). LTP can be induced in the spinal cord by peripheral trauma, inflammation, or nerve injury. LTP of synaptic strength in nociceptive pathways may aggravate postoperative hyperalgesia and allodynia but may also lead to phantom pain and chronic postsurgical pain.¹ Although postoperative pain and chronicity of surgical pain are of enormous clinical importance, the mechanisms of the induction and the prevention of LTP are not fully understood.

Calcium influx into spinal cord neurons is required for LTP induction. Strong excitation of dorsal horn neurons is essential for calcium influx through voltage-gated calcium channels and voltage- and ligand-gated glutamatergic receptors of the *N*-methyl-D-aspartate (NMDA) subtype. Consequently, activity-dependent calcium influx can be reduced by any means that prevents strong excitation of nociceptive spinal dorsal horn neurons, *e.g.*, by agonists at μ -opioid receptors (MORs). Opioid agonists were shown to inhibit voltage-gated Ca^{2+} currents in neurons⁷ and to reduce excitation by presynaptic and postsynaptic inhibition.⁸ Therefore, blockade opioid receptors could be clinically important to prevent LTP induction and thereby spinal sensitization in patients. In our investigation, we tested the clinically used MOR agonist fentanyl on its ability to influence the induction and maintenance of LTP.

Methods

Animals

After obtaining approval from the Institutional Animal Care Committee (Regierungspräsidium Karlsruhe, Germany), the experiments were performed on male Sprague-Dawley rats weighing 250–320 g (Charles River Deutschland, Sulzfeld, Germany). Animals were kept under temperature-controlled environmental conditions on a 12:12 h light:dark cycle and were fed a standard diet (Altromin C 1000; Altromin, Lage, Germany) with access to food and water *ad libitum*.

Preparation of the Animals

To establish LTP under different anesthetics, halothane (3.5 vol% inspiratory) in two thirds N_2O and one third O_2

* Resident, † Ph.D. Student, ‡ Professor and Chairman, Klinik für Anaesthesiologie, Universität Heidelberg. § Professor, Institut für Hirnforschung, Universität Wien, Wien, Austria.

Received from the Klinik für Anaesthesiologie, Universität Heidelberg, Heidelberg, Germany. Submitted for publication April 29, 2002. Accepted for publication January 8, 2004. Supported by the Pain Research Program of the Medical Faculty of the University of Heidelberg, Heidelberg, Germany; grant No. Sa 435/10-2 from the Deutsche Forschungsgemeinschaft, Bonn, Germany; and institutional sources of the Klinik für Anaesthesiologie, University of Heidelberg, Heidelberg, Germany. Presented at the 30th Annual Meeting of the Society for Neuroscience, New Orleans, Louisiana, November 4–9, 2000, and the 31st Annual Meeting of the Society for Neuroscience, San Diego, California, November 10–15, 2001. Dr. Benrath and Ms. Brechtel contributed equally to the article.

Address correspondence to Dr. Benrath: Klinik für Anästhesie und Allgemeine Intensivmedizin B, Universität Wien, Währinger Gürtel 18-20, AKH, A-1090 Wien, Austria. Address electronic mail to: justus.benrath@univie.ac.at. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

was used to induce and urethane was used (1.2 g/kg intravenous) to maintain anesthesia, or isoflurane in two thirds N_2O and one third O_2 was used to induce (3.5 vol% inspiratory) and maintain (1.0 vol% expiratory) anesthesia, or sevoflurane in two thirds N_2O and one third O_2 was used to induce (6 vol% inspiratory) and to maintain (1.7 vol% inspiratory) anesthesia. Halothane in two thirds N_2O and one third O_2 was used to induce and urethane was used to maintain anesthesia in all experiments in which fentanyl was used. Concentrations of inhalational anesthetics were monitored with a gas analyser (Capnomac Ultima; Datex-Ohmeda, Duisburg, Germany). The surgical level of anesthesia was verified by stable arterial blood pressure and constant heart rate during noxious stimulation of the interdigital area of the paw.

A polyethylene catheter (Portex Ltd., Hythe, Kent, United Kingdom) was inserted into the right femoral vein for application of drugs and for constant infusion of 0.9% normal saline and 0.1% sodium bicarbonate. The right femoral artery was cannulated with a polyethylene catheter and was continuously infused by a heparin solution (2.5 IE/ml) to monitor arterial blood pressure and heart rate (Plugsys; Hugo Sachs, March, Germany). Mean arterial blood pressure and heart rate were continuously recorded with a pen recorder (L 120E; Linseis, Selb, Germany). The trachea was intubated with a 14-gauge intravenous cannula to allow mechanical ventilation. Muscle relaxation was achieved with $1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ pancuronium bromide. Atropin (0.05 mg/kg) was given once at the beginning of the preparation to reduce tracheal secretion. Glucose (5%, 10 ml/kg) was injected subcutaneously for nutrition of the animals during the 8–12 h of experimental procedure. Blood gas analyses were performed at regular intervals during the experiment (Rapidlap 348; Bayer Diagnostics, Fernwald, Germany), and ventilation parameters were adapted if necessary. Body temperature was kept constant between 37° and 38°C by means of a feedback-controlled heating blanket.

Lumbar segments L4 and L5 of the spinal cord were exposed by laminectomy, and the dura mater was incised longitudinally. All exposed tissue was covered with warm agar (1.6%) except the recording segments of the spinal cord column.

The left sciatic nerve was dissected free for bipolar electrical stimulation with a silver hook electrode. This electrode was separated from the surrounding muscles by a sheet of elastic plastic film (Parafilm, Chicago, IL). All exposed nervous tissues were then covered with warm paraffin oil.

Electrophysiologic Recordings and Nerve Stimulation

In response to electrical stimulation of the sciatic nerve, field potentials were recorded with glass microelectrodes (3–5 M Ω) 300–600 μm from the dorsal sur-

face of the spinal cord dorsal horn (laminae I and II). Microelectrodes were pulled on a Flaming/Brown micropipette puller (Sutter Instrument Co., Novato, CA) and moved into the spinal cord by an electronically controlled microstepper. Recordings were made with an ISO-DAM-amplifier (World Precision Instruments Inc., Sarasota, FL). Corner frequencies of the band-pass filter were set to 1 kHz and 0.1 Hz, respectively. The signal output was monitored on a digital oscilloscope (Hameg HM 250, Oceanside, CA) and digitized at a sampling rate of 5 kHz by an A/D-converter card (DT281-F-16SE). Data were stored in a personal computer for off-line analysis. Test stimuli (10–25 V, 0.5 ms) were applied at intervals of 5 min to the sciatic nerve. For induction of LTP, high-frequency stimulation (HFS) consisting of four trains of 100-Hz, 40- to 50-V, 0.5-ms pulses for 1 s at 10-s intervals was applied. The distance between the stimulation site at the sciatic nerve and the recording site in the lumbar spinal dorsal horn was 8 cm.

Drug Administration

Fentanyl, atropin, 8.4% sodium hydrogen carbonate, naloxone, and pancuronium were given intravenously. Heparin was given intraarterially, and 5% glucose was injected subcutaneously. All stock solutions were diluted in 0.9% saline. Volumes for intravenous injection were 200 μl . The total continuous injection volume per hour (intravenous plus intraarterial) was 0.8% per body weight.

Experimental Procedures

Test stimuli (10–25 V, 0.5 ms) were applied to the sciatic nerve for 60 min at intervals of 5 min. The following experimental protocols were performed:

1. To establish LTP under different anesthetics ($n = 13$ urethane; $n = 26$ isoflurane; $n = 7$ sevoflurane), test stimuli were given to achieve baseline recordings for 60 min as mentioned above. Then, the conditioning HFS was applied to the sciatic nerve, and responses to test stimuli were recorded for another 60 min.
2. To test the effect of the MOR agonist fentanyl on the magnitude of C fiber-evoked potentials (acute nociception), increasing doses of fentanyl were given intravenously, which in all cases consisted of a bolus followed by an infusion. Baseline recordings were performed for 60 min as mentioned above. Then, the bolus was given, and the infusion was started. Recordings were continued for another 60 min. The following dose regimens (one per animal) were used:
 - fentanyl bolus of 5 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 6 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 7$)
 - fentanyl bolus of 10 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 12 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 7$)
 - fentanyl bolus of 20 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 24 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$)

- fentanyl bolus of 40 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 48 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$)
 - fentanyl bolus of 40 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 96 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$)
 - fentanyl bolus of 40 $\mu\text{g}/\text{kg}$ followed by fentanyl infusion of 192 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 6$)
 - bolus injections of 200 μl saline followed by saline infusion of 0.2 ml/h to serve as control ($n = 6$)
3. To test the effect of these different doses of fentanyl on the induction of LTP (preemptive effect), HFS was applied to the sciatic nerve under the above-mentioned running fentanyl infusions. After HFS, fentanyl infusions were continued for another 60 min, during which responses of test stimuli were recorded.
 4. Naloxone was used as the opioid receptor antagonist to test whether the effect of fentanyl on C fiber-evoked potentials depends on the opioid receptor. Baseline recordings were performed for 60 min, and then a fentanyl bolus of 40 $\mu\text{g}/\text{kg}$ was applied, which was followed by a fentanyl infusion of 96 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ for 120 min. Then, 100 $\mu\text{g}/\text{kg}$ naloxone was given intravenously three times in 15-min intervals, and recordings were continued for another 60 min ($n = 3$).

Data Analysis and Statistics. The area under the curve of the C fiber-evoked field potentials was determined off-line by parameter extraction, which was implemented by the software Experimenter's Workbench (DataWave Technologies, Longmont, CO). The area of the C fiber-evoked field potentials was determined as the integral of the waveform. Measurements were normalized for each rat such that the mean of the respective baseline values was equal to 100. Changes of C fiber-evoked potentials on acute nociception were characterized by post minus baseline differences (values 30–60 min before start of fentanyl infusion minus values 30–60 min after start of fentanyl infusion). These changes within each dose were compared with the change in the zero dose using analysis of variance and *post hoc* tests according to Dunnett. This procedure controls the type I error for comparisons of all doses to the zero dose within each experiment. The grand mean and its corresponding 95% confidence interval (CI) were estimated to test whether there was an effect of fentanyl on C fiber-evoked potentials or of HFS on C fiber-evoked potentials under fentanyl infusion.

The same model was applied to test the changes after application of HFS during fentanyl infusion. Changes of C fiber-evoked potentials in experiments in which HFS was given during fentanyl infusion were characterized by postcomparison minus precomparison (values 30–60 min before HFS compared with values 30–60 min after application of HFS). In addition, the corresponding 95% CIs were defined for each dose respectively. Calculations were performed using the SAS software system version 8.2 (SAS Institute Inc., Cary, NC). A significant difference

was considered when P was less than 0.05. All values are expressed as mean \pm SEM.

Results

Recording of C Fiber-evoked Potentials and LTP of C Fiber-evoked Potentials under Inhalational Anesthesia

A β and C fiber-evoked field potentials could clearly be distinguished after stimulation of the sciatic nerve (fig. 1A, *inset*). C fiber-evoked field potentials that were characterized by a negative focus in superficial spinal dorsal horn, long latencies (90–120 ms, corresponding to conduction velocities < 1.2 m/s), and high thresholds (10–25 V) were investigated further.

High-frequency stimulation consistently induced LTP in urethane-anesthetized rats to $149 \pm 12\%$ of control ($n = 13$), in isoflurane-anesthetized rats to $152 \pm 6\%$ of control ($n = 26$), and in sevoflurane-anesthetized rats to $132 \pm 6\%$ of control ($n = 7$) (figs. 1A–C).

C fiber-evoked potentials remained unchanged when isoflurane was increased for 20 min from 1 vol% inspiratory to 2 vol% inspiratory and decreased back to 1 vol% inspiratory in four animals (data not shown).

Fentanyl Decreases C Fiber-evoked Potentials

The MOR agonist fentanyl reduced C fiber-evoked potentials in a dose-dependent manner: At its lowest dose (bolus 5 $\mu\text{g}/\text{kg}$, infusion 6 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 7$), fentanyl had no effect on C fiber-evoked potentials (fig. 2A). Increasing doses of fentanyl led to significant reductions of C fiber-evoked potentials (figs. 2B–F and fig. 3). At its highest dose (bolus 40 $\mu\text{g}/\text{kg}$, infusion 192 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 6$), fentanyl reduced responses to 20% of control 60 min after the start of the fentanyl injection regimen (fig. 2F).

The reduction due to fentanyl was fully reversible. Naloxone (100 $\mu\text{g}/\text{kg}$) that was given 2 h after the start of the fentanyl infusion (bolus 40 $\mu\text{g}/\text{kg}$, infusion 96 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 3$) completely antagonized the reduction in C fiber-evoked potentials by fentanyl (data not shown).

Differential Effects of Increasing Doses of Fentanyl on LTP Induction

One hour after the beginning of fentanyl infusion regimens, HFS was applied to the sciatic nerve to investigate the effect of fentanyl on the induction of LTP. There was a significant overall (intercept) increase of the C fiber-evoked potentials of 26.5% (95% CI: 16.8, 36.1). The mean changes of the separate doses were as follows:

- saline infusion ($n = 6$): mean change, 50.7 (95% CI: 25.7, 75.7)
- fentanyl infusion of 6 $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 7$): mean change, 33.3 (95% CI: 10.2, 56.5)




Fig. 1. Induction of long-term potentiation of synaptic strength between primary afferent C fibers and neurons in superficial spinal dorsal horn by high-frequency stimulation (HFS) of sciatic nerve fibers at C-fiber strength. In each experiment, C fiber-evoked potentials were provoked by electrical stimulation of the sciatic nerve. Responses were normalized to the area of C fiber-evoked potentials recorded 60 min before HFS was applied. Mean areas \pm SEMs were expressed as percent of controls and were plotted *versus* time. HFS resulted in long-term potentiation during urethane anesthesia, isoflurane, and sevoflurane. Original recordings from field potentials immediately before and 30 min after HFS during isoflurane anesthesia are shown above the graphs.^{1,2}

- fentanyl infusion of $48 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$): mean change, 28.3 (95% CI: 1, 55.7)
- fentanyl infusion of $96 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$): mean change, 35.3 (95% CI: 8, 62.8)
- fentanyl infusion of $192 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 6$): mean change, 18.8 (95% CI: 6, 43.8)

A CI that overlaps 0 reveals a significant difference, and a CI that does not overlap 0 reveals no significant difference. Therefore, the lowest dose of fentanyl (bolus $5 \mu\text{g}/\text{kg}$, infusion $6 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 7$), which revealed no effect on C fiber-evoked potentials, also failed to block the induction of LTP (fig. 3). The next higher dosages of fentanyl used (bolus $10 \mu\text{g}/\text{kg}$, infusion $12 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 7$, and bolus $20 \mu\text{g}/\text{kg}$, infusion $24 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 5$) blocked the induction of LTP (fig. 3). Even higher fentanyl dosages (bolus $40 \mu\text{g}/\text{kg}$, infusion 48 – $192 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$, $n = 5$ or $n = 6$, respectively) failed to block the induction of LTP (fig. 3). Although these high doses of fentanyl did not block the induction of LTP, the potentiated C fiber-evoked potentials all clearly remained below the control values before the start of fentanyl infusion (figs. 2D–F and fig. 3).

Discussion

The main result of this study is a U-shaped dose-response relation of the MOR agonist fentanyl on the induction of synaptic LTP in nociceptive pathways. At a very low dose, fentanyl was not able to block C fiber-evoked field potentials or prevent the induction of LTP of synaptic strength on C fibers. At intermediate doses, fentanyl blocked C fiber-evoked responses and LTP induction. At the highest doses tested, fentanyl showed an increasing antinociceptive effect but did not block the induction of LTP.

Induction of LTP under Different Anesthetics

In our experiments, surgical levels of anesthesia by volatile anesthetic isoflurane in combination with nitrous oxide failed to prevent synaptic LTP in nociceptive pathways, a form of central sensitization. There is dis-

- fentanyl infusion of $12 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 7$): mean change, 7.4 (95% CI: -15.8 , 30.5)
- fentanyl infusion of $24 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ ($n = 5$): mean change, 11.4 (95% CI: -16 , 38.8)




Fig. 2. Effects of increasing doses of fentanyl on the C fiber-evoked field potentials and on the induction of long-term potentiation. One hour after the start of fentanyl infusion, high-frequency stimulation (HFS) was given. Responses were normalized to the area of C fiber-evoked potentials recorded 60 min before HFS was applied. Mean responses \pm SEMs were expressed as percent of controls and were plotted versus time.

cussion about the exact mechanisms of action of volatile anesthetics. Suppression of neuronal electrical activity by volatile anesthetics has been ascribed to interactions with the A type of the γ -aminobutyric acid (GABA_A) receptor and glycine receptors.⁹ After binding of GABA to the GABA_A receptor, the channel opens to allow chloride ions to diffuse across the postsynaptic membrane. Volatile anesthetics prolong channel opening of GABA_A and glycine receptors and consequently increase postsynaptic inhibition.⁹ Isoflurane blocks the induction of LTP in the hippocampus *in vitro*¹⁰ and suppresses somatosensory information transfer to the thalamic level *in vivo* by enhancing thalamic GABA_A receptor-mediated inhibition.¹¹ The major action of volatile anesthetics seems to be supraspinal because isoflurane has a variable and minimal depressant effect on dorsal horn cell responses to noxious mechanical stimulation.¹² In our experiments, an increase of isoflurane concentration from 1 to 2 vol% did not influence nociceptive synaptic transmission in the spinal dorsal horn. Furthermore, nei-

ther isoflurane nor sevoflurane nor nitrous oxide prevented the induction of LTP. Nitrous oxide may have some potency as an NMDA receptor antagonist,¹³ but in our experiments, nitrous oxide at the concentration used did not prevent the induction of LTP, which requires the activation of NMDA receptors.^{14,15}

Differential Effects of Fentanyl on the Induction of LTP

The activation of opioid receptors results in at least three actions on synaptic transmission: presynaptic inhibition of Ca²⁺ channels, postsynaptically inhibition of adenylyl cyclase, and membrane hyperpolarization *via* activation of K⁺ channels (see Ingram⁸ for review). These mechanisms may synergistically act to reduce the C fiber-evoked potentials by fentanyl in the current study. None of the doses of fentanyl tested blocked the input onto the dorsal horn neurons completely. This is in accord with studies that have shown that opioids reduce but do not abolish noxious evoked




Fig. 3. Dose-dependent reduction of C fiber-evoked potentials by fentanyl (black bars). Bars show means of C fiber-evoked potentials \pm SEMs *** $P < 0.001$; values 30–60 min before start of fentanyl infusion regimen vs. values 30–60 min after start of infusion regimen; $n = 5-7$. Gray bars show mean C fiber-evoked potentials 30–60 min after high-frequency stimulation (values 30–60 min before high-frequency stimulation during fentanyl infusion vs. values 30–60 min after high-frequency stimulation during fentanyl infusion; $n = 5-7$; cf. fig. 2).

release of substance P^{16,17} and glutamate¹⁸ in the dorsal horn of the spinal cord.

At the lowest dose used, fentanyl revealed neither antinociceptive nor preemptive action. Higher doses of fentanyl, however, significantly depressed C fiber-evoked potentials, proving their potent antinociceptive action. This confirms the spinal action of opioids in animal models of nociception^{19–21} and provides a synaptic mechanism of action as strong analgesics in the clinical situation.

The induction of LTP was fully blocked during the infusion of doses of fentanyl that were not fully antinociceptive. This suggests that fentanyl is more potent to block central sensitization than nociception. Because LTP induction requires release of glutamate and activation of postsynaptic NMDA receptor, the presynaptic and postsynaptic inhibition by fentanyl well explains block of LTP induction. Our results are in line with an *in vitro* study that shows that the MOR agonist DAMGO inhibited NMDA receptor-dependent LTP in spinal cord slices of neonatal rats.²²

However, at higher doses, fentanyl did not prevent LTP induction. This phenomenon may be explained by the interactions between postsynaptic NMDA receptor and postsynaptic opioid receptors.²³ Fentanyl may lead to enhanced NMDA receptor-mediated currents in trigeminal dorsal horn neurons²⁴ and in nucleus accumbens neurons.²⁵ In both studies, the enhancement of NMDA currents by μ opioids was mediated by protein kinase C, an intracellular enzyme that can be activated by intracellular Ca^{2+} or phospholipase C pathway.^{24,25} Activation of protein kinase C reduces the Mg^{2+} block of NMDA receptor and thereby considerably facilitates currents through NMDA receptor.²⁶ Removal or significant reduction of voltage-dependent Mg^{2+} block by NMDA

receptor may explain how strongly reduced glutamate release and reduced postsynaptic depolarization may still be sufficient to induce NMDA receptor-dependent LTP under high doses of fentanyl in the current study.

There are animal studies that show that acute tolerance can occur during anesthesia²⁷ and even after a single dose of opioids.²⁸ In a clinical trial, high doses of fentanyl led to higher postoperative pain intensity and greater fentanyl consumption.²⁹ Whether the failure of high doses of fentanyl in our study to block LTP contributes to acute opioid tolerance remains to be determined.^{28,30}

It is important to note that at all doses that did not block LTP C fiber-evoked responses always remained below the control value. We can only speculate which level of C fiber-evoked potentials would be reached after the washout of fentanyl. Fentanyl accumulates as a result of its slow release by liver metabolism, and redistribution is less effective in removing fentanyl from its site of action, such as the central nervous system. Because the time course of complete washout of the lipophilic fentanyl in the rat spinal cord *in vivo* is not known but likely may last for many hours, it was not tested here.

Taken together, fentanyl in dosages with low analgesic potency completely blocks LTP but does not block LTP in higher doses.

The authors thank Magister Stephan Lehr (Institut für Medizinische Computereissenschaften, Abteilung für Klinische Biometrie, Universität Wien, Wien, Austria) for his valuable help in preparing the statistical analysis.

References

1. Crombie IK, Davies HT, Macrae WA: Cut and thrust: Antecedent surgery and trauma among patients attending a chronic pain clinic. *Pain* 1998; 76:167–71

2. Reeh PW, Pethö G: Nociceptor excitation by thermal sensitization: A hypothesis, Nervous System Plasticity of Chronic Pain. Progress in Brain Research. Vol 219. Edited by Sandkühler J, Bromm B, Gebhart GF. Amsterdam, Elsevier, 2000, pp 39-50
3. Woolf CJ: Evidence for a central component of post injury pain hypersensitivity. *Nature* 1983; 306:686-8
4. Sandkühler J: Learning and memory in pain pathways. *Pain* 2000; 88:113-8
5. Sandkühler J, Benrath J, Brechtel C, Ruscheweyh R, Heinke B: Synaptic mechanisms of hyperalgesia, Nervous System Plasticity and Chronic Pain. Progress in Brain Research. Vol 129. Edited by Sandkühler J, Bromm B, Gebhart GF. Amsterdam, Elsevier, 2000, pp 81-100
6. Ikeda H, Heinke B, Ruscheweyh R, Sandkühler J: Synaptic plasticity in spinal lamina I projection neurons that mediate hyperalgesia. *Science* 2003; 299:1237-40
7. Schroeder JE, McCleskey EW: Inhibition of Ca^{2+} currents by a μ -opioid in a defined subset of rat sensory neurons. *J Neurosci* 1993; 13:867-73
8. Ingram SL: Cellular and molecular mechanisms of opioid action. Progress in Brain Research. Vol 129. Edited by Sandkühler J, Bromm B, Gebhart GF. Amsterdam, Elsevier, 2000, pp 483-92
9. Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harris RA, Harrison NL: Sites of alcohol and volatile anaesthetic action on GABA(A) and glycine receptors. *Nature* 1997; 389:385-9
10. Simon W, Hapfelmeier G, Kochs E, Zieglgänsberger W, Rammes G: Isoflurane blocks synaptic plasticity in the mouse hippocampus. *ANESTHESIOLOGY* 2001; 94:1058-65
11. Vahle-Hinz C, Detsch O, Siemers M, Kochs E, Bromm B: Local GABA(A) receptor blockade reverses isoflurane's suppressive effects on thalamic neurons in vivo. *Anesth Analg* 2001; 92:1578-84
12. Antognini JF, Carstens E: Increasing isoflurane from 0.9 to 1.1 minimum alveolar concentration minimally affects dorsal horn cell responses to noxious stimulation. *ANESTHESIOLOGY* 1999; 90:208-14
13. Jevtovic-Todorovic V, Todorovic SM, Mennerick S, Powell S, Dikranian K, Benshoff N, Zorumski CF, Olney JW: Nitrous oxide (laughing gas) is an NMDA antagonist, neuroprotectant and neurotoxin. *Nature Med* 1998; 4:460-3
14. Liu X-G, Sandkühler J: Long-term potentiation of C-fiber-evoked potentials in the rat spinal dorsal horn is prevented by spinal N-methyl-D-aspartic acid receptor blockage. *Neurosci Lett* 1995; 191:43-6
15. Liu X-G, Sandkühler J: Activation of spinal N-methyl-D-aspartate or neurokinin receptors induces long-term potentiation of spinal C-fibre-evoked potentials. *Neuroscience* 1998; 86:1209-16
16. Hirota N, Kuraishi Y, Hino Y, Sato Y, Satoh M, Takagi H: Met-enkephalin and morphine but not dynorphin inhibit noxious stimuli induced release of substance P from rabbit dorsal horn in situ. *Neuropharmacol* 1985; 24:567-70
17. Aimone LD, Yaksh TL: Opioid modulation of capsaicin-evoked release of substance P from rat spinal cord in vivo. *Peptides* 1989;10: 1127-31
18. Kangrga I, Randic M: Outflow of endogenous aspartate and glutamate from the rat spinal dorsal horn in vitro by activation of low- and high-threshold primary afferent fibers: Modulation by μ -opioids. *Brain Res* 1991; 553:347-52
19. Kumeta Y, Murata K, Kitahata LM, Aoki M, Nishio Y, Collins JG: Fentanyl suppression of nociceptive neurons in the superficial dorsal horn of the cat. *ANESTHESIOLOGY* 1988; 69:371-6
20. Serrao JM, Stubbs SC, Goodchild CS, Gent JP: Intrathecal midazolam and fentanyl in the rat: Evidence for different spinal antinociceptive effects. *ANESTHESIOLOGY* 1989; 70:780-6
21. Kozlowski CM, Bountra C, Grundy D: The effect of fentanyl, DNQX and MK-801 on dorsal horn neurones responsive to colorectal distension in the anaesthetized rat. *Neurogastroenterol Motil* 2000; 12:239-47
22. Terman GW, Eastman CL, Chavkin C: Mu opiates inhibit long-term potentiation induction in the spinal cord slice. *J Neurophysiol* 2001; 85:485-94
23. Mao J: NMDA and opioid receptors: Their interactions in antinociception, tolerance and neuroplasticity. *Brain Res Reviews* 1999; 30:289-304
24. Chen L, Huang L-YM: Sustained potentiation of NMDA receptor-mediated glutamate responses through activation of protein kinase C by a μ opioid. *Neuron* 1991; 7:319-26
25. Martin G, Nie Z, Siggins GR: μ -Opioid receptors modulate NMDA receptor-mediated responses in nucleus accumbens neurons. *J Neurosci* 1997; 17:11-22
26. Chen L, Huang L-YM: Protein kinase C reduces Mg^{2+} block of NMDA-receptor channels as a mechanism of modulation. *Nature* 1992; 356:521-3
27. Askitopoulou H, Whitwam JG, Al-Khudhairi D, Chakrabarti M, Bower S, Hull CJ: Acute tolerance to fentanyl during anesthesia in dogs. *ANESTHESIOLOGY* 1985; 63:255-61
28. Larcher A, Laulin JP, Celerier E, Le Moal M, Simonnet G: Acute tolerance associated with a single opiate administration: Involvement of N-methyl-D-aspartate-dependent pain facilitatory systems. *Neuroscience* 1998; 84:583-9
29. Chia YY, Liu K, Wang JJ, Kuo MC, Ho ST: Intraoperative high dose fentanyl induces postoperative fentanyl tolerance. *Can J Anaesth* 1999; 46:872-7
30. Fairbanks CA, Wilcox GL: Spinal plasticity of acute opioid tolerance. *J Biomed Sci* 2000; 7:200-12