- 16. Kucukemre F, Kunt N, Kaygusuz K, Kiliccioglu F, Gurelik B, Cetin A: Remifentanil compared with morphine for postoperative patient-controlled analgesia after major abdominal surgery: A randomized controlled trial. Eur J Anaesthesiol 2005; 22:378–85
- 17. Lee LH, Irwin MG, Lui SK: Intraoperative remifentanil infusion does not increase postoperative opioid consumption compared with 70% nitrous oxide. ANESTHESIOLOGY 2005; 102:398-402
- 18. Gomez de Segura IA, Criado AB, Santos M, Tendillo FJ: Aspirin synergistically potentiates isoflurane minimum alveolar concentration reduction produced by morphine in the rat. Anesthesiology 1998; 89:1489-94
- 19. Eger El 2nd, Saidman LJ, Brandstater B: Minimum alveolar anesthetic concentration: A standard of anesthetic potency. Anesthesiology 1965; 26:756-63
- 20. Criado AB, Gomez de Segura IA: Reduction of isoflurane MAC by fentanyl or remifentanil in rats. Vet Anaesth Analg 2003; 30:250-6
- 21. Crawford MW, Hickey C, Zaarour C, Howard A, Naser B: Development of acute opioid tolerance during infusion of remifentanil for pediatric scoliosis surgery. Anesth Analg 2006; 102:1662-7
  - 22. Eger EI 2nd, Johnson BH, Weiskopf RB, Holmes MA, Yasuda N, Targ A,

- Rampil IJ: Minimum alveolar concentration of I-653 and isoflurane in pigs: Definition of a supramaximal stimulus. Anesth Analg 1988; 67:1174-6
- 23. Katoh T, Ikeda K: The minimum alveolar concentration (MAC) of sevoflurane in humans. Anisthesiology 1987: 66:301-3
- 24. Criado AB, Gómez de Segura IA, Tendillo FJ, Marsico F: Reduction of isoflurane MAC with buprenorphine and morphine in rats. Lab Anim 2000; 34:252-9
- 25. Lindstedt SL, Schaeffer PJ: Use of allometry in predicting anatomical and physiological parameters of mammals. Lab Anim 2002; 36:1-19
- 26. Lang E, Kapila A, Shlugman D, Hoke JF, Sebel PS, Glass PS: Reduction of isoflurane minimal alveolar concentration by remifentanil. Anesthesiology 1996; 85:721-8
- 27. Castanelli DJ, Splinter WM, Clavel NA: Remifentanil decreases sevoflurane requirements in children. Can J Anaesth 2005; 52:1064-70
- 28. Gustorff B, Nahlik G, Hoerauf KH, Kress HG: The absence of acute tolerance during remifentanil infusion in volunteers. Anesth Analg 2002; 94: 1223-8

## ANESTHESIOLOGY REFLECTIONS

## The Edison Etherizer



To electrify North America, Thomas A. Edison (1847-1931) proposed using direct current (D.C.) rather than the alternating current (A.C.) suggested by his rival, George Westinghouse, Jr. (1846-1914). To undermine acceptance of A.C. for household use, Edison terminally "Westinghoused" test animals in 1887 and then advocated similar use of A.C. upon death-row inmates. When the State of New York tried Edison's "Westinghouse [electric] chair" in 1888, the first victim survived a 17-second electrocution before succumbing to a 72-second one. This debacle and D.C.'s economic costs backfired on Edison, and America adopted his rival's A.C. Over a half century after the botched electrocution, Thomas A. Edison, Inc., of New Jersey manufactured an apparatus for passing "multitudinous air streams" through an ether-adsorbing "channeled carbon mass" whose heat-conducting container was surrounded by a "crystallizable liquid." Heats of "adsorption and solidification," designed into this apparatus by 1946, were counteracting the chill of vaporizing ether, the inefficiency of which had plagued earlier bubble-through vaporizers. After inscribing the signature of "Thomas A. Edison" on the front of their "Edison Etherizer" (pictured above from the Wood Library-Museum Gallery), the New Jersey team powered it with A.C. Quite an ironic posthumous salute to one-time D.C.-advocate, Thomas Alva Edison! – (Copyright © the American Society of Anesthesiologists, Inc. This image appears in color in the Anesthesiology Reflections online collection available at www.anesthesiology.org.)

George S. Bause, M.D., M.P.H., Honorary Curator, ASA's Wood Library-Museum of Anesthesiology, Park Ridge, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Obio. UJYC@aol.com.