

5. Davies CSL: The youth and education of Christopher Wren. *Engl Hist Rev* 2008; 123:300-27
6. Bennet JA: A Study of parenthalia, with two unpublished letters of Sir Christopher Wren. *Ann Sci* 1973; 30:129-47
7. Bergman NA: Early intravenous anesthesia: An eyewitness account. *ANESTHESIOLOGY* 1990; 72:185-6

8. Frank RG Jr: John Aubrey, F.R.S., John Lydall, and science at Commonwealth Oxford. *Notes Rec R Soc Lond* 1973; 27:193-217
9. McKie D: The origins and foundation of the Royal Society of London. *Notes Rec R Soc Lond* 1960; 15:1-37

(Accepted for publication June 24, 2009.)

Anesthesiology 2009; 111:924

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

In Reply:—I thank Professor Jorge Dagnino, M.D., for his chronology of the founding of the Royal Society. With my six-sentence limitation on caption space for *Anesthesiology Reflections*, I thought that I had dealt reasonably with the Royal Society's nebulous origins by writing that Wren, Boyle, and others had met "by" (not "first met") in November of 1660.

Just as I acknowledged 21 yr ago, Wren was the "brains" behind the intravenous goose quill experiment of 1656.¹ So I concur with Professor Dagnino on these facts. In "Boyle, a Most Skeptical Chemist," the 1659 date in the caption was my typographical error.²

My thanks to Professor Dagnino for his thoughtful feedback on my telegraphic captions.

George S. Bause, M.D., M.P.H., Case Western Reserve University, Cleveland, Ohio. ujyc@aol.com

References

1. Bause GS: An historical backdrop to cardiovascular monitoring. *Anesth Clin N Am* 1988; 6:672
2. Bause GS: Boyle, a most skeptical chemist. *ANESTHESIOLOGY* 2009; 110:610

(Accepted for publication June 24, 2009.)

Anesthesiology 2009; 111:924-5

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Esmolol May Abolish Volatile Anesthetic-induced Postconditioning by Scavenging Reactive Oxygen Species

To the Editor:—We read with great interest the article recently published by Lange *et al.*¹ In an *in vivo* rabbit model of myocardial ischemia-reperfusion induced by 30 min of coronary occlusion and 180 min reperfusion, the authors observed that β -adrenergic receptor blockade during early reperfusion with either the β 1-adrenergic blocker esmolol or the β 2-adrenergic blocker ICI 118,551 abolished desflurane-induced postconditioning cardioprotection manifested as reduced myocardial infarct size. However, neither esmolol nor ICI 118,551 had a significant effect on postischemic myocardial infarct size when used alone during the first 30 min of early reperfusion, in the absence of desflurane. This is a very interesting finding. However, the more interesting point of the study, as commented on by Dr. Riess in an editorial² accompanying this article, is that sustained β 1-blockade with esmolol during the entire period of reperfusion not only failed to abolish desflurane-induced postconditioning cardioprotection, but instead actually conferred a similar degree of cardioprotection. We want to join Dr. Riess² in congratulating the authors for this comprehensive study detailing the role of β -blockers in volatile anesthetic postconditioning. However, we do not entirely agree that the energy-sparing effect of β -blockade, mainly heart rate reduction, may have been the principal reason for the infarct size reduction. We propose that β -blockers may have conferred cardioprotection primarily by reducing the production of reactive oxygen species (ROS)^{3,4} during reperfusion, and that esmolol may have abolished desflurane-induced postconditioning by scavenging ROS.

ROS has been shown to play an essential role in β -adrenergic signaling in cardiac myocytes.⁵ Volatile anesthetic-induced generation of small amounts of ROS plays a critical role in anesthetic preconditioning,^{6,7} and likely in anesthetic postconditioning as well, since they share similar mechanisms. Esmolol has been shown to increase antioxidant activity and reduce ROS-induced lipid peroxidation in patients with acute myocardial infarction.⁸ Therefore, it is reasonable to postulate that esmolol abolished desflurane-induced postconditioning *via* its antioxidant action in the study of Lange *et al.*¹ If this is the case, it could be possible that the cardioprotection conferred by a combination of desflurane postconditioning and delayed β -adrenergic blockade

during reperfusion could be superior to desflurane postconditioning or β -adrenergic blockade alone. We are interested in the authors' opinion on this possibility, and the clinical relevance of their findings.

It should be noted that the volatile anesthetic isoflurane-induced ROS production and anesthetic preconditioning cardioprotection is attenuated in senescent hearts,⁹ likely because ROS production is already increased in the senescent. Information regarding the age or body weight of the study animal (New Zealand White rabbits) is not provided in the study of Lange *et al.*¹ Presumably, the study was conducted in young animals. It would be also interesting if the authors could provide this information and comment on the potential effect of aging on the effectiveness of anesthetic postconditioning.

Zhengyuan Xia, Ph.D., M.D.,* Michael G. Irwin, M.D. *The University of Hong Kong, Hong Kong, China. zyxia@hku.hk

References

1. Lange M, Redel A, Lotz C, Smul TM, Blomeyer C, Frank A, Stumpner J, Roewer N, Kehl F: Desflurane-induced postconditioning is mediated by beta-adrenergic signaling: Role of beta 1- and beta 2-adrenergic receptors, protein kinase A, and calcium/calmodulin-dependent protein kinase II. *ANESTHESIOLOGY* 2009; 110:516-28
2. Riess ML: The rocky road from bench to bedside: Beta-blockers and anesthetic postconditioning. *ANESTHESIOLOGY* 2009; 110:451-2
3. Röth E, Matos G, Guarnieri C, Papp B, Varga J: Influence of the beta-blocker therapy on neutrophil superoxide generation and platelet aggregation in experimental myocardial ischemia and reflow. *Acta Physiol Hung* 1995; 83:163-70
4. Röth E, Török B: Effect of the ultrashort-acting beta-blocker Brevibloc on free-radical-mediated injuries during the early reperfusion state. *Basic Res Cardiol* 1991; 86:422-33
5. Remondino A, Kwon SH, Communal C, Pimentel DR, Sawyer DB, Singh K, Colucci WS: Beta-adrenergic receptor-stimulated apoptosis in cardiac myocytes is mediated by reactive oxygen species/c-Jun NH2-terminal kinase-dependent activation of the mitochondrial pathway. *Circ Res* 2003; 92:136-8
6. Kevin LG, Novalija E, Riess ML, Camara AK, Rhodes SS, Stowe DF: Sevoflurane exposure generates superoxide but leads to decreased superoxide during ischemia and reperfusion in isolated hearts. *Anesth Analg* 2003; 96:949-55
7. Tanaka K, Weihsrauch D, Kehl F, Ludwig LM, LaDisa JF Jr, Kersten JR, Pagel PS, Warltier DC: Mechanism of preconditioning by isoflurane in rabbits: A direct role for reactive oxygen species. *ANESTHESIOLOGY* 2002; 97:1485-90

8. Daga MK, Chaudhary M, Sharma B, Bhattacharjee J, Ghambhir DS, Arora N, Sudha R: Effect of esmolol on oxidant status and antioxidant activity in acute myocardial infarction. *J Assoc Physicians India* 2003; 51:677-80

9. Nguyen LT, Rebecchi MJ, Moore LC, Glass PS, Brink PR, Liu L: Attenuation

of isoflurane-induced preconditioning and reactive oxygen species production in the senescent rat heart. *Anesth Analg* 2008; 107:776-82

(Accepted for publication June 29, 2009.)

Anesthesiology 2009; 111:925

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

In Reply:—We thank Drs. Xia and Irwin for their interest in our study on the role of β -adrenergic signaling in anesthetic postconditioning¹ and in the accompanying editorial view.² We agree with Drs. Xia and Irwin that, besides their energy-sparing effect, several alternative mechanisms of β blockers might be responsible for their infarct size-reducing capacity. Apart from their effect on the interaction between β receptor activation and reactive oxygen species production³ and scavenging,⁴ β blockers can inhibit calcium/calmodulin-dependent protein kinase II⁵ and phospholipase A,⁶ exert membrane stabilizing effects,⁷ and may even have direct effects on mitochondrial electron transport and reactive oxygen species production.⁸ The role of alternative mechanisms is certainly supported by the finding that infarct size reduction by β blockade is independent of heart rate, the main determinant of myocardial oxygen consumption.⁹ It is entirely conceivable that the combination of different cardioprotective principles at different time points during reperfusion might provide additive protective effects. In this context, it is of interest that calcium/calmodulin-dependent protein kinase II is necessary for desflurane-induced postconditioning, whereas prolonged postischemic calcium/calmodulin-dependent protein kinase II blockade might attenuate adverse effects of ischemia/reperfusion injury, including remodeling.¹⁰ Thus, it might be reasonable to apply anesthetic postconditioning at the onset of reperfusion and to initiate β blockade later during reperfusion. However, further basic research and clinical studies will be necessary to determine an optimized cardioprotective approach and to identify the possible clinical consequences of these experimental findings.

The rabbits used in this study were between 8 and 12 weeks of age and weighed between 2.5 and 3.0 kg. Although cardioprotection by ischemic¹¹ and pharmacological¹² preconditioning can be attenuated or lost in senescent hearts, there is some evidence of preserved ischemic postconditioning in the aged myocardium.¹³ Thus, the impact of aging on the cardioprotective effects of β blockade, anesthetic postconditioning and their interaction with reactive oxygen species needs to be determined in future studies.

Markus Lange, M.D.,* Matthias L. Riess, M.D., Ph.D. *Klinikum der Bayerischen Julius-Maximilians-Universität, Würzburg, Germany. lange_m@klinik.uni-wuerzburg.de

References

1. Lange M, Redel A, Lotz C, Smul TM, Blomeyer C, Frank A, Stumpner J, Roewer N, Kehl F: Desflurane-induced postconditioning is mediated by beta-adrenergic signaling: role of beta 1- and beta 2-adrenergic receptors, protein kinase A, and calcium/calmodulin-dependent protein kinase II. *ANESTHESIOLOGY* 2009; 110:516-28
2. Riess ML: The rocky road from bench to bedside: Beta-blockers and anesthetic postconditioning. *ANESTHESIOLOGY* 2009; 110:451-2
3. Moniri NH, Daaka Y: Agonist-stimulated reactive oxygen species formation regulates beta2-adrenergic receptor signal transduction. *Biochem Pharmacol* 2007; 74:64-73
4. Srivastava S, Chandrasekar B, Gu Y, Luo J, Hamid T, Hill BG, Prabhu SD: Downregulation of CuZn-superoxide dismutase contributes to beta-adrenergic receptor-mediated oxidative stress in the heart. *Cardiovasc Res* 2007; 74:445-55
5. Lange M, Smul TM, Redel A, Lotz C, Jazbutyte V, Schnupp V, Roewer N, Kehl F: Differential role of calcium/calmodulin-dependent protein kinase II in desflurane-induced preconditioning and cardioprotection by metoprolol: Metoprolol blocks desflurane-induced preconditioning. *ANESTHESIOLOGY* 2008; 109:72-80
6. Trotz M, Jellison EJ, Hostetler KY: Propranolol inhibition of the neural phospholipases A of rat heart mitochondria, sarcoplasmic reticulum and cytosol. *Biochem Pharmacol* 1987; 36:4251-6
7. Boucher M, Chapuy E, Duchêne-Marullaz P: Membrane stabilizing activity and β -adrenoceptor antagonist-induced bradycardia in conscious dogs. *Eur J Pharmacol* 1992; 211:343-9
8. Kametani R, Miura T, Harada N, Shibuya M, Wang R, Tan H, Fukagawa Y, Kawamura S, Matsuzaki M: Carvedilol inhibits mitochondrial oxygen consumption and superoxide production during calcium overload in isolated heart mitochondria. *Circ J* 2006; 70:321-6
9. Wartier DC, Gross GJ, Jesmok GJ, Brooks HL, Hardman HF: Protection of ischemic myocardium: Comparison of effects of propranolol, bevantolol and N-dimethyl propranolol on infarct size following coronary artery occlusion in anesthetized dogs. *Cardiology* 1980; 66:133-46
10. Backs J, Backs T, Neef S, Kreusser MM, Lehmann LH, Patrick DM, Grueter CE, Qi X, Richardson JA, Hill JA, Katus HA, Bassel-Duby R, Maier LS, Olson EN: The delta isoform of CaM kinase II is required for pathological cardiac hypertrophy and remodeling after pressure overload. *Proc Natl Acad Sci USA* 2009; 106:2342-7
11. Fenton RA, Dickson EW, Meyer TE, Dobson JG: Aging reduces the cardioprotective effect of ischemic preconditioning in the rat heart. *J Mol Cell Cardiol* 2000; 32:1371-5
12. Sniecinski R, Liu H: Reduced efficacy of volatile anesthetic preconditioning with advanced age in isolated rat myocardium. *ANESTHESIOLOGY* 2004; 100:589-97
13. Lauzier B, Delemasure S, Debin R, Collin B, Sicard P, Acar N, Bretillon L, Joffe C, Bron A, Creuzot-Garcher C, Vergely C, Rochette L: Beneficial effects of myocardial postconditioning are associated with reduced oxidative stress in a senescent mouse model. *Transplantation* 2008; 85:1802-8

(Accepted for publication June 29, 2009.)

Anesthesiology 2009; 111:925-6

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

Goals Neither Validated Nor Met in Goal-directed Colloid versus Crystalloid Therapy

To the Editor:—Kimberger *et al.*¹ and the editors² are to be commended for attempting to shed light on an important topic: What is the optimal intraoperative fluid and resuscitation target?

Many experienced physicians, including us, who provide anesthesia for major intraabdominal surgery have evolved over time from crystal-

loid-only, "show me the proof" physicians to those being in philosophical agreement with both the author and the editorial writers—goal-directed therapy with colloid is best in intestinal cases. We believe this produces less gut edema without compromising gut or other critical organ perfusion (not to mention reducing the anesthesiologist's aural discomfort from the oft repeated surgeon lament that the anesthesia team is "drowning" the patient). Indeed, Victor Hugo once said, "All the forces in the world are not so powerful as an idea whose time has come."³

Unfortunately, despite our hope to the contrary, all the forces in the world may have to wait a little longer, because this study does not

The above letter was sent to the authors of the referenced Editorial. The authors did not feel that a response was required. —James C. Eisenach, M.D., Editor-in-Chief.

*Available at http://famouspoetsandpoems.com/poets/victor_hugo/quotes. Accessed May 29, 2009.