1386 CORRESPONDENCE

prospective and retrospective studies that would allow for the control of comorbidity and variations in anesthetic management, the examination of effects according to surgical procedure, the determination of effect by LD type, and more comprehensive measures of academic achievement, cognitive/memory functions, and quality of life. This study represents an initial attempt at unraveling this complex and difficult issue. Other studies planned and currently under way will, no doubt, add to the slowly accumulating body of clinical data that we hope will help to resolve this important and difficult issue.

Randall P. Flick, M.D., M.P.H., Robert T. Wilder, M.D., Ph.D., Juraj Sprung, M.D., Ph.D.,* Slavica K. Katusic, M.D., Robert Voigt, M.D., Robert Colligan, M.D., Darrell R. Schroeder, M.S., Amy L. Weaver, M.S., David O. Warner, M.D. *Mayo Clinic, Rochester, Minnesota. sprung.juraj@mayo.edu

References

- Wilder RT, Flick RP, Sprung J, Katusic SK, Barbaresi WJ, Mickelson C, Gleich SJ, Schroeder DR, Weaver AL, Warner DO: Early exposure to anesthesia and learning disabilities in a population-based birth cohort. Anesthesiology 2009; 110:796-804
- Loepke AW, McGowan FX Jr, Soriano SG: Con: The toxic effects of anesthetics in the developing brain—The clinical perspective. Anesth Analg 2008; 106:1664-9
- 3. Jevtovic-Todorovic V, Olney JW: Pro: Anesthesia-induced developmental neuroapoptosis—Status of the evidence. Anesth Analg 2008; 106:1659-63

- Sun L: Anesthesia and neurodevelopment in children: Time for an answer. Anesthesiology 2008; 109:5
- 5. Soriano SG, Anand KJ, Rovnaghi CR, Hickey PR: Of mice and men: Should we extrapolate rodent experimental data to the care of human neonates? (letter). Anesthesiology 2005; 102:866-8, reply 868-9
- 6. Olney JW, Young C, Wozniak DF, Ikonomidou C, Jevtovic-Todorovic V: Anesthesia-induced developmental neuroapoptosis. Does it happen in humans? Anesthesiology 2004; 101:273–5
- 7. Hansen TG, Danish Registry Study Group, Flick R, Mayo Clinic Pediatric Anesthesia and Learning Disabilities Study Group: Anesthetic effects on the developing brain: Insights from epidemiology. Anesthesiology 2009; 110:1-3
- 8. Mitchell RB, Kelly J: Behavior, neurocognition and quality-of-life in children with sleep-disordered breathing. Int J Pediatr Otorhinolaryngol 2006; 70:395–406
- 9. Mitchell RB: Adenotonsillectomy for obstructive sleep apnea in children: Outcome evaluated by pre- and postoperative polysomnography. Laryngoscope 2007; 117:1844-54
- 10. Hutton JB: Incidence of learning problems among children with middle ear pathology. J Learn Disabil 1984; 17:41-2
- 11. Campbell TF, Dollaghan CA, Rockette HE, Paradise JL, Feldman HM, Shriberg LD, Sabo DLL, Kurs-Lasky M: Risk factors for speech delay of unknown origin in 3-year-old children. Child Dev 2003; 74:346-57
- 12. Cote CJ, Rolf N, Liu LM, Goudsouzian NG, Ryan JF, Zaslavsky A, Gore R, Todres TD, Vassallo S, Polaner D, Alifimoff JK: A single-blind study of combined pulse oximetry and capnography in children. Anesthesiology 1991; 74:980-7
- 13. Deulofeut R, Critz A, Adams-Chapman I, Sola A: Avoiding hyperoxia in infants < or = 1250 g is associated with improved short- and long-term outcomes. J Perinatol 2006: 26:700-5
- 14. Tin W, Milligan DW, Pennefather P, Hey E: Pulse oximetry, severe retinopathy, and outcome at one year in babies of less than 28 weeks gestation. Arch Dis Child Fetal Neonatal Ed 2001; 84:F106-10

(Accepted for publication July 31, 2009.)

Anesthesiology 2009; 111:1386

Copyright © 2009, the American Society of Anesthesiologists, Inc. Lippincott Williams & Wilkins, Inc.

The Need for Perspective

To the Editor:-We were disappointed that Anesthesiology chose to publish the articles by Kalkman et al.1 and Wilder et al.2 without an accompanying cautionary editorial. Kalkman et al.1 state, "children undergoing urologic surgery at age less than 24 months showed more behavioral disturbances . . . although the results were not statistically significant." We disagree with this statement; namely, because statistical significance was not achieved, more behavioral disturbances were not observed. Furthermore, they go on to perform a sample size calculation to determine the number of patients that would be required to detect a statistically significant effect of the effect size they found. Their estimate for such a potential association between anesthesia and behavioral problems could be explained by chance alone, and using such an estimate to guide future studies is misleading. Wilder et al.² were unable to separate out the effects of multiple anesthetics from the effects of the underlying clinical problems requiring multiple procedures. By publishing these two studies as part of a larger series including several animal models, Anesthesiology seems to send the message that two independent teams reported similar findings in humans. At a minimum, a cautionary editorial putting these studies into context was warranted. Studies such as these, reported on by the lay media, may cause an already wary public much alarm and put pediatric anesthesiologists in an impossible position. Parental concerns regarding the possible deleterious effects of anesthesia will not be assuaged by statistical explanations. Anesthesiology has an obligation beyond merely reporting interesting studies. We are sure that, like us, other readers are looking for perspective.

Karthik Raghunathan, M.D., M.P.H.,* Donald A. Schwartz, M.D., Neil Roy Connelly, M.D. *Tufts University School of Medicine, Baystate Medical Center, Springfield, Massachusetts. karthik.raghunathan@bhs.org

References

- Wilder RT, Flick RP, Sprung J, Katusic SK, Barbaresi WJ, Mickelson C, Gleich SJ, Schroeder DR, Weaver AL, Warner DO: Early exposure to anesthesia and learning disabilities in a population-based birth cohort. Anesthesiology 2009; 110:796–804
- 2. Kalkman CJ, Peelen L, Moons KG, Veenhuizen M, Bruens M, Sinnema G, de Jong TP: Behavior and development in children and age at the time of first anesthetic exposure. Anesthesiology 2009; 110:805–12

(Accepted for publication August 7, 2009.)

Anesthesiology 2009; 111:1386-7

 $Copyright @\ 2009, the\ American\ Society\ of\ Anesthesiologists,\ Inc.\ Lippincott\ Williams\ \&\ Wilkins,\ Inc.\ Lippincott\ Williams\ \&\ Wilkins,\ Lippincott\ Williams\ Wilkins,\ Wilkins,\ Lippincott\ Williams\ Wilkins,\ Wilkins,\ Lippincott\ Williams\ Wilkins,\ Wilki$

In Reply:—We thank Dr. Raghunathan et al. for their letter regarding their disappointment that we did not publish a cautionary editorial regarding the reports by Wilder et al.¹ and Kalkman et al.² in the April issue of Anesthesiology. These clinical articles, which were published with laboratory work presented at the Anesthesiology/Foundation for Anesthesia Education and Research session at the 2008 Annual Meeting of the American Society of Anesthesiologists, were accompanied by an editorial by Drs. Patel and Sun,³ thought leaders in research regarding

the mechanisms and clinical relevance of neurodevelopment after exposure to anesthetics. Regarding the clinical article, they concluded in their editorial, "Although two retrospective studies herein suggest that a correlation between anesthetic exposure early in life is associated with learning and behavioral abnormalities later in life, the data cannot be considered to be evidence of the existence of anesthetic neurotoxicity in humans. The absence of rigorously conducted prospective randomized trials precludes recommendations on clinical