

Immune Modulation by Volatile Anesthetics

Lindsay M. Stollings, M.D., Li-Jie Jia, M.D., Pei Tang, Ph.D., Huanyu Dou, M.D., Binfeng Lu, Ph.D., Yan Xu, Ph.D.

ABSTRACT

Volatile general anesthetics continue to be an important part of clinical anesthesia worldwide. The impact of volatile anesthetics on the immune system has been investigated at both mechanistic and clinical levels, but previous studies have returned conflicting findings due to varied protocols, experimental environments, and subject species. While many of these studies have focused on the immunosuppressive effects of volatile anesthetics, compelling evidence also exists for immunoactivation. Depending on the clinical conditions, immunosuppression and activation due to volatile anesthetics can be either detrimental or beneficial. This review provides a balanced perspective on the anesthetic modulation of innate and adaptive immune responses as well as indirect effectors of immunity. Potential mechanisms of immunomodulation by volatile anesthetics are also discussed. A clearer understanding of these issues will pave the way for clinical guidelines that better account for the impact of volatile anesthetics on the immune system, with the ultimate goal of improving perioperative management. (**ANESTHESIOLOGY 2016; 125:399-411**)

IN recent decades, the field of immunology has progressed substantially, elucidating many of the cellular and molecular mechanisms underlying human immune responses.^{1,2} During the same period, significant insights have been gained to better understand the action of general anesthetics.³⁻⁶ General anesthesia can be administered using inhalational anesthetics, intravenous medications, or most frequently a combination of both. All of these forms of anesthesia have been found to modulate the immune system and exert effects on innate and adaptive immunity.⁷⁻¹² A comprehensive evaluation of the impact of anesthetic regimens on the immune system can help refine current perioperative management.

This review focuses on the effects of inhalational anesthetics, specifically volatile anesthetics, on the immune response. Volatile anesthetics play a significant role in clinical anesthesia throughout the world. Furthermore, the volatile nature of these compounds extends their influence to not only the immune system of the patients, but also that of the physicians, nurses, and other personnel in the perioperative setting.¹³ Both *in vitro* and *in vivo* studies^{9,14-17} have been conducted to answer various questions regarding how volatile anesthetics impact immunity. For example, what is

the effect of anesthesia on postoperative infection? Is it the emotional and physical stress before and after the surgery or general anesthesia *per se* that predisposes patients to postoperative complications? Should the impact on immune system be part of the decision-making when choosing anesthesia regimens? Is there a benefit from immunosuppression in the ischemic setting? Do volatile anesthetics hasten the process of metastatic disease? Compared to *in vitro* and animal studies, there are fewer human studies due to the challenge in isolating a single variable in the clinical setting. Nevertheless, we sought to present a comprehensive review of *in vitro* studies and *in vivo* investigations with both small animals and humans. Figure 1 shows some of the direct immune modulations by volatile anesthetics. Direct modulation impacts innate and adaptive immunity, in which the majority of effector immune cells are natural killer (NK) cells, dendritic cells (DCs), neutrophils, macrophages, and lymphocytes. Table 1 summarizes the effects of some of the commonly used volatile anesthetics on these cells.^{7,9,18-44} In addition to the direct effects, volatile anesthetics also impact the neuroendocrine response from the hypothalamic–pituitary–adrenal axis, thereby indirectly influencing the immune response through the secretions of

This article is featured in "This Month in Anesthesiology," page 1A. Figures 1 and 2 were enhanced by Annemarie B. Johnson, C.M.I., Medical Illustrator, Vivo Visuals, Winston-Salem, North Carolina. L.M.S. and L.-J.J. equally contributed as co-first authors.

Submitted for publication September 28, 2015. Accepted for publication March 10, 2016. Departments of Anesthesiology (L.M.S., P.T., Y.X.), Immunology (B.L.), Pharmacology and Chemical Biology (P.T., Y.X.), Computational and Systems Biology (P.T.), and Structural Biology (Y.X.), University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania; Department of Anesthesiology, Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China (L.-J.J.); and Department of Biomedical Sciences, Texas Tech University Health Sciences Center, El Paso, Texas (H.D.)

Copyright © 2016, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. Anesthesiology 2016; 125:399–411




Fig. 1. Direct immune modulations by volatile anesthetics (VAs). Depicted here are immune cells responsible for the innate (*shaded*) and adaptive (*unshaded*) immunity. VAs have been shown to suppress innate immunity by impairing or suppressing neutrophil adhesion, monocytes, macrophages and natural killer (NK) cells and affecting resident cells in tissues, such as platelets and microglial cells. VAs also suppress adaptive immunity by decreasing lymphocyte proliferation, such as cluster of differentiation (CD) 4 positive (CD4⁺) and 8 positive (CD8⁺) T cells as well as B cells. Note that VAs can have both inhibition (shown as a line with a dot) and potentiation (shown as a line with an arrowhead) effects on macrophages, depending on the site of infection or inflammation. Ag = antigen; T_h1 = T helper cell type 1; T_h2 = T helper cell type 2; T_h17 = T helper cell type 17.

immunomodulator hormones such as catecholamines and glucocorticoids.⁴⁵ Both direct and indirect modulations of immunity by volatile anesthetics are covered in this review.

Innate Immunity

The innate immune response is mediated by innate immune cells that are activated when protective barriers, such as the skin or other mucosa, have been compromised due to infection or injury. Cytokines, chemokines, and inflammatory mediators are secreted by both resident tissue cells and the recruited innate immune cells. They bring forth initial responders: neutrophils, monocytes, and NK cells, as well as the complement pathways that enhance or amplify the immune responses. Surgery, sepsis, ischemia, and even the stress of being in the hospital or undergoing surgery can trigger reactions of the innate immune system. Volatile anesthetics have been found to exert a number of effects on innate immunity,^{9,18,46–48} mainly through neutrophils, DCs, NK cells, and resident tissue macrophages.

Neutrophils

Neutrophils are the most abundant granulocytes, and their numerous functions play a significant role in an inflammatory reaction.⁴⁹ They are generally the first and most lethal effector cells recruited to an inflammation site. Phagocytosis and oxidative burst, which leads to rapid production of

Table 1. Immunosuppressive and Immunoactivating Effects of Volatile Anesthetics*

Immune Cell Type	Effect	Volatile Anesthetic
Neutrophil	Decreased cell number, adhesion	Sevoflurane ^{18–22} Isoflurane ^{19–21,23} Halothane ^{18–21}
	Increase cell number	Desflurane ²⁴
PBMC/ macrophage	Decreased cytokine release (interleukin-1B, TNF- α , interleukin-6, interleukin-8, interleukin-10)	Sevoflurane ^{25,26} Isoflurane ^{25,27}
	Decreased phagocytosis, ROS, chemotaxis	Halothane ¹⁸
	Decreased cell number	Sevoflurane ²⁸ Desflurane ²⁸
	Reversed nitrous oxide immune suppression	Sevoflurane ⁷
	Enhanced nitrite production	Isoflurane ²⁷
	Increased cell number, respiratory burst	Halothane ²⁹
NK cell	Decreased cytotoxicity	Sevoflurane ^{9,30–32} Isoflurane ^{9,30–33} Halothane ^{9,30–32} Isoflurane ^{25,30,34–36} Halothane ^{25,30,34–36} Sevoflurane ³⁴
	Decreased response to interferon- γ	
	Decreased cytokine release	
T lymphocyte	Biphasic (increase then decrease cell number)	Desflurane ²⁴
	Decreased cell number, proliferation, change in T _h 1/T _h 2 ratio	Isoflurane ^{37,38}
	Decreased T _h 1	Sevoflurane ³¹
	Increased T _h 1	Desflurane ²⁴ Sevoflurane ³⁹
	Promoted cell-mediated immunity	Sevoflurane ³⁹
B lymphocyte	Decreased cell number, increased B-cell damage	Sevoflurane ^{40,41} Isoflurane ⁴⁰ Desflurane ⁴⁰
Other	Increased cortisol	Isoflurane ³⁷
	Decreased platelet-immune cell adhesion	Desflurane ⁴²
	Decreased microglial cytokine release	Isoflurane ⁴³
	Decreased monocyte chemoattractant	Isoflurane ⁴⁴ Halothane ⁴⁴
	Increased platelet-immune cell adhesion	Sevoflurane ⁴²

*Unshaded and shaded entries are considered immunosuppressive and immunoactivating, respectively.

NK = natural killer; PBMC = peripheral blood mononuclear cell; ROS = reactive oxygen species; T_h1 = T helper cell type 1; T_h2 = T helper cell type 2; TNF- α = tumor necrosis factor- α .

oxygen radicals, destroy foreign entities and damage native tissues. With this in mind, the impact of volatile anesthetics on this aspect of innate immunity can be viewed both positively and negatively.

The impaired function of neutrophils after exposure to volatile anesthetics was observed in several studies.^{18,26,47}

Sevoflurane was found to decrease the number of reacting polymorphonuclear cells (PMNs).¹⁸ Reactive oxygen species (ROS) production and chemotaxis were affected after exposure to sevoflurane, desflurane, halothane, and enflurane.¹⁸ Since halothane and enflurane are no longer used clinically, we will deemphasize their discussions in this review. Isoflurane and sevoflurane at clinical concentrations decreased neutrophil adhesion to human endothelial cells by inhibiting activation of PMNs.¹⁹ However, in the active state, PMNs are stimulated to roll and adhere to the endothelium of the vasculature within the inflamed tissue; thus, free neutrophils may not accurately reflect the active population.⁵⁰ In contrast, the suppression of neutrophil adhesion after exposure to volatile anesthetics may positively affect the deleterious effects of PMNs in the ischemic setting. Isoflurane and sevoflurane have been shown to impair the postischemic adhesion of PMNs in the intact coronary system of isolated reperfused guinea pig hearts, and their inhibitory effects may be beneficial to cardiac function during general anesthesia.^{20,21}

The results from *in vivo* studies largely parallel those from *in vitro* investigations. Exposing mice to 1.4% isoflurane before or after stimulation with lipopolysaccharide for 30 min decreased PMN levels in the bronchial alveolar fluid.²³ Neutrophils were noted to concentrate perivascularly, but were inhibited from migrating directly to the affected site. The neutrophil-attracting C-X-C motif chemokine ligand 1 (CXCL1) and CXCL2/3, which belong to the early signaling molecules for PMN recruitment in immune response, were also found to decrease in the same study. Mice injected with a sublethal dose of influenza A showed fewer physical signs and symptoms of infection after exposure to halothane.⁵¹ A delay in appearance of neutrophils in lung tissue was demonstrated. This protective effect from halogenated volatile anesthetics was shown in a recent study⁵² to be the anesthetic-induced reduction in type I and type II interferon production. Similarly, in a rat model of liver transplantation, sevoflurane was found to attenuate neutrophil renal injury and decrease neutrophil infiltration, as well as decrease plasma tumor necrosis factor- α (TNF- α) and interleukin-6 levels.⁵³ In a murine model of zymosan-induced peritonitis, isoflurane diminished the amplitude of PMN infiltration and down-regulated a panel of proinflammatory cytokines.⁵⁴ In a human study, sevoflurane at twice the minimum alveolar concentration (MAC) induced leukocyte rolling, but decreased neutrophils in the peripheral blood samples.²² Despite the aforementioned evidence of the suppressive effects of volatile anesthetics on PMNs, contradictory data exist due to the immune complexity and variation inherent in the clinical setting. A study comparing sevoflurane and propofol in combination with fentanyl noted an overall similar inflammatory response, including increased interleukin-8, decreased interleukin-17, and decreased cellular adhesion.⁵⁵ Additional research, particularly human studies, is

necessary to determine the clinical impact of volatile anesthetic effects on neutrophil function.

Macrophages

Peripheral blood mononuclear cells (PBMCs) include both lymphocytes and monocytes, which become macrophages upon migration into a tissue. Macrophages are phagocytic scavengers of innate immunity, similar to neutrophils.⁴⁹ As resident cells in the tissues, however, macrophages are often the first responders to infection, sending recruitment signals to other effector cells.

In vitro studies revealed suppressive effects of volatile anesthetics on PBMCs and macrophages. Sevoflurane and isoflurane at concentrations of 1.5 to 2.5 MAC suppressed the release of interleukin-1 β and TNF- α from human peripheral mononuclear cells stimulated by NK-sensitive tumor cells.²⁵ Despite its potent inhibitory effect on inflammatory cytokines, sevoflurane does not reduce the proliferation of human PBMC.⁷ Interestingly, the same study suggested that sevoflurane might have a beneficial effect by alleviating the immunosuppressive effect of nitrous oxide, which inhibits the proliferation of PBMC.⁷

A number of *in vivo* studies also demonstrated that volatile anesthetics could be either detrimental or beneficial, depending on the setting of inflammation with or without an infection. In a ventilated pig model, sevoflurane and desflurane were shown to decrease macrophage levels in bronchial alveolar fluid, and the overall cellular infiltration was also reduced.²⁸ Isoflurane at 1.0 MAC after lipopolysaccharide exposure decreased macrophage release of TNF- α and interleukin-1 β .²⁷ Sevoflurane decreased cytokine release, specifically interleukin-6, interleukin-8, and interleukin-10, in patients undergoing cardiac surgery.²⁶ Decreased pulmonary sequestration of white blood cells was noted as well. In contrast to suppressions caused by volatile anesthetics, a study involving isoflurane and sevoflurane administered at 1.5 MAC for more than 2 h showed significant increases in interleukin-1 β , macrophage inflammatory protein-2 (MIP-2), interferon- γ , and TNF- α in rat alveolar macrophages under mechanical ventilation.⁵⁶ Postexposure of 1.0 MAC isoflurane 4 h after lipopolysaccharide-induced endotoxemia in rats attenuated the systemic release of TNF- α and interleukin-1 β , but simultaneously enhanced the nitrite production in cultured alveolar macrophages.²⁷ Altogether, it appears that the complexity of *in vivo* studies has resulted in uncertainties regarding the relationship between macrophage function and volatile anesthetic exposure. Table 2 summarizes some of the reported effects of volatile anesthetics on several proinflammatory and antiinflammatory cytokines.^{22,24,43,52,57–109} Future in-depth mechanistic studies are needed to reconcile many of the observed discrepancies.

Natural Killer Cells

NK cells, unlike neutrophils and monocytes, are a component of innate immunity originating from the lymphoid

Table 2. General Anesthetic Effects on Several Key Pro- and Antiinflammatory Cytokines*

Anesthetics	Proinflammatory Cytokines				Antiinflammatory Cytokines			
	Interleukin-1 β	Interleukin-6	TNF- α	Interferon- γ	Interleukin-1 α	Interleukin-4	Interleukin-10	TGF- β
Isoflurane	↑57-63 ↓43,63-68	↓61,64,65	↑59-61,63 ↓62,64-67	↑62,63	—	—	↑63,69,70	↑71-73
Sevoflurane	↓74-79	↑80-82 ↓22,76,79,83-86	↑81,82,87 ↓74-79,83,84,88-90	↓91	—	No effect ⁹²⁻⁹⁴	↑74,77,87 ↓79,94	↑95,96 ↓88,97
Desflurane	↓89,98 No effect ⁶³	↑99 No effect ^{82,98}	↓89,98	↑100	↑101	No effect ²⁴	↑102 ↓100	—
Halothane	↑103	↑103	↑103,104	↑105 ↓52,106	—	↑107,108	No effect ¹⁰⁹ ↓106	↑73

*Up-arrows indicate activation or potentiation, and down-arrows indicate suppression or inhibition.

TGF = transforming growth factor; TNF = tumor necrosis factor.

lineage of white blood cells.⁴⁹ They are large granular lymphocytes that play a critical role in the defense against viral infection as well as oncologic disease. Because surgery and general anesthesia are often necessary in the treatment of cancer, extensive research has been conducted to determine the effect of volatile anesthetics on the NK cell population.

In vitro studies show that isoflurane and sevoflurane suppress NK cell cytotoxicity and cytokine-associated NK cell activation.^{9,30-32} Isoflurane decreased the NK cells' response to interferon; sevoflurane decreased cytokine release, specifically TNF- α .^{25,30,34} It was unclear whether NK cell functions could be fully restored postoperatively with supplemental interferon.^{32,110} A previous human study indicated that NK cell activity decreased for several days postoperatively.⁴⁷ A more recent *in vivo* study on dogs also showed a significant decrease in NK cytotoxic activity, measured by the percentage of NK cell-induced apoptosis and narcosis in canine thyroid adenocarcinoma cell line 24 h after isoflurane anesthesia compared to the baseline values and the control group without anesthesia.³³ The decreased responses to interferon after exposure to isoflurane were also supported by other *in vivo* studies.^{30,31,34-36} A decrease in NK cell number and a shift in cell-mediated immunity away from NK cell promotion were reported.^{31,34} A recent meta-analysis of NK cell function and anesthetic exposure in 189 patients noted significant data heterogeneity without a conclusive association between anesthetic modulation and NK cell functions and called for further clinical investigations.¹¹¹

Other Resident Tissue Cells

Resident cells in tissues, such as alveolar macrophages, platelets, and glial cells, can also be affected by volatile anesthetics, thereby affecting the immune response. Alveolar cells are in direct contact with volatile anesthetics. In rat alveolar type II cells in primary culture, isoflurane reduced cell secretions of interleukin-6, MIP-2, and monocyte chemoattractant protein-1, but did not change total protein secretion.⁴⁴ Although levels of interleukin-6 and MIP-2 were largely restored to baseline in 4 to 24 h after anesthetic exposure, monocyte chemoattractant protein-1 remained suppressed at 24 h.⁴⁴

Platelets also play a significant role in the immune response, as they are critical for cellular adhesion. After blood was incubated with 1 or 2 MAC sevoflurane for more than an hour, the binding of platelets to lymphocytes, neutrophils, and monocytes was enhanced, and the expression of P-selectin on platelets increased.⁴² However, the same treatment with desflurane resulted in a reduction in lymphocyte-platelet, neutrophil-platelet, and monocyte-platelet conjugates.⁴² Similar phenomena were also observed in a previous study.¹¹² Another independent study suggested that neither desflurane nor sevoflurane caused significant changes in adenosine diphosphate-stimulated platelets even though sevoflurane increased the expression of P-selectin in unstimulated platelets.¹¹³

Microglia, which are resident neural immune cells, were recognized recently to contribute to neuroinflammation and postoperative delirium and cognitive decline. The immune-activated microglia not only changed cell number, size, and shape but also released the proinflammatory cytokines such as TNF- α , interleukin-1 β , interleukin-6, and interferon- γ . *In vivo* experiments in young mice with repeated exposures to clinical concentrations of sevoflurane showed activation of microglia and accumulation of interleukin-6 and TNF- α , with associated cognitive impairment.⁸² The same study also found that these detrimental changes were absent in adult mice, suggesting selective vulnerability in a particular age group. Repeated exposures to desflurane did not lead to microglia activation and interleukin-6 and TNF- α accumulation in either young or old mice. In cultured H4 human neuroglioma cells, isoflurane was found to induce caspase-3 activation, cause mitochondrial dysfunction, promote ROS accumulation, induce apoptosis, and reduce cell viability.¹¹⁴⁻¹¹⁶ Strategies to target these isoflurane-induced events have been demonstrated.^{115,116}

In other studies, beneficial effects to reduce neuroinflammation are noted from preconditioning with volatile anesthetics. Isoflurane suppressed the proinflammatory cytokine interleukin-1 β in the mouse brain after intraperitoneal injection of lipopolysaccharide.⁴³ In adult mice, exposure to isoflurane was found not to produce neuroapoptosis but

reduce astroglial processes.¹¹⁷ A more recent study⁶⁴ showed that isoflurane preconditioning inhibited the up-regulation of toll-like receptor 4, which is known to regulate microglia activation and microglia production of proinflammatory factors.

Overall, because of the complexity of various tissue responses to volatile anesthetics, the anesthetic effects on innate immunity remain an active area of investigation.

Adaptive Immunity

Adaptive immune responses are distinct from innate immunity because they are generated by clonal selection of lymphocytes.¹¹⁸ There are two broad classes of adaptive immunity: humoral immune responses, which are mediated by macromolecules (such as antibodies and antimicrobial peptides) made by B lymphocytes, and cell-mediated immune responses, which are carried out mainly by T lymphocytes. Given the variety of lymphocytes and the multiple mechanisms involved in their recognition and response to antigens, investigations into the impact of volatile anesthetics on the adaptive immune system have been challenging. In general, volatile anesthetics induced a decrease in proliferation of lymphocytes or an increase in lymphocyte apoptosis.^{9,119–121}

T Lymphocytes

Cell-mediated immunity within the adaptive response includes T lymphocytes (T cells), distinguished from other lymphocytes by the T-cell receptor, which can be modified and tailored for specific antigens.⁴⁹ T-cell precursors originate in the bone marrow and then travel in an immature state to the thymus to fully mature. From that point, they circulate in the blood and throughout the secondary lymphoid tissues, such as lymph nodes, in search of antigens sequestered there by antigen-presenting cells that have traveled from infected sites. Upon activation, T cells proliferate and differentiate. T helper (T_h) cells remain in the lymph nodes. There are three subsets of T_h cells: T_h1 cells that magnify inflammation *via* soluble protein secretion and macrophage stimulation; T_h2 cells that stimulate B lymphocytes to mature and produce antibodies; and the more recently discovered T_h17 cells that produce interleukin-17, interleukin-17F, and interleukin-22 and secrete interleukin-21 to communicate with the cells in the immune system.^{122,123}

Different anesthetics have been found to produce varied effects on T_h cells^{37,100} even though exposure to volatile anesthetics has generally resulted in a decrease in the number and proliferation of T cells.^{9,47,124} It is often difficult to discern whether the decrease in the number and proliferation of T cells is related to a decrease in interferon- γ , an increase in cortisol, impaired antigen presentation, surgical insult, or a combination of all these factors. Patients exposed to isoflurane or propofol had drastically different T-cell responses.³⁷ Those exposed to isoflurane showed no change in their T_h1/T_h2 ratio, but did show an increase in cortisol, a known promoter

of T_h2 cells.³⁷ A recent clinical trial on 40 breast cancer surgeries²⁴ showed that desflurane can preserve T_h1/T_h2 ratio as well as the ratio of their cytokine products interleukin-2/interleukin-4. A separate study, however, reported a decrease in the T_h1/T_h2 ratio in patients who underwent isoflurane anesthesia.³⁸ Another study compared patients who received spinal anesthesia and desflurane general anesthesia¹⁰⁰ and showed that desflurane, but not bupivacaine, increased the T_h1/T_h2 ratio, mainly due to an increase in T_h1 responses in patients.¹⁰⁰ Exposure to sevoflurane alone was associated with a decrease in T_h1 cells and in the T_h1/T_h2 ratio postoperatively.³¹ However, adding a spinal block to sevoflurane general anesthesia in surgery was noted to preserve the T_h1/T_h2 balance and thereby reduce the promotion of tumor metastasis in a mouse tumor model.³¹ In hepatocellular carcinoma patients, neuraxial anesthesia, specifically epidural, combined with general anesthesia, was found to be superior to general anesthesia alone in promoting antitumor T_h polarization, including shifting the T_h1/T_h2 balance toward T_h1 and decreasing T_h17 .³⁹ Although further investigation is certainly needed to clarify the inconsistent effects of volatile anesthetics on the T-cell-mediated immune responses, accumulating evidence seems to suggest that a proper selection of suitable anesthetic methods can mediate the balances of T_h subsets or even benefit the balance of antitumor responses.

B Lymphocytes and Complement System

Similar to T cells, B lymphocytes (B cells) can modify their cell surface receptors or immunoglobulins to recognize specific pathogens.⁴⁹ Data regarding the effects of volatile anesthetics on B cells, however, are relatively scarce. A previous study implied that surgical trauma or associated perioperative conditions, not the specific anesthetic agent employed, was the dominant factor responsible for most postoperative specific humoral immunity impairment.⁴⁷ A more recent study, however, seemed to suggest that isoflurane, sevoflurane, and desflurane could induce B-cell damage due to calcium release from the endoplasmic reticulum.⁴⁰ A study on mice also found that sevoflurane significantly decreased the level of splenic B cells.⁴¹

Complement-mediated immunity plays a role in both innate and adaptive immunity. It can act as an extension of the immunity provided by the B cells and the antibodies that they produce. To date, there are few reports concerning the effects of volatile anesthetics on the complement system. The combination of anesthesia and surgery was recognized as being associated with a decrease in complement levels, which may represent complement pathway activation.⁴⁷ Patients exposed to halogenated volatile anesthetics developed specific IgG1 autoantibodies that were likely cleared by classical activation of the complement system, while anesthetic-induced hepatitis patients developed specific IgG4 autoantibodies that escaped clearance because of their small size or by direct inhibition of complement activation.¹²⁵

Indirect Effectors of Immunity

Volatile anesthetics can indirectly affect immunity through their impact on stress hormone levels as well as other effectors of immunity. Stress is inherent in the perioperative setting and a known modulator of the immune system. The major stress hormones include endogenous glucocorticoid (*e.g.*, cortisol in humans and corticosterone in nonhuman animals) and catecholamines (*e.g.*, epinephrine and norepinephrine), which can be released to result in systemic immune activations. Surgery-induced inflammatory response and alteration in cell-mediated immunity were found to be more pronounced after a balanced volatile anesthesia when compared to total intravenous anesthesia.¹²⁰ The effects were attributed to the enhanced stress response in patients undergoing anesthesia with a volatile agent.¹²⁰ Volatile anesthetics also often cause hypotension and transient hypoxia,^{126,127} which promote tissue inflammation and increase cellular adhesion. These can, in turn, depress the T_H1 phenotype or promote cell-mediated immunity.⁴⁶ Glycemic control in the perioperative environment is another topic of significant debate, and volatile anesthetics can exert direct effects on immunity by manipulating glucose control. Blood glucose levels were found to be higher in patients anesthetized with a combination of sevoflurane and fentanyl *versus* those anesthetized with propofol and fentanyl.⁵⁵ Isoflurane was also

noted to inhibit normal insulin production and produce a hyperglycemic response.^{128–130} The observed hyperglycemic response in isoflurane anesthesia was thought to be a consequence of both impaired glucose clearance and increased glucose production.¹²⁹

Potential Mechanisms of Immunomodulation by Volatile Anesthetics

Although specific targets of volatile anesthetics in the immune system have not been well defined, molecular and cellular events involved in immune modulation by volatile anesthetics have been identified, including a reduction in the number of immune cells due to cell death and the suppression of immune activities (fig. 2). In reality, with the heterogeneity in immune responses, immunomodulation is likely more complicated than what is shown in figure 2. For instance, cross talks may occur among different pathways, such as interactions between ROS and inducible nitric oxide synthase (iNOS). Understanding individual pathways and their relationships will facilitate mechanistic understanding of immune modulation by volatile anesthetics.

Lymphocytes are more prone to apoptosis than other immune cells.¹³¹ Apoptosis is initiated by the mitochondria-triggered pathway (intrinsic pathway) or the




Fig. 2. Potential mechanisms involved in the immunomodulation by volatile anesthetics (VAs). Depicted here are schematic representations of major immunomodulation pathways affected by VAs. The *thick solid line* shows the cytoplasmic membrane and the *dashed line* marks the nuclear membrane. The *pink shaded areas* are cytoplasmic and extracellular space, and the *light purple shaded area* is the cell nucleus. *Lines with arrowheads and dots at the end* represent "activation" and "inhibition," respectively. $\Delta\Psi_m$ = mitochondrial membrane potential; AIF = apoptosis-inducing factor; AP-1 = activator protein 1; APC = antigen-presenting cell; cGMP = cyclic 3',5'-guanosine monophosphate; DR = antigen D related; ERK = extracellular signal-regulated kinases; GTP = guanosine triphosphate; HLA = human leukocyte antigen; HO-1 = heme oxygenase 1; ICAM-1 = intercellular adhesion molecule; IL-1 = interleukin-1; IL-3 = interleukin-3; iNOS = inducible nitric oxide synthase; LFA-1 = lymphocyte function-associated antigen 1; NO = nitric oxide; PKC = protein kinase C; ROS = reactive oxygen species.

death-receptor-triggered pathway (extrinsic pathway).^{132,133} Sevoflurane and isoflurane were found to decrease mitochondrial membrane potential ($\Delta\Psi_m$) in a dose-dependent manner, subsequently triggering the release of cytochrome C from the mitochondrial intermembrane space into the cytosol and eventually inducing apoptosis *via* activation of caspase-3.^{14,121,134} The irreversible pan-caspase inhibitor Z-VAD-fmk was shown to block sevoflurane-induced apoptosis.¹⁴ Another distinct mitochondria-mediated molecule, apoptosis-inducing factor (AIF), also initiates apoptosis.^{132,133} AIF was originally identified as a mitochondrial flavoprotein that was released into the cytoplasm and subsequently entered the nucleus to cause cell death.^{132,133} A recent human study¹³⁵ showed that sevoflurane increased AIF in cardiac surgery patients, who also exhibited decreased lymphocyte counts. ROS is another major signaling molecule in the mitochondrial pathway for apoptosis.⁷⁰ Sevoflurane was shown to increase the production of intracellular ROS and promote lymphocyte apoptosis.¹³⁴ Interestingly, the same study also suggested that propofol might attenuate the sevoflurane-induced mitochondria-related apoptosis.¹³⁴ Compared to the mitochondria-triggered pathway, the death receptor-signaling pathway played little role in sevoflurane-induced lymphocyte apoptosis.¹⁴ Thus, it is reasonable to believe that mitochondria are central mediators of volatile anesthetic-associated apoptosis. In addition to apoptosis, cell necrosis could also contribute to the isoflurane-induced decrease in immune cell count.¹⁴

Adhesion molecules are important for immune cell recruitment and accumulation at inflammatory sites. The human leukocyte antigen heterodimers are cell surface antigen for the T-cell receptor. Volatile anesthetics may interact directly with these molecules to modify their functions or reduce their expression. Immune cell trafficking and penetration depend predominantly on integrin lymphocyte function-associated antigen-1 (LFA-1).¹³⁶ Isoflurane and sevoflurane bind to LFA-1 and allosterically block the coupling of LFA-1 to its major interaction partner intercellular adhesion molecule-1 found on antigen-presenting cells. As a result, immune cell adhesion is inhibited.^{16,17} It was found recently¹³⁷ that isoflurane, but not sevoflurane, had the same inhibitory effect on macrophage-1 antigen (MAC-1), a LFA-1 homologous protein. Structural biology approaches, combined with computational docking and mutations of key residues at the anesthetic binding site in LFA-1 and MAC-1, have shed new lights on how volatile anesthetic binding to a functionally important protein domain (the so-called I domain in LFA-1 and the homologous MAC-1) can allosterically change the binding pocket at a remote location on these immune signaling proteins to change their interaction with intercellular adhesion molecule-1, thereby inhibiting the downstream events of leukocyte recruitment and migration.^{137,138} Cluster of differentiation (CD) 11b (CD11b) is another pivotal integrin on the surface of leukocytes. Isoflurane and sevoflurane at clinical concentrations abolished the

upward regulation of CD11b on neutrophils and resulted in reduced neutrophil adhesion.^{19,21} L-selectin, a cell adhesion molecule belonging to the selectin family, can be found in most leukocytes. Sevoflurane decreased L-selectin expression by 25%, indicating an increased threshold for cellular activation.¹³⁹

Volatile anesthetics mostly suppress, but in some cases up-regulate, iNOS expression and nitric oxide production.^{48,140} The suppressive effect is followed by the alteration of the nitric oxide-cyclic 3',5'-guanosine monophosphate system, which is a major signaling transduction pathway implicated in a wide range of physiologic functions.^{140,141} Evidence showed that volatile anesthetics interacted with several upstream mediators of iNOS, including calcium, protein kinase C, and heme oxygenase-1 (HO-1). Isoflurane and desflurane at clinically relevant concentrations mediated the inhibitory effect on iNOS expression by inhibiting mobilization of cytosolic-free calcium, which occurred upon macrophage activation.⁴⁸ Treatment or pretreatment with 2% isoflurane induced HO-1 protein expression and caused an induction of HO activity, which correlated with a decrease in iNOS expression and nitric oxide production in lipopolysaccharide-stimulated macrophages.¹⁴² Blockade of HO activity reversed these effects.¹⁴² Pretreatment with 2% isoflurane inhibited overexpression of iNOS and accumulation of nitrite induced by lipopolysaccharide and interferon- γ in macrophages.¹⁴³ The isoflurane preconditioning effect may be mediated by isoform protein kinase C- ϵ .¹⁴³ It was noted that lipopolysaccharide stimulation in combination with interferon- γ resulted in increased nitrite release after exposure to isoflurane,⁴⁸ indicating that supplementary interferon- γ is able to overcome any inhibition of normal macrophage function. The notion was further reinforced by the study showing the reversal of volatile anesthetic-induced impairment to macrophage chemotaxis and H₂O₂ production upon addition of interferon- γ .⁴⁷

Mitogen-activated protein kinases (MAPK), including extracellular signal-regulated kinase (ERK), c-Jun *N*-terminal protein kinase, and p38 MAPK, have been implicated in proinflammatory cytokine release.¹⁴⁴ In isolated T cells, sevoflurane inhibited activation of the transcription factor activator protein-1 (AP-1), which was associated with the inhibition of p38 activity and resulted in a decreased interleukin-3 expression.¹⁵ Activation of p38 is known to regulate several inflammation-related genes, including TNF- α , interleukin- β , and interleukin-6. Isoflurane, but not halothane, can activate p38 MAPK in a concentration-dependent manner.¹⁴⁵ Most interestingly, both isoflurane and halothane can greatly enhance the proinflammatory cytokine-induced p38 activations but have little effects on oxidative stress-induced p38 activations,¹⁴⁵ suggesting that the anesthetic action might be upstream of phosphorylation of p38 MAPK. Similarly, phosphorylation of ERK can activate the transcription factor cyclic adenosine monophosphate response element-binding protein, which in turn modulates

many cyclic adenosine monophosphate response element-binding protein-targeted genes. In glial cells, particularly microglia, isoflurane suppressed lipopolysaccharide-induced phosphorylation of ERK 1/2 and the high expression of interleukin-1 β mRNA and protein but did not affect nuclear factor- κ B or activator protein-1 activation.⁴³ More molecular biology investigations aimed at dissecting anesthetic effects on each of these pathways will help us better understand the molecular mechanisms of immunomodulation by volatile anesthetics.

Additional Considerations

Oncologic Considerations

Surgical intervention remains a primary treatment for cancer. Anesthetics used during the perioperative period may influence the immune system, directly affect cancer cells, and ultimately modify oncologic outcomes.^{146,147} Investigations on whether anesthetics affect the outcome and prognosis of cancer have been carried out on various types of cancer, including ovarian, colon, breast, prostate, and rectal cancer. Several retrospective studies suggested that, in comparison to general anesthesia, regional anesthesia was associated with better outcomes.^{148–152} In one study, patients who had radical prostatectomy during general anesthesia using a combination of sevoflurane and nitrous oxide showed more than 20% higher mortality rate than those who received epidural anesthesia.¹⁴⁸ A similar study also showed a decrease in time to tumor recurrence after primary cancer surgery or even to death when using an anesthetic regimen of sevoflurane alone compared to that of intraoperative neuraxial anesthesia combined with postoperative analgesia.¹⁵³ However, conflicting data exist. The beneficial effect of regional anesthesia on cancer recurrence was not observed in all types of cancer.^{152,154,155} Some studies found no association between volatile anesthetics and death or length of cancer-free survival time.^{156,157} Thus, more prospective randomized controlled trials, with careful designs of various anesthetic regimens, are warranted. In addition, in order to optimize anesthesia management strategies for oncologic surgical patients, more mechanistic studies at the cellular level are also needed.^{7,158–161}

Positive Immune Modulation by Volatile Anesthetics

Despite the suppressive effects of volatile anesthetics on the function of neutrophils, macrophages, DCs, T cells, B cells, and NK cells, as reviewed above under the subheadings “Innate Immunity” and “Adaptive Immunity,” is it possible for volatile anesthetics to enhance the immune system or possibly generate therapeutic benefits? A recent review by Fukazawa and Lee¹⁶² presented compelling evidence of protective effects of volatile anesthetics against ischemic acute kidney injury in both preclinical and clinical studies. The cellular mechanisms of volatile anesthetic-induced kidney protection lie in the anesthetic activation of multiple

pathways synthesizing antiinflammatory and cytoprotective signaling molecules, such as releasing transforming growth factor β 1, activating CD73 and inducing interleukin-11, generating adenosine, and producing sphingosine kinase and sphingosine-1-phosphate.^{71,96,163–165} Antiischemic and antiinflammatory effects of volatile anesthetics were also shown to affect other organs such as the heart, liver, and brain through either pretreatment before prolonged ischemia or after completion of an ischemic insult.^{166–169} The protective mechanisms on these organs, however, may differ from those observed in renal protection.¹⁷⁰ Nevertheless, systematic investigations of protective effects of volatile anesthetics against acute kidney injury have led to an important discovery that volatile anesthetics can positively modulate immunity and provide off-label therapeutic effects if dose and exposure time of volatile anesthetics are optimized.¹⁶²

Closing Remarks

The majority of studies reported thus far show that volatile anesthetics have immunosuppressive effects. Whether a short period of immunosuppression has a prolonged effect on patient outcomes merits further investigation. Well-controlled randomized clinical trials are highly desirable, although isolating effects of volatile anesthetics from other factors in a perioperative setting remains challenging. Future studies should take into consideration the surgical procedures involved, the anesthetics and other medications used, and the time dependence in immune modulation and resolution. Because immunosuppression is in general detrimental for cancer patients, but potentially beneficial for septic patients, the choice of anesthesia regimens should be carefully evaluated in the overall planning for the perioperative care.

In vitro– and *in vivo*–level mechanistic studies focusing on how volatile anesthetics modulate various immune responses should continue. These studies not only can provide valuable clues to initiate more complex clinical trials, but also can identify useful biomarkers to detect the detrimental effects of volatile anesthetics.

Desirable off-label effects of volatile anesthetics have been demonstrated in a few areas, but on a large scale, they are underexplored.¹⁶² Whether and how volatile anesthetics positively modulate immune responses and subsequently generate therapeutic benefits to patients warrants further investigations.

Research Support

Supported by a National Institutes of Health (NIH; Bethesda, Maryland) Ruth Kirschstein National Service Award (to Dr. Stollings) through the NIH training grant T32GM075770 (to Dr. Xu) and by the NIH grants R01GM066358 (to Dr. Tang) and R01GM114851 (to Drs. Dou and Xu).

Competing Interests

The authors declare no competing interests.

Correspondence

Address correspondence to Dr. Xu: Department of Anesthesiology, University of Pittsburgh, 2048 Biomedical Science Tower 3, 3501 Fifth Avenue, Pittsburgh, Pennsylvania 15260. xuy@anes.upmc.edu. Information on purchasing reprints may be found at www.anesthesiology.org or on the masthead page at the beginning of this issue. ANESTHESIOLOGY's articles are made freely accessible to all readers, for personal use only, 6 months from the cover date of the issue.

References

- Satpathy AT, Wu X, Albring JC, Murphy KM: Re(de)fining the dendritic cell lineage. *Nat Immunol* 2012; 13:1145–54
- Iwasaki A, Medzhitov R: Control of adaptive immunity by the innate immune system. *Nat Immunol* 2015; 16:343–53
- Forman SA, Miller KW: Anesthetic sites and allosteric mechanisms of action on cys-loop ligand-gated ion channels. *Can J Anaesth* 2011; 58:191–205
- Franks NP: General anaesthesia: From molecular targets to neuronal pathways of sleep and arousal. *Nat Rev Neurosci* 2008; 9:370–86
- Hemmings HC Jr, Akabas MH, Goldstein PA, Trudell JR, Orser BA, Harrison NL: Emerging molecular mechanisms of general anesthetic action. *Trends Pharmacol Sci* 2005; 26:503–10
- Tang P, Xu Y: Large-scale molecular dynamics simulations of general anesthetic effects on the ion channel in the fully hydrated membrane: The implication of molecular mechanisms of general anesthesia. *Proc Natl Acad Sci USA* 2002; 99:16035–40
- Schneemilch CE, Hachenberg T, Ansorge S, Ittenson A, Bank U: Effects of different anaesthetic agents on immune cell function *in vitro*. *Eur J Anaesthesiol* 2005; 22:616–23
- Homburger JA, Meiler SE: Anesthesia drugs, immunity, and long-term outcome. *Curr Opin Anaesthesiol* 2006; 19:423–8
- Kurosawa S, Kato M: Anesthetics, immune cells, and immune responses. *J Anesth* 2008; 22:263–77
- Pirbudak Cocelli L, Ugur MG, Karadasli H: Comparison of effects of low-flow sevoflurane and desflurane anesthesia on neutrophil and T-cell populations. *Curr Ther Res Clin Exp* 2012; 73:41–51
- Gong L, Qin Q, Zhou L, Ouyang W, Li Y, Wu Y, Li Y: Effects of fentanyl anesthesia and sufentanil anesthesia on regulatory T cells frequencies. *Int J Clin Exp Pathol* 2014; 7:7708–16
- Alsina E, Matute E, Ruiz-Huerta AD, Gilsanz F: The effects of sevoflurane or remifentanyl on the stress response to surgical stimulus. *Curr Pharm Des* 2014; 20:5449–68
- Bargellini A, Rovesti S, Barbieri A, Vivoli R, Roncaglia R, Righi E, Borella P: Effects of chronic exposure to anaesthetic gases on some immune parameters. *Sci Total Environ* 2001; 270:149–56
- Loop T, Dovi-Akue D, Frick M, Roesslein M, Egger L, Humar M, Hoetzel A, Schmidt R, Borner C, Pahl HL, Geiger KK, Pannen BH: Volatile anesthetics induce caspase-dependent, mitochondria-mediated apoptosis in human T lymphocytes *in vitro*. *ANESTHESIOLOGY* 2005; 102:1147–57
- Loop T, Scheiermann P, Doviakue D, Musshoff F, Humar M, Roesslein M, Hoetzel A, Schmidt R, Madea B, Geiger KK, Pahl HL, Pannen BH: Sevoflurane inhibits phorbol-myristate-acetate-induced activator protein-1 activation in human T lymphocytes *in vitro*: Potential role of the p38-stress kinase pathway. *ANESTHESIOLOGY* 2004; 101:710–21
- Yuki K, Astrof NS, Bracken C, Soriano SG, Shimaoka M: Sevoflurane binds and allosterically blocks integrin lymphocyte function-associated antigen-1. *ANESTHESIOLOGY* 2010; 113:600–9
- Zhang H, Astrof NS, Liu JH, Wang JH, Shimaoka M: Crystal structure of isoflurane bound to integrin LFA-1 supports a unified mechanism of volatile anesthetic action in the immune and central nervous systems. *FASEB J* 2009; 23:2735–40
- Fröhlich D, Rothe G, Schwall B, Schmid P, Schmitz G, Taeger K, Hobbhahn J: Effects of volatile anaesthetics on human neutrophil oxidative response to the bacterial peptide FMLP. *Br J Anaesth* 1997; 78:718–23
- Möbert J, Zahler S, Becker BF, Conzen PF: Inhibition of neutrophil activation by volatile anesthetics decreases adhesion to cultured human endothelial cells. *ANESTHESIOLOGY* 1999; 90:1372–81
- Kowalski C, Zahler S, Becker BF, Flaucher A, Conzen PF, Gerlach E, Peter K: Halothane, isoflurane, and sevoflurane reduce postischemic adhesion of neutrophils in the coronary system. *ANESTHESIOLOGY* 1997; 86:188–95
- Heindl B, Reichle FM, Zahler S, Conzen PF, Becker BF: Sevoflurane and isoflurane protect the reperfused guinea pig heart by reducing postischemic adhesion of polymorphonuclear neutrophils. *ANESTHESIOLOGY* 1999; 91:521–30
- Morisaki H, Aoyama Y, Shimada M, Ochiai R, Takeda J: Leucocyte distribution during sevoflurane anaesthesia. *Br J Anaesth* 1998; 80:502–3
- Reutershan J, Chang D, Hayes JK, Ley K: Protective effects of isoflurane pretreatment in endotoxin-induced lung injury. *ANESTHESIOLOGY* 2006; 104:511–7
- Woo JH, Baik HJ, Kim CH, Chung RK, Kim DY, Lee GY, Chun EH: Effect of propofol and desflurane on immune cell populations in breast cancer patients: A randomized trial. *J Korean Med Sci* 2015; 30:1503–8
- Mitsuhata H, Shimizu R, Yokoyama MM: Suppressive effects of volatile anesthetics on cytokine release in human peripheral blood mononuclear cells. *Int J Immunopharmacol* 1995; 17:529–34
- Cho EJ, Yoon JH, Hong SJ, Lee SH, Sim SB: The effects of sevoflurane on systemic and pulmonary inflammatory responses after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 2009; 23:639–45
- Flondor M, Hofstetter C, Boost KA, Betz C, Homann M, Zwissler B: Isoflurane inhalation after induction of endotoxemia in rats attenuates the systemic cytokine response. *Eur Surg Res* 2008; 40:1–6
- Kalimeris K, Christodoulaki K, Karakitsos P, Batistatou A, Lekka M, Bai M, Kitsioulis E, Nakos G, Kostopanagiotou G: Influence of propofol and volatile anaesthetics on the inflammatory response in the ventilated lung. *Acta Anaesthesiol Scand* 2011; 55:740–8
- Colucci D, Harvey G, Gayol MC, Elena G, Puig N: Halothane anesthesia in mice: Effect on the phagocytic activity and respiratory burst of peritoneal macrophages. *Neuroimmunomodulation* 2011; 18:11–8
- Melamed R, Bar-Yosef S, Shakhar G, Shakhar K, Ben-Eliyahu S: Suppression of natural killer cell activity and promotion of tumor metastasis by ketamine, thiopental, and halothane, but not by propofol: Mediating mechanisms and prophylactic measures. *Anesth Analg* 2003; 97:1331–9
- Wada H, Seki S, Takahashi T, Kawarabayashi N, Higuchi H, Habu Y, Sugahara S, Kazama T: Combined spinal and general anesthesia attenuates liver metastasis by preserving TH1/TH2 cytokine balance. *ANESTHESIOLOGY* 2007; 106:499–506
- Welden B, Gates G, Mallari R, Garrett N: Effects of anesthetics and analgesics on natural killer cell activity. *AANA J* 2009; 77:287–92
- Miyata T, Kodama T, Honma R, Nezu Y, Harada Y, Yogo T, Hara Y, Tagawa M: Influence of general anesthesia with isoflurane following propofol-induction on natural killer cell cytotoxic activities of peripheral blood lymphocytes in dogs. *J Vet Med Sci* 2013; 75:917–21
- Markovic SN, Knight PR, Murasko DM: Inhibition of interferon stimulation of natural killer cell activity in mice

- anesthetized with halothane or isoflurane. *ANESTHESIOLOGY* 1993; 78:700–6
35. Sessler DI: Does regional analgesia reduce the risk of cancer recurrence? A hypothesis. *Eur J Cancer Prev* 2008; 17:269–72
 36. Brand JM, Kirchner H, Poppe C, Schmucker P: The effects of general anesthesia on human peripheral immune cell distribution and cytokine production. *Clin Immunol Immunopathol* 1997; 83:190–4
 37. Ren XF, Li WZ, Meng FY, Lin CF: Differential effects of propofol and isoflurane on the activation of T-helper cells in lung cancer patients. *Anaesthesia* 2010; 65:478–82
 38. Inada T, Yamanouchi Y, Jomura S, Sakamoto S, Takahashi M, Kambara T, Shingu K: Effect of propofol and isoflurane anaesthesia on the immune response to surgery. *Anaesthesia* 2004; 59:954–9
 39. Zhou D, Gu FM, Gao Q, Li QL, Zhou J, Miao CH: Effects of anesthetic methods on preserving anti-tumor T-helper polarization following hepatectomy. *World J Gastroenterol* 2012; 18:3089–98
 40. Yang H, Liang G, Hawkins BJ, Madesh M, Pierwola A, Wei H: Inhalational anesthetics induce cell damage by disruption of intracellular calcium homeostasis with different potencies. *ANESTHESIOLOGY* 2008; 109:243–50
 41. Puig NR, Ferrero P, Bay ML, Hidalgo G, Valenti J, Amerio N, Elena G: Effects of sevoflurane general anesthesia: Immunological studies in mice. *Int Immunopharmacol* 2002; 2:95–104
 42. Horn NA, de Rossi L, Robitzsch T, Hecker KE, Hutschenreuter G, Rossaint R: The effects of sevoflurane and desflurane *in vitro* on platelet-leukocyte adhesion in whole blood. *Anaesthesia* 2003; 58:312–9
 43. Tanaka T, Kai S, Matsuyama T, Adachi T, Fukuda K, Hirota K: General anesthetics inhibit LPS-induced IL-1 β expression in glial cells. *PLoS One* 2013; 8:e82930
 44. Giraud O, Molliex S, Rolland C, Leçon-Malas V, Desmonts JM, Aubier M, Dehoux M: Halogenated anesthetics reduce interleukin-1 β -induced cytokine secretion by rat alveolar type II cells in primary culture. *ANESTHESIOLOGY* 2003; 98:74–81
 45. Marana E, Annetta MG, Meo F, Parpaglion R, Galeone M, Maussier ML, Marana R: Sevoflurane improves the neuroendocrine stress response during laparoscopic pelvic surgery. *Can J Anaesth* 2003; 50:348–54
 46. Eltzschig HK, Carmeliet P: Hypoxia and inflammation. *N Engl J Med* 2011; 364:656–65
 47. Stevenson GW, Hall SC, Rudnick S, Seleny FL, Stevenson HC: The effect of anesthetic agents on the human immune response. *ANESTHESIOLOGY* 1990; 72:542–52
 48. Tschaikowsky K, Ritter J, Schröppel K, Kühn M: Volatile anesthetics differentially affect immunostimulated expression of inducible nitric oxide synthase: Role of intracellular calcium. *ANESTHESIOLOGY* 2000; 92:1093–102
 49. Parham P: *The Immune System*, 2nd edition. New York, Garland Science, 2005
 50. Leonard SA, Redmond HP: Effects of volatile and intravenous anesthetic agents on neutrophil function. *Int Anesthesiol Clin* 2003; 41:21–9
 51. Penna AM, Johnson KJ, Camilleri J, Knight PR: Alterations in influenza A virus specific immune injury in mice anesthetized with halothane or ketamine. *Intervirology* 1990; 31:188–96
 52. MacDonald BA, Chakravarthy KV, Davidson BA, Mullan BA, Alluri R, Hakansson AP, Knight PR: Halothane modulates the type I interferon response to influenza and minimizes the risk of secondary bacterial pneumonia through maintenance of neutrophil recruitment in an animal model. *ANESTHESIOLOGY* 2015; 123:590–602
 53. Kong HY, Zhu SM, Wang LQ, He Y, Xie HY, Zheng SS: Sevoflurane protects against acute kidney injury in a small-size liver transplantation model. *Am J Nephrol* 2010; 32:347–55
 54. Chiang N, Schwab JM, Fredman G, Kasuga K, Gelman S, Serhan CN: Anesthetics impact the resolution of inflammation. *PLoS One* 2008; 3:e1879
 55. Tylman M, Sarbinowski R, Bengtson JP, Kvarnström A, Bengtsson A: Inflammatory response in patients undergoing colorectal cancer surgery: The effect of two different anesthetic techniques. *Minerva Anesthesiol* 2011; 77:275–82
 56. Kotani N, Takahashi S, Sessler DI, Hashiba E, Kubota T, Hashimoto H, Matsuki A: Volatile anesthetics augment expression of proinflammatory cytokines in rat alveolar macrophages during mechanical ventilation. *ANESTHESIOLOGY* 1999; 91:187–97
 57. Zhang J, Tan H, Jiang W, Zuo Z: The choice of general anesthetics may not affect neuroinflammation and impairment of learning and memory after surgery in elderly rats. *J Neuroimmune Pharmacol* 2015; 10:179–89
 58. Kong FJ, Ma LL, Zhang HH, Zhou JQ: Alpha 7 nicotinic acetylcholine receptor agonist GTS-21 mitigates isoflurane-induced cognitive impairment in aged rats. *J Surg Res* 2015; 194:255–61
 59. Yang C, Zhu B, Ding J, Wang ZG: Isoflurane anesthesia aggravates cognitive impairment in streptozotocin-induced diabetic rats. *Int J Clin Exp Med* 2014; 7:903–10
 60. Li XM, Zhou MT, Wang XM, Ji MH, Zhou ZQ, Yang JJ: Resveratrol pretreatment attenuates the isoflurane-induced cognitive impairment through its anti-inflammation and -apoptosis actions in aged mice. *J Mol Neurosci* 2014; 52:286–93
 61. Wu X, Lu Y, Dong Y, Zhang G, Zhang Y, Xu Z, Culley DJ, Crosby G, Marcantonio ER, Tanzi RE, Xie Z: The inhalation anesthetic isoflurane increases levels of proinflammatory TNF- α , IL-6, and IL-1 β . *Neurobiol Aging* 2012; 33:1364–78
 62. Wang HL, Ma RH, Fang H, Xue ZG, Liao QW: Impaired spatial learning memory after isoflurane anesthesia or appendectomy in aged mice is associated with microglia activation. *J Cell Death* 2015; 8:9–19
 63. Schläpfer M, Piegeler T, Dull RO, Schwartz DE, Mao M, Bonini MG, Z'Graggen BR, Beck-Schimmer B, Minshall RD: Propofol increases morbidity and mortality in a rat model of sepsis. *Crit Care* 2015; 19:45
 64. Xiang HF, Cao DH, Yang YQ, Wang HQ, Zhu LJ, Ruan BH, Du J, Wang MC: Isoflurane protects against injury caused by deprivation of oxygen and glucose in microglia through regulation of the toll-like receptor 4 pathway. *J Mol Neurosci* 2014; 54:664–70
 65. Wang H, Wang L, Li NL, Li JT, Yu F, Zhao YL, Wang L, Yi J, Wang L, Bian JF, Chen JH, Yuan SF, Wang T, Lv YG, Liu NM, Zhu XS, Ling R, Yun J: Subanesthetic isoflurane reduces zymosan-induced inflammation in murine Kupffer cells by inhibiting ROS-activated p38 MAPK/NF- κ B signaling. *Oxid Med Cell Longev* 2014; 2014:851692
 66. Liang Y, Li Z, Mo N, Li M, Zhuang Z, Wang J, Wang Y, Guo X: Isoflurane preconditioning ameliorates renal ischemia-reperfusion injury through antiinflammatory and antiapoptotic actions in rats. *Biol Pharm Bull* 2014; 37:1599–605
 67. Altay O, Suzuki H, Hasegawa Y, Ostrowski RP, Tang J, Zhang JH: Isoflurane on brain inflammation. *Neurobiol Dis* 2014; 62:365–71
 68. Fallor S, Strosing KM, Ryter SW, Buerkle H, Loop T, Schmidt R, Hoetzel A: The volatile anesthetic isoflurane prevents ventilator-induced lung injury *via* phosphoinositide 3-kinase/Akt signaling in mice. *Anesth Analg* 2012; 114:747–56
 69. Tomihari M, Nishihara A, Shimada T, Yanagawa M, Miyoshi M, Miyahara K, Oishi A: A comparison of the immunological effects of propofol and isoflurane for maintenance of anesthesia in healthy dogs. *J Vet Med Sci* 2015; 77:1227–33
 70. Canakçi CF, Çiçek Y, Canakçi V: Reactive oxygen species and human inflammatory periodontal diseases. *Biochemistry (Mosc)* 2005; 70:619–28

71. Kim M, Ham A, Kim JY, Brown KM, D'Agati VD, Lee HT: The volatile anesthetic isoflurane induces ecto-5'-nucleotidase (CD73) to protect against renal ischemia and reperfusion injury. *Kidney Int* 2013; 84:90–103
72. Kim M, Park SW, Kim M, D'Agati VD, Lee HT: Isoflurane post-conditioning protects against intestinal ischemia-reperfusion injury and multiorgan dysfunction *via* transforming growth factor- β 1 generation. *Ann Surg* 2012; 255:492–503
73. Lee HT, Kim M, Kim J, Kim N, Emala CW: TGF- β 1 release by volatile anesthetics mediates protection against renal proximal tubule cell necrosis. *Am J Nephrol* 2007; 27:416–24
74. Rancan L, Huerta L, Cusati G, Erquicia I, Isea J, Paredes SD, García C, Garutti I, Simón C, Vara E: Sevoflurane prevents liver inflammatory response induced by lung ischemia-reperfusion. *Transplantation* 2014; 98:1151–7
75. Qiao S, Xie H, Wang C, Wu X, Liu H, Liu C: Delayed anesthetic preconditioning protects against myocardial infarction *via* activation of nuclear factor- κ B and upregulation of autophagy. *J Anesth* 2013; 27:251–60
76. Ferrando C, Aguilar G, Piqueras L, Soro M, Moreno J, Belda FJ: Sevoflurane, but not propofol, reduces the lung inflammatory response and improves oxygenation in an acute respiratory distress syndrome model: A randomised laboratory study. *Eur J Anaesthesiol* 2013; 30:455–63
77. Zhang Y, Zhang FG, Meng C, Tian SY, Wang YX, Zhao W, Chen J, Zhang XS, Liang Y, Zhang SD, Xing YJ: Inhibition of sevoflurane preconditioning against cerebral ischemia reperfusion-induced oxidative injury in rats. *Molecules* 2012; 17:341–54
78. Bedirli N, Bagriacik EU, Emmez H, Yilmaz G, Unal Y, Ozkose Z: Sevoflurane and isoflurane preconditioning provides neuroprotection by inhibition of apoptosis-related mRNA expression in a rat model of focal cerebral ischemia. *J Neurosurg Anesthesiol* 2012; 24:336–44
79. Zhang Y, Tian SY, Li YW, Zhang L, Yu JB, Li J, Chen YY, Wang YX, Liang Y, Zhang XS, Wang WS, Liu HG: Sevoflurane preconditioning improving cerebral focal ischemia-reperfusion damage in a rat model *via* PI3K/Akt signaling pathway. *Gene* 2015; 569:60–5
80. Sun Z, Satomoto M, Makita K: Therapeutic effects of intravenous administration of bone marrow stromal cells on sevoflurane-induced neuronal apoptosis and neuroinflammation in neonatal rats. *Korean J Anesthesiol* 2015; 68:397–401
81. Qiao Y, Feng H, Zhao T, Yan H, Zhang H, Zhao X: Postoperative cognitive dysfunction after inhalational anesthesia in elderly patients undergoing major surgery: The influence of anesthetic technique, cerebral injury and systemic inflammation. *BMC Anesthesiol* 2015; 15:154
82. Shen X, Dong Y, Xu Z, Wang H, Miao C, Soriano SG, Sun D, Baxter MG, Zhang Y, Xie Z: Selective anesthesia-induced neuroinflammation in developing mouse brain and cognitive impairment. *ANESTHESIOLOGY* 2013; 118:502–15
83. Romero A, Moreno A, García J, Sánchez C, Santos M, García J: Effects of sevoflurane on ventilator induced lung injury in a healthy lung experimental model. *Rev Esp Anesthesiol Reanim* 2016; 63:22–8
84. Otsuki T, Ishikawa M, Hori Y, Goto G, Sakamoto A: Volatile anesthetic sevoflurane ameliorates endotoxin-induced acute lung injury *via* microRNA modulation in rats. *Biomed Rep* 2015; 3:408–12
85. Luo C, Yuan D, Zhao W, Chen H, Luo G, Su G, Hei Z: Sevoflurane ameliorates intestinal ischemia-reperfusion-induced lung injury by inhibiting the synergistic action between mast cell activation and oxidative stress. *Mol Med Rep* 2015; 12:1082–90
86. Yu QB, Li HM, Li LL, Wang SY, Wu YB: Sevoflurane downregulates interleukin-6 and interleukin-8 levels in patients after cardiopulmonary bypass surgery: A meta-analysis. *Genet Mol Res* 2015; 14:19016–27
87. Shen QY, Fang L, Wu HM, Wu L, He F, Liu RY: [Effect of Toll-like receptor 2 on the inhibition role of sevoflurane on airway inflammation in asthmatic mice]. *Zhonghua Yi Xue Za Zhi* 2016; 96:138–41
88. Burburan SM, Silva JD, Abreu SC, Samary CS, Guimarães IH, Xisto DG, Morales MM, Rocco PR: Effects of inhalational anaesthetics in experimental allergic asthma. *Anaesthesia* 2014; 69:573–82
89. Schilling T, Kozian A, Senturk M, Huth C, Reinhold A, Hedenstierna G, Hachenberg T: Effects of volatile and intravenous anesthesia on the alveolar and systemic inflammatory response in thoracic surgical patients. *ANESTHESIOLOGY* 2011; 115:65–74
90. Lai J, Zhang L, Wang H, Lin P, Chen W: [Effects of sevoflurane preconditioning on cardiomyocyte apoptosis and myocardial inflammation in rats with sepsis]. *Nan Fang Yi Ke Da Xue Xue Bao* 2014; 34:1680–3
91. Polak PE, Dull RO, Kalinin S, Sharp AJ, Ripper R, Weinberg G, Schwartz DE, Rubinstein I, Feinstein DL: Sevoflurane reduces clinical disease in a mouse model of multiple sclerosis. *J Neuroinflammation* 2012; 9:272
92. Cata JP, Bauer M, Sokari T, Ramirez MF, Mason D, Plautz G, Kurz A: Effects of surgery, general anesthesia, and perioperative epidural analgesia on the immune function of patients with non-small cell lung cancer. *J Clin Anesth* 2013; 25:255–62
93. Kvarnström AL, Sarbinowski RT, Bengtson JP, Jacobsson LM, Bengtsson AL: Complement activation and interleukin response in major abdominal surgery. *Scand J Immunol* 2012; 75:510–6
94. Deegan CA, Murray D, Doran P, Moriarty DC, Sessler DI, Mascha E, Kavanagh BP, Buggy DJ: Anesthetic technique and the cytokine and matrix metalloproteinase response to primary breast cancer surgery. *Reg Anesth Pain Med* 2010; 35:490–5
95. Lee HT, Kim M, Song JH, Chen SW, Gubitosa G, Emala CW: Sevoflurane-mediated TGF- β 1 signaling in renal proximal tubule cells. *Am J Physiol Renal Physiol* 2008; 294:F371–8
96. Lee HT, Chen SW, Doetschman TC, Deng C, D'Agati VD, Kim M: Sevoflurane protects against renal ischemia and reperfusion injury in mice *via* the transforming growth factor- β 1 pathway. *Am J Physiol Renal Physiol* 2008; 295:F128–36
97. Lee HJ, Kwon JY, Shin SW, Baek SH, Choi KU, Jeon YH, Kim WS, Bae JH, Choi HJ, Kim HK, Baik SW: Effects of sevoflurane on collagen production and growth factor expression in rats with an excision wound. *Acta Anaesthesiol Scand* 2010; 54:885–93
98. Boost KA, Hofstetter C, Flondor M, Betz C, Homann M, Pfeilschifter J, Muehl H, Zwissler B: Desflurane differentially affects the release of proinflammatory cytokines in plasma and bronchoalveolar fluid of endotoxemic rats. *Int J Mol Med* 2006; 17:1139–44
99. Baki ED, Aldemir M, Kokulu S, Koca HB, Ela Y, Sivaci RG, Ozturk NK, Emmiler M, Adali F, Uzel H: Comparison of the effects of desflurane and propofol anesthesia on the inflammatory response and s100 β protein during coronary artery bypass grafting. *Inflammation* 2013; 36: 1327–33
100. Koksoy S, Sahin Z, Karsli B: Comparison of the effects of desflurane and bupivacaine on Th1 and Th2 responses. *Clin Lab* 2013; 59:1215–20
101. Sheen MJ, Yang CP, Liu YC, Borel CO, Wong CS, Ho ST, Wu CT: Comparing the effects of minimal low-flow desflurane with that of semi-close high flow desflurane on perioperative cytokine response in patients undergoing gastrectomy. *Acta Anaesthesiol Taiwan* 2006; 44:5–10
102. Sun Z, Lv J, Zhu Y, Song D, Zhu B, Miao C: Desflurane preconditioning protects human umbilical vein endothelial cells against anoxia/reoxygenation by upregulating NLRP12 and inhibiting non-canonical nuclear factor- κ B signaling. *Int J Mol Med* 2015; 36:1327–34

103. You Q, Cheng L, Reilly TP, Wegmann D, Ju C: Role of neutrophils in a mouse model of halothane-induced liver injury. *Hepatology* 2006; 44:1421–31
104. Dugan CM, MacDonald AE, Roth RA, Ganey PE: A mouse model of severe halothane hepatitis based on human risk factors. *J Pharmacol Exp Ther* 2010; 333:364–72
105. Dugan CM, Fullerton AM, Roth RA, Ganey PE: Natural killer cells mediate severe liver injury in a murine model of halothane hepatitis. *Toxicol Sci* 2011; 120:507–18
106. Kobayashi E, Kobayashi M, Tsuneyama K, Fukami T, Nakajima M, Yokoi T: Halothane-induced liver injury is mediated by interleukin-17 in mice. *Toxicol Sci* 2009; 111:302–10
107. Chakraborty M, Fullerton AM, Semple K, Chea LS, Proctor WR, Bourdi M, Kleiner DE, Zeng X, Ryan PM, Dagur PK, Berkson JD, Reilly TP, Pohl LR: Drug-induced allergic hepatitis develops in mice when myeloid-derived suppressor cells are depleted prior to halothane treatment. *Hepatology* 2015; 62:546–57
108. Proctor WR, Chakraborty M, Fullerton AM, Korrapati MC, Ryan PM, Semple K, Morrison JC, Berkson JD, Chea LS, Yang Q, Li AP, Spolski R, West EE, Rochman Y, Leonard WJ, Bourdi M, Pohl LR: Thymic stromal lymphopoietin and interleukin-4 mediate the pathogenesis of halothane-induced liver injury in mice. *Hepatology* 2014; 60:1741–52
109. Simeonova GP, Slavov E, Usunov R, Halacheva K, Dinev DN: Increased apoptosis of peripheral blood mononuclear cells (PBMC) during general and epidural anaesthesia in dogs. *Vet Res Commun* 2008; 32:619–26
110. Kutza J, Gratz I, Afshar M, Murasko DM: The effects of general anesthesia and surgery on basal and interferon stimulated natural killer cell activity of humans. *Anesth Analg* 1997; 85:918–23
111. Conrick-Martin I, Kell MR, Buggy DJ: Meta-analysis of the effect of central neuraxial regional anesthesia compared with general anesthesia on postoperative natural killer T lymphocyte function. *J Clin Anesth* 2012; 24:3–7
112. Fröhlich D, Rothe G, Schmitz G, Hansen E: Volatile anaesthetics induce changes in the expression of P-selectin and glycoprotein Ib on the surface of platelets *in vitro*. *Eur J Anaesthesiol* 1998; 15:641–8
113. Koroglu A, Cicek M, Toprak HI, Karakoc Y, Noyan F, Ersoy OM: Comparison of the effects of desflurane and sevoflurane on the expression of platelet surface glycoproteins in unstimulated and adenosine diphosphate-induced platelets *in vitro*. *J Clin Anesth* 2007; 19:328–33
114. Zhang Y, Zhen Y, Dong Y, Xu Z, Yue Y, Golde TE, Tanzi RE, Moir RD, Xie Z: Anesthetic propofol attenuates the isoflurane-induced caspase-3 activation and A β oligomerization. *PLoS One* 2011; 6:e27019
115. Sun Y, Zhang Y, Cheng B, Dong Y, Pan C, Li T, Xie Z: Glucose may attenuate isoflurane-induced caspase-3 activation in H4 human neuroglioma cells. *Anesth Analg* 2014; 119:1373–80
116. Cheng B, Zhang Y, Wang A, Dong Y, Xie Z: Vitamin C attenuates isoflurane-induced caspase-3 activation and cognitive impairment. *Mol Neurobiol* 2015; 52:1580–9
117. Dallasen RM, Bowman JD, Xu Y: Isoflurane does not cause neuroapoptosis but reduces astroglial processes in young adult mice. *Med Gas Res* 2011; 1:27
118. Janeway C: *Immunobiology: The immune system in health and disease*, 6th edition. New York, Garland Science, 2005
119. Picq CA, Clarençon D, Sinniger VE, Bonaz BL, Mayol JF: Impact of anesthetics on immune functions in a rat model of vagus nerve stimulation. *PLoS One* 2013; 8:e67086
120. Schneemilch CE, Itenson A, Ansorge S, Hachenberg T, Bank U: Effect of 2 anesthetic techniques on the postoperative proinflammatory and anti-inflammatory cytokine response and cellular immune function to minor surgery. *J Clin Anesth* 2005; 17:517–27
121. Matsuoka H, Kurosawa S, Horinouchi T, Kato M, Hashimoto Y: Inhalation anesthetics induce apoptosis in normal peripheral lymphocytes *in vitro*. *ANESTHESIOLOGY* 2001; 95:1467–72
122. Korn T, Bettelli E, Oukka M, Kuchroo VK: IL-17 and Th17 cells. *Annu Rev Immunol* 2009; 27:485–517
123. Miossec P, Korn T, Kuchroo VK: Interleukin-17 and type 17 helper T cells. *N Engl J Med* 2009; 361:888–98
124. Hamra JG, Yaksh TL: Halothane inhibits T cell proliferation and interleukin-2 receptor expression in rats. *Immunopharmacol Immunotoxicol* 1996; 18:323–36
125. Njoku DB, Mellerson JL, Talor MV, Kerr DR, Faraday NR, Outschoorn I, Rose NR: Role of CYP2E1 immunoglobulin G4 subclass antibodies and complement in pathogenesis of idiosyncratic drug-induced hepatitis. *Clin Vaccine Immunol* 2006; 13:258–65
126. Rossi A, Falzetti G, Donati A, Orsetti G, Pelaia P: Desflurane *versus* sevoflurane to reduce blood loss in maxillofacial surgery. *J Oral Maxillofac Surg* 2010; 68:1007–12
127. Dal D, Celiker V, Ozer E, Başgöl E, Salman MA, Aypar U: Induced hypotension for tympanoplasty: A comparison of desflurane, isoflurane and sevoflurane. *Eur J Anaesthesiol* 2004; 21:902–6
128. van den Brom CE, Bulte CS, Loer SA, Bouwman RA, Boer C: Diabetes, perioperative ischaemia and volatile anaesthetics: Consequences of derangements in myocardial substrate metabolism. *Cardiovasc Diabetol* 2013; 12:42
129. Lattermann R, Schricker T, Wachter U, Georgieff M, Goertz A: Understanding the mechanisms by which isoflurane modifies the hyperglycemic response to surgery. *Anesth Analg* 2001; 93:121–7
130. Desborough JP, Jones PM, Persaud SJ, Landon MJ, Howell SL: Isoflurane inhibits insulin secretion from isolated rat pancreatic islets of Langerhans. *Br J Anaesth* 1993; 71:873–6
131. Hotchkiss RS, Nicholson DW: Apoptosis and caspases regulate death and inflammation in sepsis. *Nat Rev Immunol* 2006; 6:813–22
132. Hengartner MO: The biochemistry of apoptosis. *Nature* 2000; 407:770–6
133. Jäättelä M, Tschopp J: Caspase-independent cell death in T lymphocytes. *Nat Immunol* 2003; 4:416–23
134. Zhou Y, Li E, Li Y, Liu S: Attenuating sevoflurane-induced cellular injury of human peripheral lymphocytes by propofol in a concentration-dependent manner. *Arch Pharm Res* 2011; 34:1535–43
135. Jia L, Dong R, Zhang F, Wang W, Lu H, Luo Y, Xue Q, Yu B: Propofol provides more effective protection for circulating lymphocytes than sevoflurane in patients undergoing off-pump coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth* 2015; 29:1172–9
136. Evans R, Patzak I, Svensson L, De Filippo K, Jones K, McDowall A, Hogg N: Integrins in immunity. *J Cell Sci* 2009; 122(pt 2):215–25
137. Jung S, Yuki K: Differential effects of volatile anesthetics on leukocyte integrin macrophage-1 antigen. *J Immunotoxicol* 2016; 13:148–56
138. Yuki K, Bu W, Shimaoka M, Eckenhoof R: Volatile anesthetics, not intravenous anesthetic propofol bind to and attenuate the activation of platelet receptor integrin α IIb β 3. *PLoS One* 2013; 8:e60415
139. Lucchinetti E, Aguirre J, Feng J, Zhu M, Suter M, Spahn DR, Härter L, Zaugg M: Molecular evidence of late preconditioning after sevoflurane inhalation in healthy volunteers. *Anesth Analg* 2007; 105:629–40
140. Toda N, Toda H, Hatano Y: Anesthetic modulation of immune reactions mediated by nitric oxide. *J Anesth* 2008; 22:155–62

141. Vulliamoz Y: The nitric oxide-cyclic 3',5'-guanosine monophosphate signal transduction pathway in the mechanism of action of general anesthetics. *Toxicol Lett* 1998; 100-101:103-8
142. Li QF, Zhu YS, Jiang H, Xu H, Sun Y: Heme oxygenase-1 mediates the anti-inflammatory effect of isoflurane preconditioning in LPS-stimulated macrophages. *Acta Pharmacol Sin* 2009; 30:228-34
143. Xu X, Feng J, Zuo Z: Isoflurane preconditioning reduces the rat NR8383 macrophage injury induced by lipopolysaccharide and interferon gamma. *ANESTHESIOLOGY* 2008; 108:643-50
144. Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, Berman K, Cobb MH: Mitogen-activated protein (MAP) kinase pathways: Regulation and physiological functions. *Endocr Rev* 2001; 22:153-83
145. Itoh T, Hirota K, Hisano T, Namba T, Fukuda K: The volatile anesthetics halothane and isoflurane differentially modulate proinflammatory cytokine-induced p38 mitogen-activated protein kinase activation. *J Anesth* 2004; 18:203-9
146. Xie Z: Cancer prognosis: Can anesthesia play a role? *ANESTHESIOLOGY* 2013; 119:501-3
147. Gottschalk A, Sharma S, Ford J, Durieux ME, Tiourine M: Review article: The role of the perioperative period in recurrence after cancer surgery. *Anesth Analg* 2010; 110:1636-43
148. Lin L, Liu C, Tan H, Ouyang H, Zhang Y, Zeng W: Anaesthetic technique may affect prognosis for ovarian serous adenocarcinoma: A retrospective analysis. *Br J Anaesth* 2011; 106:814-22
149. Christopherson R, James KE, Tableman M, Marshall P, Johnson FE: Long-term survival after colon cancer surgery: A variation associated with choice of anesthesia. *Anesth Analg* 2008; 107:325-32
150. Exadaktylos AK, Buggy DJ, Moriarty DC, Mascha E, Sessler DI: Can anesthetic technique for primary breast cancer surgery affect recurrence or metastasis? *ANESTHESIOLOGY* 2006; 105:660-4
151. Biki B, Mascha E, Moriarty DC, Fitzpatrick JM, Sessler DI, Buggy DJ: Anesthetic technique for radical prostatectomy surgery affects cancer recurrence: A retrospective analysis. *ANESTHESIOLOGY* 2008; 109:180-7
152. Gupta A, Björnsson A, Fredriksson M, Hallböök O, Eintrei C: Reduction in mortality after epidural anaesthesia and analgesia in patients undergoing rectal but not colonic cancer surgery: A retrospective analysis of data from 655 patients in central Sweden. *Br J Anaesth* 2011; 107:164-70
153. de Oliveira GS Jr, Ahmad S, Schink JC, Singh DK, Fitzgerald PC, McCarthy RJ: Intraoperative neuraxial anesthesia but not postoperative neuraxial analgesia is associated with increased relapse-free survival in ovarian cancer patients after primary cytoreductive surgery. *Reg Anesth Pain Med* 2011; 36:271-7
154. Gottschalk A, Ford JG, Regelin CC, You J, Mascha EJ, Sessler DI, Durieux ME, Nemergut EC: Association between epidural analgesia and cancer recurrence after colorectal cancer surgery. *ANESTHESIOLOGY* 2010; 113:27-34
155. Ismail H, Ho KM, Narayan K, Kondalsamy-Chennakesavan S: Effect of neuraxial anaesthesia on tumour progression in cervical cancer patients treated with brachytherapy: A retrospective cohort study. *Br J Anaesth* 2010; 105:145-9
156. Myles PS, Peyton P, Silbert B, Hunt J, Rigg JR, Sessler DI, ANZCA Trials Group Investigators: Perioperative epidural analgesia for major abdominal surgery for cancer and recurrence-free survival: Randomised trial. *BMJ* 2011; 342:d1491
157. Fleischmann E, Marschalek C, Schlemitz K, Dalton JE, Gruenberger T, Herbst F, Kurz A, Sessler DI: Nitrous oxide may not increase the risk of cancer recurrence after colorectal surgery: A follow-up of a randomized controlled trial. *BMC Anesthesiol* 2009; 9:1
158. Huang H, Benzonana LL, Zhao H, Watts HR, Perry NJ, Bevan C, Brown R, Ma D: Prostate cancer cell malignancy *via* modulation of HIF-1 α pathway with isoflurane and propofol alone and in combination. *Br J Cancer* 2014; 111:1338-49
159. Benzonana LL, Perry NJ, Watts HR, Yang B, Perry IA, Coombes C, Takata M, Ma D: Isoflurane, a commonly used volatile anesthetic, enhances renal cancer growth and malignant potential *via* the hypoxia-inducible factor cellular signaling pathway *in vitro*. *ANESTHESIOLOGY* 2013; 119:593-605
160. Tavares AN, Perry NJ, Benzonana LL, Takata M, Ma D: Cancer recurrence after surgery: Direct and indirect effects of anesthetic agents. *Int J Cancer* 2012; 130:1237-50
161. Buckley A, McQuaid S, Johnson P, Buggy DJ: Effect of anaesthetic technique on the natural killer cell anti-tumour activity of serum from women undergoing breast cancer surgery: A pilot study. *Br J Anaesth* 2014; 113 Suppl 1:i56-62
162. Fukazawa K, Lee HT: Volatile anesthetics and AKI: Risks, mechanisms, and a potential therapeutic window. *J Am Soc Nephrol* 2014; 25:884-92
163. Ham A, Kim M, Kim JY, Brown KM, Yeh J, D'Agati VD, Lee HT: Critical role of interleukin-11 in isoflurane-mediated protection against ischemic acute kidney injury in mice. *ANESTHESIOLOGY* 2013; 119:1389-401
164. Bakar AM, Park SW, Kim M, Lee HT: Isoflurane protects against human endothelial cell apoptosis by inducing sphingosine kinase-1 *via* ERK MAPK. *Int J Mol Sci* 2012; 13:977-93
165. Kim M, Kim M, Park SW, Pitson SM, Lee HT: Isoflurane protects human kidney proximal tubule cells against necrosis *via* sphingosine kinase and sphingosine-1-phosphate generation. *Am J Nephrol* 2010; 31:353-62
166. Pagel PS: Postconditioning by volatile anesthetics: Salvaging ischemic myocardium at reperfusion by activation of prosurvival signaling. *J Cardiothorac Vasc Anesth* 2008; 22:753-65
167. Lv X, Yang L, Tao K, Liu Y, Yang T, Chen G, Yu W, Lv H, Wu F: Isoflurane preconditioning at clinically relevant doses induce protective effects of heme oxygenase-1 on hepatic ischemia reperfusion in rats. *BMC Gastroenterol* 2011; 11:31
168. Xiong L, Zheng Y, Wu M, Hou L, Zhu Z, Zhang X, Lu Z: Preconditioning with isoflurane produces dose-dependent neuroprotection *via* activation of adenosine triphosphate-regulated potassium channels after focal cerebral ischemia in rats. *Anesth Analg* 2003; 96:233-7, table of contents
169. Warltier DC, al-Wathiqui MH, Kampine JP, Schmeling WT: Recovery of contractile function of stunned myocardium in chronically instrumented dogs is enhanced by halothane or isoflurane. *ANESTHESIOLOGY* 1988; 69:552-65
170. Lee HT, Ota-Setlik A, Fu Y, Nasr SH, Emala CW: Differential protective effects of volatile anesthetics against renal ischemia-reperfusion injury *in vivo*. *ANESTHESIOLOGY* 2004; 101:1313-24