- Griffee MJ, Tonna JE, McKellar SH, Zimmerman JM: Echocardiographic guidance and troubleshooting for venovenous extracorporeal membrane oxygenation using the dual-lumen bicaval cannula. J Cardiothorac Vasc Anesth 2018; 32:370–8
- Fierro MA, Daneshmand MA, Bartz RR: Perioperative management of the adult patient on venovenous extracorporeal membrane oxygenation requiring noncardiac surgery. ANESTHESIOLOGY 2018; 128:181–201

ANESTHESIOLOGY REFLECTIONS FROM THE WOOD LIBRARY-MUSEUM

Would Chest Compressions from 57 Rescuers Save Albert Frick from Landry's Paralysis?



Four days after undergoing two dental extractions, 22-yr-old Albert R. Frick (1904 to 1927) found himself hospitalized in a wheelchair as first his legs weakened and then his diaphragm became completely flaccid from acute ascending or Landry's paralysis, now better known as Guillain-Barré syndrome. For 4 days in St. Francis Hospital, Evanston, Illinois, Frick's employer (*far right*), Frick's father (*next to the employer*), and scores of friends and coworkers toiled in pairs to rhythmically compress and release Frick's chest. Finally, at 10:02 PM on Thursday, March 10, 1927, Albert Frick was pronounced dead. Before critical care and ventilators were commonplace, salesmen, bookkeepers, stenographers—all told, 57 rescuers—had tried to keep young Frick breathing during his 108h of respiratory paralysis. (Copyright © the American Society of Anesthesiologists' Wood Library-Museum of Anesthesiology.)

George S. Bause, M.D., M.P.H., Honorary Curator and Laureate of the History of Anesthesia, Wood Library-Museum of Anesthesiology, Schaumburg, Illinois, and Clinical Associate Professor, Case Western Reserve University, Cleveland, Ohio. UJYC@aol.com.