


# INFOGRAPHICS IN ANESTHESIOLOGY

Complex Information for Anesthesiologists Presented Quickly and Clearly



Infographic created by Jonathan P. Wanderer, Vanderbilt University Medical Center, and James P. Rathmell, Brigham and Women's Health Care/Harvard Medical School. Illustration by Annemarie Johnson, Vivo Visuals. Address correspondence to Dr. Wanderer: jon.wanderer@vumc.org.

1. Farag E, Ghobrial M, Sessler DI, Dalton JE, Liu J, Lee JH, Zaky S, Benzal E, Bingaman W, Kurz A: Effect of perioperative intravenous lidocaine administration on pain, opioid consumption, and quality of life after complex spine surgery. *ANESTHESIOLOGY* 2013; 119:932–4
2. Loftus RW, Yeager MP, Clark JA, Brown JR, Abdu WA, Sengupta DK, Beach ML: Intraoperative ketamine reduces perioperative opiate consumption in opiate-dependent patients with chronic back pain undergoing back surgery. *ANESTHESIOLOGY* 2010; 113:639–46
3. Toms L, McQuay HJ, Derry S, Moore RA: Single dose oral paracetamol (acetaminophen) for postoperative pain in adults. *Cochrane Database Syst Rev* 2008; CD004602
4. Turan A, Karamanlioglu B, Memiş D, Hamamcioglu MK, Tükenmez B, Pamukçu Z, Kurt I: Analgesic effects of gabapentin after spinal surgery. *ANESTHESIOLOGY* 2004; 100:935–8
5. Maheshwari K, Avitsian R, Sessler DI, Makarova N, Tanios M, Raza S, Traul D, Rajan S, Manlapaz M, Machado S, Krishnaney A, Machado A, Rosenquist R, Kurz A: Multimodal analgesic regimen for spine surgery: A randomized placebo-controlled trial. *ANESTHESIOLOGY* 2020; 132:992–1002