who underwent open heart operations with extracorporeal circulation disclosed the following changes in the postoperative period: mild anemia, minimal hemolysis of crythrocytes, leukocytosis, "atypical" lymphocytes, slight reticulocytosis and minimal prolongation of the prothrombin time. (Battle, J. D. and Heulett, J. S.: Hematologic Changes Observed After Extracorporeal Circulation During Open Heart Operations, Clereland Clinic Quart. 25: 112 (April) 1958.)

CARDIAC PACEMAKER To undertake open heart operations without a pacemaker at hand no longer seems justifiable. The electrodes should be placed on the heart of any patient in whom atrioventricular block occurs during the operation, even though the ventricular rate appears satisfactory at the time. The pacemaker should be used in children whose ventricular rates fall below 90, and in adults whose rates drop helow 80. (Olmsted, F., Kolff, W. J., and Effler, D. R.: Electronic Cardiac Pacemaker After Open Heart Operations, Cleveland Clinic Quart. 25: 81 (April) 1958.)

PULMONARY COMPLICATIONS Temporary overloading of the pulmonary circulation is the most important single factor in the initiation of capillary damage that marks the beginning of severe pulmonary complications after open heart operations. Overloading may occur by forward overfilling, through collateral vessels and by retrograde overfilling. Other possible factors are pre-existing pulmonary vascular disease, oxygen intoxication of alveolar and capillary cells and exsicention of the lungs. (Kolff, W. J., and others: Pulmonary Complications of Open Heart Operations: Their Pathogenesis and Avoidance, Cleveland Clinic Quart. 25: 65 (April) 1958.)

OPEN HEART SURGERY Extracorporeal circulation and hypothermia were used for open heart surgery in a series of 46 patients. Low flow extracorporeal circulation and hypothermia have proven to be complementary for open heart surgery. This procedure is supported by the high venous oxygen saturation and the minor alteration in the lactic acid levels in blood during perfusion. Difficulties on temperature control have heen solved by the use of a heat exchanger in the extence corporeal system. Cardiac irritability has not been a serious problem. (Seally, W. E., Brouen, I. W., and Young, W. G., JE; Report on Use of Both Extracorporeal Cigalitation and Hypothermia for Open Heat Surgery, Ann. Surg. 117: 603 (May) 1958.)

OPEN HEART MORTALITY With the use of extracorporeal circulation techniques, the mortality rate is now well under 5 per cent in the less serious cardiac defects. DeWall achieved a rate of 2.5 per cent in 40 consecutive cases; Spencer 4-ports thirteen consecutive nortic commissurotomics with no mortality; Lillehei Esports a mortality rate of 8 per cent in 20c last 25 consecutive patients undergoing complete correction of tetralogy of Fallet. (Heimbecker, R. O.: Heart-Lung Machige in Open Heart Surgery, Canad. M. A. Z. 73: 534 (April 1) 1938.)

AORTIC VALVE SURGERY Two techniques for aortic valve surgery under direct exposure have been devised in dogs. Both utilize a pump-oxygenator which x8turns blood to the femoral artery while the norta is clamped two inches distal to the aortic valve. To maintain myocardial 🕏 tegrity in one method, oxygenated blood is perfused through the coronary system in a retrograde fashion after inserting a casnula into the coronary sinus. The secost method utilizes the induction of cardiac standstill with potassium to prevent myo-Both methods permet cardial damage. restoration of normal unsupported circula-(State. D., affil tion in most instances. others: Direct Visualization of Aortic Valre in Dogs, West. J. Surg. 66: 29 (March-April) 1958.)

MYOCARDIAL CONTRACTILITY
The effect of cardine bypass with potassium induced arrest and right ventriculotory was investigated in fourteen dogs and tenter. Direct measurements of mygacardial contractility in these studies showed that the heart was still capable of doing the same amount of work following re-