

THE METABOLIC PRODUCTION OF CARBON DIOXIDE¹

MIDNEY FLEISCHER, PH.D.

ALL organic compounds when burned or combusted to completion yield carbon dioxide as one of the final products. Animal systems without exception contain the biochemical apparatus for burning organic compounds by molecular oxygen and thus massive CO₂ production is the hallmark of the animal body. We shall address ourselves in this review to the following questions: (1) which organic compounds can be combusted to CO₂; (2) how are these compounds formed; and (3) how does the combustion process provide energy for the needs of the body.

WHICH ORGANIC SUBSTANCES ARE COMBUSTED TO CARBON DIOXIDE?

The three main classes of organic materials in the animal body, *viz.*, proteins, fats and carbohydrates, are all ultimately combustible. That is not to say that proteins, fats and carbohydrates are combustible as such. An elaborate preliminary preparatory process is mandatory before the stage is set for the actual combustion. For example, proteins must first be hydrolyzed to peptides; in turn the peptides must be hydrolyzed to amino acids; these in turn must be degraded to particular keto acids and only these can funnel into the cellular furnace. The nature of the preliminary manipulation to which the complex molecules have to be subjected will vary from one class to another and quite separate biochemical mechanisms are involved in the degradation of each of the three main classes of organic materials. The unifying and simplifying fact is that proteins, carbohydrates and fats ultimately give rise to much the same small molecules and thus the same cellular furnace can be used to burn the fragmentation or degradation products of each of the three classes of foodstuffs.

Dr. Fleischer is Assistant Professor, Institute for Enzyme Research, University of Wisconsin, Madison, Wisconsin.

HOW PROTEINS, LIPIDS AND CARBOHYDRATES ARE PREPARED FOR COMBUSTION

In essence, the preparative process is simply a matter of breaking the bonds which hold together the component molecules of the large macromolecules. Proteins can be looked upon as large molecular weight compounds composed of as many as several hundred amino acids. In the protein the component amino acids are linked one to the other by means of

peptide bonds $\left(\begin{array}{c} \text{O} \quad \text{H} \\ \parallel \quad | \\ -\text{C}-\text{N}- \end{array} \right)$. In the gut there

are specific enzymes which hydrolyze these peptide bonds, and as a result of this proteolytic action the protein is reduced to the large number of component amino acids. Not all of the amino acids can be directly transformed into derivative molecules that are capable of entering the cellular furnace. However, almost without exception, all amino




FIG. 1. The structural formulas of compounds capable of being directly combusted by the cell: (a) pyruvic acid, (b) acetic acid, (c) citric acid, (d) isocitric acid, (e) α -ketoglutaric acid, (f) succinic acid, (g) fumaric acid, (h) malic acid and (i) oxaloacetic acid. (see figure 3, The Citric Acid Cycle.)

TABLE I
FORMATION OF CITRIC ACID CYCLE INTER-
MEDIATES AND CO₂ FROM AMINO ACIDS

Amino Acid	Key Degradation Product
alanine	} → pyruvate
serine	
cysteine	
methionine	
tyrosine	} → fumarate + acetyl CoA
phenylalanine	
glutamic acid	} → α-ketoglutarate
proline	
arginine	
histidine	
lysine	→ α-ketoglutarate + CO ₂
leucine	→ acetyl CoA + CO ₂
isoleucine	→ *propionyl CoA + acetyl CoA + CO ₂
threonine	→ *propionyl CoA + CO ₂
valine	→ *propionyl CoA + CO ₂
aspartic acid	→ oxaloacetate
glycine	→ CO ₂

* Propionyl CoA is converted to a citric acid cycle intermediate, succinic acid.

acids can be manipulated chemically in such a way that they will give rise, in part, if not in whole, to some molecule which is directly combustible. The particular molecules which are capable of direct combustion are pyruvic acid, acetic acid, citric acid, isocitric acid, α-ketoglutaric acid, succinic acid, fumaric acid, malic acid and oxaloacetic acid (structural formulas, fig. 1). As shown in table I, it is possible to convert each of 16 amino acids into at least one of the nine compounds capable of direct combustion.

Lipids are, in essence, esters of polyalcohols, such as glycerol or inositol. Both fatty acids and substituted phosphoric acids, such as choline phosphate and ethanolamine phosphate (formulas, fig. 2) form the ester linkages with the polyalcohols. The preparation of lipid for ultimate combustion involves essentially the hydrolysis of the ester bonds which hold the molecule together. After hydrolysis by a group of special enzymes known as esterases, the lipid molecule is reduced to the individual chemical building stones, namely, the free fatty acids, glycerol phosphoric acid and the nitrogenous bases. Of these, only the fatty acids are of any practical importance as far as combustion is concerned, although it is


also possible to convert the polyalcohols and the nitrogenous bases to combustible molecules. The fatty acids can be oxidized to acetic acid which, as mentioned above, is one of the organic fuels for the cellular furnace.

The carbohydrates are in essence polymer of sugar molecules held together by glycosidic linkages. A whole group of special enzymes are available which can hydrolyze polysaccharides like starch or glycogen into disaccharides and eventually into monosaccharides. In turn, the monosaccharides can now be degraded ultimately to pyruvic acid in a complex sequence of reactions collectively referred to as glycolysis.² Pyruvic acid is then combusted to CO₂ and water in the cellular furnace.²

THE MECHANISMS OF THE COMBUSTION PROCESS

There are three complex processes of a cyclical nature which largely define the combustion of organic molecules in the animal body. These are, respectively, glycolysis,² the citric acid cycle³ and the pentose cycle.⁴

In glycolysis, sugar in the form of glucose is cleaved into two molecules of lactic acid.



No CO₂ is formed in this sequence but the product of the cleavage is a first cousin to pyruvic acid which is one of the ultimate fuels. Thus glycolysis is a device for preparing the sugar molecule for combustion to CO₂ and H₂O. However, in this particular preparatory process, energy is harnessed even though true combustion involving molecular oxygen has not intervened.

The citric acid cycle is the key mechanism for cellular combustion. In this cycle pyruvic





FIG. 2. Structural formulae of: (a) choline phosphate and (b) ethanolamine phosphate.

acid is combusted to CO_2 and H_2O in a series of 5 oxidative steps:




This is also the major system in animal tissues for trapping chemical energy in a utilizable form.

The pentose cycle is a cyclical sequence of reactions which, in effect, can oxidize glucose to CO_2 and H_2O . No energy is harnessed in the process. However, the pentose cycle rarely, if at all, operates like the citric cycle. It would be more accurate to say that the pentose cycle is yet another device for converting glucose to intermediates which are capable of entering the citric acid cycle. There are other intermediates formed in the pentose cycle which are not relevant to the combustion process and much of the meaning of the cycle has to be found in these additional functions.


At this point it may be desirable to consider in more detail the three major cycles referred

to above. In this review the primary emphasis will be on exposing the chemical strategy, while the actual chemical changes will take a place of secondary importance.

Glycolysis. The most significant chemical event in glycolysis is the fragmentation of a 6-carbon sugar into two molecules, each containing 3 carbon atoms. It is not glucose as such which is fragmented but rather a phosphoric ester of an isomer of glucose, *viz.*, fructose. A specific phosphorylating agent known as adenosine triphosphate (ATP) and three separate enzymes are required for this conversion of glucose to fructose-1,6-diphosphate:



Fructose-1,6-diphosphate is cleaved into two similar but not identical halves:



Each of the two fragmentation products (which are interconvertible) can in turn be converted to pyruvic acid in the following sequence of reactions:



When anaerobic conditions prevail the product of glycolysis is not pyruvate but its reduction product, lactate. Under aerobic conditions, which are more pertinent to the reactions considered in the present review, not lactate, but pyruvate, is at the end of the glycolytic sequence.

The Citric Acid Cycle.³ The strategy of

the citric cycle is readily apparent from a consideration of figure 3.

Pyruvate is oxidatively decarboxylated to acetate and CO_2 :



"Acetate" condenses with oxalacetate (7) to form citrate (1) which in turn is converted






FIG. 3. The complete combustion of pyruvate by way of the citric acid cycle. The intermediates of the cycle are represented by numbers as follows: 1—citric acid, 2—isocitric acid, 3— α -ketoglutaric acid, 4—succinic acid, 5—fumaric acid, 6—malic acid, 7—oxaloacetic acid; "acetate" = acetyl-CoA. (Structural formulas are shown in fig. 1.)

back to oxalacetate in a sequence of 4 oxidative reactions (2→3, 3→4, 4→5, and 6→7). This point is more clearly shown below:




At the beginning of each cycle "acetate" is fed into the cycle via combination with oxalacetate. At the end of each cycle oxalacetate is formed. Thus each cycle achieves the complete combustion of acetate to CO_2 and water.


The condensation of "acetate" and oxalacetate to form citrate is a simple aldol type condensation:⁵




Citric acid rearranges to form isocitric acid:



Then isocitric acid is decarboxylated to α -ketoglutaric acid:



A second oxidative decarboxylation then takes place, this time of α -ketoglutaric acid to succinic acid:




Succinic acid is dehydrogenated to fumaric acid which in turn is hydrated to malic acid:



Finally, malic acid is oxidized to oxaloacetic acid and that is the end of the cycle:




In the interest of simplicity a notable omission was made. It is not acetic acid as such which enters the cycle but rather a compound of the acid with a special coenzyme known as coenzyme A (fig. 4).⁶ The active group of this coenzyme is a sulfhydryl group and we may represent the coenzyme by the notation CoASH. The active form of acetate is formed by interaction with the coenzyme in the presence of a specific enzyme and ATP:⁷⁻⁹



The thiol ester link in effect makes the esterified molecule much more reactive than the free acid.

Fatty acids are capable of being oxidatively and quantitatively converted to acetate and thus ultimately fatty acids can be burned to CO_2 and H_2O in the furnace of the citric cycle. Again, it is not the free fatty acid as such which is so converted, but rather the thiol ester of the fatty acid and coenzyme A and

he product is not free acetate but the coenzyme A ester thereof:




The oxidative fragmentation of the fatty acyl CoA ester to two carbon units proceeds by way of the well-known β -oxidation sequence.⁷⁻¹⁰

Amino acids can give rise to pyruvate, oxalacetate, α -ketoglutarate, succinate, fumarate or acetate—each one of which can be burned to CO_2 and H_2O in the citric cycle (table 1). In

this way amino acids, like fatty acids, can eventually be oxidized to CO_2 and H_2O after appropriate preparation.


Pentose Cycle. Glucose can be converted to pyruvic acid by a sequence of reactions other than that shown above for glycolysis. Again the starting point is glucose-6-phosphate which is formed by the phosphorylation of glucose by ATP. The aldehyde group of the sugar phosphate is oxidized to a carboxyl and the corresponding phosphorylated sugar acid is then oxidatively decarboxylated to a pentose phosphate:



Ribulose-5-phosphate can isomerize to another pentose xylulose-5-phosphate:



Then two molecules of pentosephosphate interact to form a 7-carbon sugar phosphate (sedoheptulose-phosphate) and a 3-carbon sugar (triosephosphate):



Triosephosphate is convertible to pyruvic acid according to equations 8 and 9.

The sequence of reactions from glucose to triosephosphate via pentose phosphate is part of the pentose cycle—an alternative to glycolysis for degrading the sugar molecule. The pentose cycle is present in many tissues but is not of the same quantitative significance as glycolysis.

COMBUSTION AND ENERGY PRODUCTION^{11, 12}

The complete oxidation of pyruvic acid to CO_2 and H_2O by molecular oxygen is an en-

ergy yielding process and in general metabolic CO_2 production runs parallel with energy generation. Part of the energy is released in




FIG. 4. The structural formula of coenzyme A.





FIG. 5. The structural formula of adenosine triphosphate (ATP).

the form of heat and the rest in the form of the bond energy of adenosine triphosphate (fig. 5). Thus when 1 molecule of pyruvate is fully oxidized to CO_2 and H_2O by oxygen 15 molecules of ATP are synthesized:



The released energy of oxidation is harnessed to a chemical synthesis, *viz.*, the esterification of ADP by inorganic phosphate with formation of ATP. The major source of biochemical energy is this coupling of citric cycle oxidations to synthesis of ATP.

The glycolysis of sugar to lactic acid also leads to synthesis of ATP. Thus the conversion of one molecule of sugar to two molecules of lactic acid leads to the synthesis of 2 molecules of ATP. By contrast the further oxidation of the two molecules of lactic acid to CO_2



FIG. 6. Electron micrograph of a section of beef heart muscle showing clusters of mitochondria along the muscle fibrils. The structures are magnified 30,300 diameters. The photograph was prepared by K. L. Filmer and Paul Kaesburg of the University of Wisconsin.

and H_2O by way of the citric cycle leads to the synthesis of 36 molecules of ATP—18 times as much as in the glycolytic sequence.

The oxidations of the citric cycle as well as the coupled synthesis of ATP are carried out in the mitochondrion—a subcellular entity. An electron micrograph of a cluster of mitochondria in heart muscle can be seen in figure 6. The thirty or more enzymes involved in the combustion and coupling processes are arranged and organized in a very precise pattern within the mitochondrion.¹² The harnessing of energy involves a rather intricate sequence of events. In each of the five oxidative steps of the citric acid cycle, electrons are transferred




FIG. 7. Oxidation-reduction components of the electron transfer system: (1) flavin adenine dinucleotide which contains a riboflavin moiety is the functional group of the flavoproteins; (2) the heme group of the hemoproteins (cytochromes *a*, *b*, *c*, and *c*₁) undergoes oxidation-reduction; (3) coenzyme Q₁₀.

from substrate to molecular oxygen through a series of catalysts capable of undergoing reduction and oxidation. This involves a structurally linked series of enzymes collectively designated as the electron transfer chain.

The electron transfer chain contains enzymes with the following functional groups: riboflavin, heme, and a quinone known as coenzyme Q.¹⁴ Each of these functional groups can undergo reversible reduction and oxidation (fig. 7) as can non-heme iron and copper^{15, 16} which is bound to some of the electron carrier enzymes. Electrons are transferred from one functional group to the next in line and each such transfer involves the oxidation of the first

and the reduction of the next component. Thus electrons are passed from one catalyst to the next in a sequence of some eight or more transfers. Oxygen is the terminal electron acceptor and the final product of the reduction is water. Some three of these transfers involve a simultaneous synthesis of ATP. How this is accomplished is still unknown and is one of the pressing problems of contemporary biochemistry.

CONCLUSION

It can be seen from the foregoing discussion that carbon dioxide is mainly a by-product of the energy supplying mechanisms of the cell. While it is true that a small amount is reutilized in urea formation¹⁸ and other carboxylation reactions,¹⁹ the bulk of the carbon dioxide is transported by the blood to the lungs where it is expired. The mechanism of this transport will be discussed in other sections of this symposium.

I am indebted to Dr. David E. Green for his help in preparing the manuscript.

REFERENCES

1. Calvin, M., and Poin, N. G.: Carboxylations and decarboxylations, *J. Cell Comp. Physiol.* **54**: (Suppl. 1) 51, 1959.
2. Stumpf, P. K.: "Glycolysis" in *Chemical Pathways of Metabolism*. D. M. Greenberg, Ed., New York, Academic Press, Inc., 1954, pp. 67-108.
3. Krebs, H. A.: The tricarboxylic acid cycle, *Harvey Lect. Series*: 165, 1948-49.
4. Wood, H. G.: Significance of alternate pathways in metabolism of glucose, *Physiol. Rev.* **35**: 841, 1955.
5. Bové, J., Martin, R. O., Ingraham, L. L., and Stumpf, P. K.: Studies of mechanism of action of condensing enzyme, *J. Biol. Chem.* **234**: 999, 1959.

6. Lipman, F.: *Biochemical mechanisms*, Harvey Lect. Series: 99, 1948-49.
7. Lynen, F.: Acetyl CoA and the fatty acid cycle, *Harvey Lect. Series* **48**: 210, 1952-53.
8. Mahler, H. R., Wakil, S. J., and Bock, R. M.: Studies of fatty acid oxidation; enzymatic activation of fatty acids, *J. Biol. Chem.* **204**: 453, 1953.
9. Green, D. E.: Fatty acid oxidation in soluble systems of animal tissues, *Biol. Rev.* **29**: 330, 1954.
10. Lynen, F., and Ochoa, S.: Enzymes of fatty acid metabolism, *Biochim. et Biophysica Acta* **12**: 299, 1953.
11. Green, D. E., and Fleischer, S.: "The Mitochondrial System of Enzymes in Chemical Pathways of Metabolism" vol. 1. David M. Greenberg, Ed., New York, Academic Press, Inc., 1960, pp. 41-96 (in press).
12. Chance, B.: Cytochromes—their nature and function in living cells in *Proceedings of the International Symposium of Enzyme Chemistry*, Tokyo-Kyoto 1957, K. Ichihara, Ed., New York, Academic Press, 1958, pp. 9-20.
13. Green, D. E.: Studies in organized enzyme systems, *Harvey Lect. Series*: 177, 1956-57.
14. Crane, F. L., Hatefi, Y., Lester, R. L., and Widmer, C.: Isolation of quinone from beef heart mitochondria, *Biochim. et Biophys. Acta* **25**: 220, 1957.
15. Beinert, H., and Sands, R. H.: On function of iron in DPNH cytochrome c reductase, *Biochem. Biophys. Res. Comm.* **1**: 171, 1959.
16. Sands, R. H., and Beinert, M.: On function of copper in cytochrome oxidase, *Biochem. Biophys. Res. Comm.* **1**: 175, 1959.
17. Wolf, D. E., Hoffman, C. H., Trenner, N. R., Arison, B. H., Shunk, C. H., Linn, B. O., McPherson, J. F., and Folkers, K.: Coenzyme Q; studies on coenzyme Q group, *J. Amer. Chem. Soc.* **80**: 4752, 1958.
18. Mackenzie, C. G., and DuVigneaud, V.: Source of urea carbon, *J. Biol. Chem.* **172**: 353, 1948.
19. Utter, M. F., and Kerch, D. B.: Formation of oxalacetate from pyruvate and CO₂, *J. Biol. Chem.* **235**: PC 17, 1960.