

Circulatory Effects of Peridural Block:

I. Effects of Level of Analgesia and Dose of Lidocaine

John J. Bonica, M.D.,* Peter U. Berges, M.D.,†
Ken-ichi Morikawa, M.D.‡

Effects of different levels of peridural analgesia on cardiovascular dynamics, blood gases and limb blood flow were studied in normovolemic man. Lidocaine, 2 per cent, was injected to achieve successively greater levels of analgesic block. During analgesia to T4 or below, MABP, TPR, and CO remained within normal limits; with T2-3 block, TPR decreased 15 per cent and CO increased 22.6 per cent, resulting in a 5 per cent increase in MABP; with blocks extending to T1 or higher, TPR decreased 18.7 per cent while CO remained slightly above normal, resulting in a 16.3 per cent reduction in MABP. Blood flow in the leg increased progressively to 292 per cent with T2-3 block, while blood flow in the arm decreased as much as 58 per cent. However, after extension of the block above T1, blood flow in the arm increased to 125 per cent above normal, while blood flow in the leg declined to 219 per cent above normal. The data suggest that cardiovascular responses to peridural block, unlike responses to subarachnoid block, are not the result of sympathetic blockade alone, but involve several other factors, including the dose of local anesthetic. When several doses of lidocaine result in arterial blood levels ranging from 4 to 7 $\mu\text{g/ml}$, most subjects develop increased CO. This may be sufficient to offset vasodilating effects of high sympathetic block so that MABP remains normal. (Key words: Peridural block; Hemodynamics; Sympathetic block; Lidocaine.)

DURING the past two decades peridural block has been used with progressively greater frequency. In many centers it has supplanted subarachnoid block as a major technique for production of regional anesthesia. Although many of the articles about it contain retrospective surveys of the incidences of arterial hypotension and other side-effects, data from well controlled studies of the effects of peridural block on body functions in man are scarce. Some aspects of the technique are still unclear, a fact emphasized by Bromage.¹ To help rectify this problem, we have studied systematically the cardiovascular, respiratory, renal, and hepatic effects of peridural block in normovolemic man.²⁻⁶ The purpose of the present study was to determine the effects of different levels of peridural analgesia on cardiovascular dynamics, blood gases, and limb blood flow in normovolemic man.

Methods and Materials

Ten healthy male volunteers, 24 to 42 years of age, were studied. Several days prior to the study each subject was examined and was thoroughly informed about the nature and risk of the proposed investigation, and a written consent was obtained. When the study began each subject had fasted for eight hours and was unmedicated. The studies were performed with the subject supine in a quiet room. Intravenous infusion of 5 per cent glucose in lactated Ringer's solution was started. Catheters were inserted under local anesthesia into the brachial artery and basilic vein and advanced into the subclavian artery and superior vena cava, respectively. The catheters were connected to strain gauges and a Gilson recorder. Using the technique previously described,⁷ two vinyl catheters were inserted into the peridural space, one with its tip located at the level of the second lumbar vertebra and

* Professor and Chairman.

† Instructor.

‡ Research Fellow.

Received from the Department of Anesthesiology and the Anesthesia Research Center, University of Washington School of Medicine, Seattle, Washington 98105. Accepted for publication September 14, 1970. Supported in part by USPHS Grants GM 15991-02 and GMO 1160-07 from the National Institute of General Medical Sciences, National Institutes of Health, and by the U. S. Army Research and Development Command, Office of the Surgeon General, Contract DA 49-13-MD-2231. Presented at the Annual Meeting of the American Society of Anesthesiologists, San Francisco, October 26-29, 1969.

Reprint requests should be addressed to Dr. Bonica.

the other with its tip at the second thoracic vertebra. Whitney gauges were then placed around one forearm and one leg to measure limb blood flow.⁸

Following a 30-minute rest period, control measurements were made. Variables recorded continuously were arterial blood pressure (ABP), central venous pressure (CVP), heart rate (HR) and electrocardiogram (ECG); serial measurements were made of cardiac output (CO), left ventricular performance, limb blood flow and arterial blood gas and lidocaine levels. CO was determined by the indicator-dilution technique with indocyanine green as indicator. Arterial blood was sampled anaerobically for determination of P_{aO_2} and pH. We calculated P_{aCO_2} by the tonometric method of Astrup and base excess by the Siggaard-Andersen method.⁹ Arterial blood levels of lidocaine were measured by gas chromatography according to a modified method of Beckett *et al.*¹⁰ Stroke volume (SV) was calculated from CO and HR. Total peripheral resistance was calculated as follows:

TPR units (dynes/sec/cm⁻⁵)

$$= \frac{\Delta ABP \text{ (mm Hg)}}{CO \text{ (ml/sec)}} \times 1,332$$

Limb blood flow was measured by venous occlusive plethysmography with mercury in a silastic strain gauge (Whitney) as the transducer.⁸ To evaluate ventricular function, maximum dP/dt in the subclavian artery was obtained by electronic differentiation.¹¹ Left ventricular minute work (LVMW) was calculated as: $CO \times MABP \times 0.0144$ (kg/m/min), and left ventricular stroke work (LVS_W) as: $SV \times MABP \times 0.0135$ (g-meters).

The achievement of total sympathetic block was ascertained with the psychogalvanic reflex (PCR)¹² in the hand. The cardiac response to moderate hypercapnia induced by allowing the subject to rebreathe in a closed system for six minutes was determined. This procedure usually resulted in a P_{aCO_2} of 65 to 75 mm Hg, which in normal man increases HR and CO significantly.

Following completion of control measurements, 10 ml of 2 per cent lidocaine without epinephrine were injected into the lumbar peri-

dural catheter at a rate of 0.5 ml/sec. This dose usually produced analgesia to T10-11. Beginning five minutes after injection, the level of analgesia was evaluated every two minutes by firm pinprick stimulation. When the level had been stable for five minutes (usually 22 to 25 minutes after injection), all variables were measured. In the last five subjects, blood samples for lidocaine measurements were obtained at 5- to 10-minute intervals. Once all measurements had been completed, the level of analgesia was extended cephalad in a stepwise manner to three or four more successively higher segments by injecting progressively greater volumes of 2 per cent lidocaine. The levels were categorized as T10-11, T8-9, T6-7, T4-5, T2-3, and T1 or above. The highest level of analgesia achieved required a double injection, one sufficiently large to produce block to T4-5 through the lumbar catheter and another of 5 to 7 ml through the upper thoracic peridural catheter. Measurements were made at each level after the block had been stable for five minutes. The times between injections averaged slightly more than an hour.

Computer analysis of the data was performed on a within-person basis, i.e., using each subject as his own control. The calculations included individual differences and paired differences from control, test results expressed as proportions of control and logarithms of tests as proportions of control. Significance of the changes was determined by the use of the paired *t* test and by the Wilcoxon matched-pairs signed-rank test of differences.¹³

Results

Complete data at four or more levels of block were obtained from each subject (table 1). The total doses of lidocaine given ranged from 880 mg to 1,860 mg, with a mean for the group of 1,586 mg. On a body-weight basis the doses ranged from 11 to 20 mg/kg, with a mean of 16.8 mg/kg. Absence of data at one or two of the six levels resulted in part from the unpredictable responses of some subjects to the average dose. Thus, whereas 14 ml produced T6-7 blocks in five subjects, it produced higher or lower blocks in the other five. No data were obtained for four indi-

viduals with blocks of T1 or above: we terminated the experiment at T2 upon the request of one volunteer and at T4 in another because of signs of systemic toxicity. In the other two subjects we produced hypalgesia above T1 but could not achieve complete sympathetic interruption even after a second injection through the upper thoracic catheter. In these subjects the durations of successive blocks were significantly shorter than the preceding ones, so that by the fifth injection intensity was inadequate and duration less than 20 minutes. This suggests that these subjects probably developed increasing tolerance to repeated doses of lidocaine (tachyphylaxis) unusually rapidly.

The effects on circulation, blood gases and limb blood flow produced by various levels of block are summarized in table 2 and figures 1 and 2. The table contains the control average responses for the ten volunteers and the mean of the individual differences and standard error of mean differences from control obtained at each block level. Thus, the values at the T10 level are the averages of the eight differences from control for those (eight) volunteers who developed T10 analgesia and had measurements made at that level. The graphs show the quantity Δ , which is the geometric average of individual per cent responses. It deserves re-emphasis that these are based on means of individually calculated per cent differences and not per cent differences of the means.

During peridural block at T4 or below, all variables remained normal except HR, which was significantly increased at all levels except T6-7. When the block was extended to T2-3, TPR fell a mean of 14.6 per cent while CO increased a mean of 22.6 per cent, resulting in a 5 per cent increase in MABP. Two subjects had decreases of CO, while the other six had increases ranging from 21 per cent to 76 per cent, with a mean 41 per cent above control. The increase in CO resulted wholly from a 22 per cent increase in HR and was associated with a 26.7 per cent increase in LVMW and a 17.8 per cent increase in dp/dt. Except for heart rate, none of the changes was statistically significant. When analgesia was raised to T1 or above, TPR fell 18.7 per cent;

TABLE 1. Dosage of Lidocaine and Level of Peridural Analgesia

| Level of Analgesic Block | Number of Subjects | Volume of 2 Per Cent Lidocaine (ml) | |
|--------------------------|--------------------|-------------------------------------|-------|
| | | Mean | Range |
| T10-11 | 8 | 10.5 | 10-12 |
| T8-9 | 9 | 12.0 | 10-14 |
| T6-7 | 5 | 14.0 | 12-16 |
| T4-5 | 8 | 15.5 | 13-18 |
| T2-3 | 8 | 20.5 | 16-25 |
| T1+ | 6 | 22.0 | 18-26 |

CO, HR, and SV remained near normal while MABP decreased a significant 16.3 per cent. There were also decreases of 19.5 per cent in LVSF, 13.7 per cent in LVMW, and 14.3 per cent in dp/dt. CVP remained normal with all levels except for increases of 11 per cent for T8-9 and 22 per cent with blocks above T1 (statistically significant).

Limb blood flows changed significantly. The increases in blood flow in the leg with different levels of analgesia were: 183 per cent at T10-11, 245 per cent at T8-9, 247 per cent at T6-7, 257 per cent at T4-5, and 292 per cent at T2-3. At these levels, blood flows in the arm were -27.5, -50, -32, -58, and +4 per cent from control, respectively. As the block was extended above T1, blood flow in the arm increased to 125 per cent above normal, while blood flow in the leg declined to 219 per cent above normal, indicating redistribution of flow.

There were no significant changes in P_{aO_2} and P_{aCO_2} with blocks of T4 or below, but a 10 per cent increase in P_{aO_2} occurred with T2-3 block, and a 20 per cent increase in P_{aO_2} and an 11 per cent decrease in P_{aCO_2} occurred with analgesia above T1. There were significant decreases in pH at T6-7 and T4-5 and in base excess at all block levels above T9.

The arterial blood levels of lidocaine followed the previously established curves,¹⁴⁻¹⁷ obviously related to the amount of the drug injected. Lidocaine was found in the blood as early as five minutes after peridural injection; a peak level was achieved in 10 to 30 minutes, followed by a rapid decay, so that 60 minutes after injection the blood level was

TABLE 2. Hemodynamic Responses and Blood Gases at Various

| | Number of Subjects | MABP (mm Hg) | CO (l/min) | HR (beats/min) | SV (ml) | TPR (dynes/sec/cm ²) | CVP (cm H ₂ O) | dP/dt (mm Hg/sec) |
|------------------|--------------------|--------------|--------------|----------------|-------------|----------------------------------|---------------------------|-------------------|
| Control value | 10 | 94.6 ± 2.2 | 6.83 ± 0.46 | 67.4 ± 0.46 | 101.1 ± 3.4 | 1159 ± 91 | 8.5 ± 0.7 | 639 ± 40 |
| Mean Differences | | | | | | | | |
| T10-11 | 8 | 1.4 ± 1.4 | 0.42 ± 0.19 | 2.5 ± 0.7* | 2.5 ± 1.9 | -38 ± 47 | 0.0 ± 0.8 | 28 ± 33 |
| T8-9 | 9 | 1.9 ± 2.3 | 0.00 ± 0.29 | 3.3 ± 1.4* | -4.6 ± 2.3 | 24 ± 41 | 0.9 ± 0.9 | -18 ± 34 |
| T6-7 | 5 | -2.0 ± 2.0 | -0.09 ± 0.43 | 2.0 ± 1.1 | -3.6 ± 5.0 | -48 ± 64 | 0.3 ± 1.1 | 63 ± 30 |
| T4-5 | 8 | 1.5 ± 3.7 | 0.15 ± 0.25 | 5.5 ± 1.0* | -5.6 ± 2.6 | -24 ± 40 | 0.9 ± 1.3 | 4 ± 30 |
| T2-3 | 8 | 4.9 ± 4.6 | 1.41 ± 0.77 | 14.5 ± 3.8* | 0.6 ± 6.2 | -201 ± 121 | 1.1 ± 1.3 | 148 ± 88 |
| T1+ | 6 | -15.3 ± 5.3* | 0.11 ± 0.55 | 4.0 ± 4.5 | -3.0 ± 7.0 | -241 ± 159 | 2.2 ± 0.9* | -74 ± 85 |

* $P < 0.05$.

30 to 40 per cent of the peak value. With subsequent injections, peak levels increased above those of preceding injections, the final

values ranging from 4.0 to 6.65 $\mu\text{g/ml}$. A curve from one subject representative of the group is shown in figure 3.




FIG. 1. Circulatory responses to various levels of peridural analgesia.

Levels of Peridural Analgesia in Ten Subjects (Mean \pm SE)

| ABF (ml/100 ml tissue/min) | IBF (ml/100 ml tissue/min) | SW (g-ineters) | MW (kg/m ² /min) | PaO ₂ (mm Hg) | Paco ₂ (mm Hg) | pH | BE (mEq/l) |
|----------------------------------|----------------------------------|-------------------|--------------------------------|-----------------------------|------------------------------|-------------------|-----------------|
| 4.1 \pm 0.5 | 3.0 \pm 0.7 | 137.0 \pm 3.3 | 8.7 \pm 0.6 | 87.3 \pm 2.9 | 37.8 \pm 0.7 | 7.38 \pm 0.00 | -2.1 \pm 0.3 |
| from Control | | | | | | | |
| -1.3 \pm 0.5* | 6.4 \pm 2.2* | 5.4 \pm 2.0* | 0.7 \pm 0.2* | 0.6 \pm 2.8 | -1.5 \pm 0.8 | 0.00 \pm 0.00 | -0.7 \pm 0.3 |
| -2.0 \pm 0.5* | 7.9 \pm 1.8* | -3.7 \pm 5.3 | 0.1 \pm 0.5 | 1.5 \pm 1.7 | -1.8 \pm 0.8 | 0.00 \pm 0.01 | -0.6 \pm 0.2* |
| -1.4 \pm 0.5 | 8.8 \pm 3.0* | -7.2 \pm 8.6 | -0.2 \pm 0.7 | 3.4 \pm 5.3 | 1.1 \pm 0.8 | -0.02 \pm 0.00* | -0.7 \pm 0.1* |
| -2.1 \pm 0.7* | 8.0 \pm 1.9* | -4.9 \pm 8.4 | 0.3 \pm 0.6 | 1.1 \pm 2.7 | -0.9 \pm 1.1 | -0.01 \pm 0.00* | -0.9 \pm 0.3* |
| 1.3 \pm 1.9 | 6.7 \pm 1.3* | 7.9 \pm 12.7 | 2.4 \pm 1.4 | 8.3 \pm 2.3* | -1.4 \pm 0.7 | 0.00 \pm 0.01 | -0.7 \pm 0.3* |
| 3.7 \pm 1.2* | 5.3 \pm 1.8* | -25.3 \pm 10.0 | -1.2 \pm 1.2 | 20.8 \pm 13.1 | -3.9 \pm 2.1 | 0.02 \pm 0.02 | -0.8 \pm 0.6 |

Discussion

Until recently, cardiovascular responses to peridural block, like those to subarachnoid block, were believed to be due primarily to the effects on sympathetic nerves.¹⁸⁻²⁰ Many assumed, therefore, that the cardiovascular changes associated with peridural anesthesia were similar to those of spinal anesthesia. Data from our previous study of human volunteers showed that this was not the case and suggested the involvement of other factors, including the use of epinephrine, the local anesthetic employed, and perhaps the increase in peridural pressure during and following injection.² Bromage¹ has suggested five ways that peridural analgesia can influence cardiovascular dynamics: 1) dilatation of resistance and capacitance vessels; 2) paralysis of cardiac sympathetic fibers from the upper four thoracic segments, resulting in bradycardia with reduction of cardiac output; 3) vascular absorption of local anesthetics, which causes a decrease in cardiac output owing to beta receptor blockade and peripheral vascular smooth muscle depression; 4) beta receptor stimulation owing to vascular absorption of epinephrine if present, increasing CO and decreasing TPR; and 5) sudden elevation of cerebrospinal fluid pressure owing to peridural injection, producing transient reflex increases in vasomotor tone and cardiac output. We have shown the important role played by epinephrine^{2,6} and the lack of effect of increased peridural pressure.⁴

The results of the present study indicate that with continuous peridural block achieved with lidocaine alone, the problem is even more complex than proposed by Bromage. Apparently, factors other than the extent of sympathetic

block and depressant effects of local anesthetics are involved. Moreover, the factors interact and the net effect varies with circumstances. Peridural block extending to T4-5 usually is not associated with significant hypotension. Normovolemic man apparently can maintain cardiac output near normal or even increase it and compensate for vasodilatation in the anesthetized part of the body by vasoconstriction in the unanesthetized part. Results of many studies of spinal anesthesia²¹ and several studies of single-dose peridural block^{2,22} confirm this observation. Thus, Shimosato and Etsten²² found that both spinal and peridural blocks to sensory levels between




FIG. 2. Arterial blood gases and acid-base balances during various levels of peridural analgesia.




FIG. 3. Blood levels of lidocaine in one subject following several peridural injections.

T4 and T7 were associated with a 16 per cent decrease in mean arterial pressure, a 12 per cent decrease in peripheral resistance, but no significant change in stroke volume index, heart rate and, consequently, cardiac index. In previous studies^{2,6} of peridural block with lidocaine without epinephrine, we found that T5 analgesia was associated with decreases of about 5 to 10 per cent in mean arterial pressure and peripheral resistance, whereas cardiac output remained within 5 per cent of normal.

The cardiovascular alterations with analgesia to T2-3 in the present study were quite unexpected. Because such blocks presumably involved three or four of the five cardiac sympathetic segments, significant reductions in HR and SV, and with them CO, were anticipated. Instead, we saw increases in HR and CO and commensurate increases in LVMV and in dP/dt . The mechanism of these unexpected cardiovascular effects of high peridural block achieved with several injections of lidocaine is not known. One may speculate that they represent the action of lidocaine on the cardiovascular system.

Although lidocaine is generally believed to produce direct cardiovascular depression, ample evidence now indicates that moderate arterial blood levels of lidocaine are usually associated with increased cardiac output and slight hypertension. This was first suggested by Kimmy and Steinhaus,²³ who found in-

creased blood pressures in patients given intravenous lidocaine (1 to 2 mg/kg body weight) during general anesthesia. This was subsequently confirmed in human volunteers by Foldes and colleagues²⁴ and Jorfeldt and associates,²⁵ who reported that blood levels of lidocaine ranging from 3 to 6 µg/ml were associated with increases in arterial pressure, cardiac output and tachycardia. Similar results were found in patients receiving single intravenous injections of lidocaine for treatment of cardiac arrhythmias.^{26,27}

The cardiac stimulating effects of lidocaine may be produced through central nervous system action or potentiation of endogenous epinephrine or norepinephrine, or may simply be a compensatory reflex response to vasomotor blockade. A central mechanism is suggested by the studies of Kao and Jalar,²⁸ who found that intravenous lidocaine in doses of 1 to 2 mg/kg body weight produced significant increases in cardiac output and blood pressure. The increase in cardiac output did not occur after midbrain transection, suggesting a central action of the drug. This hypothesis was substantiated by their results in cross-circulation experiments in which they were able to isolate the effects of lidocaine on the central nervous system from those on heart and peripheral vessels. The potentiating mechanism is suggested by studies of Van Dongen²⁹ and D'Amato and Truant,³⁰ who found that lidocaine, cocaine, and some other local anes-

thetics consistently potentiated the effects of exogenous epinephrine and norepinephrine on blood pressure. These results prompted Austen and Moran³¹ to suggest that, conceivably, the rise in blood pressure following administration of lidocaine may be due to potentiation of endogenous catecholamines. Currently we are studying this problem in man.

Our data suggest that in normal man arterial blood levels of lidocaine ranging from 4 to 7 µg/ml are usually associated with an increase in cardiac output which may be sufficient to offset the vasodilating effect of high sympathetic block, provided some cardiac sympathetic nerves remain intact. Because such blood levels usually develop following single peridural injections of 400 to 500 mg or following several injections of 200 to 300 mg of lidocaine,^{14-16, 19} these findings have obvious clinical implications. They are especially important in patients receiving "continuous" peridural block for extensive surgical intervention or prolonged labor, for which injections are usually repeated hourly. Since man requires considerably more than an hour to metabolize and eliminate lidocaine,³²⁻³⁴ the drug will accumulate to a significant level in blood and body tissues.

Our results with peridural block extending above T1 are not quite in accord with those of McLean and associates,³⁵ who published the only other hemodynamic study of total sympathetic block achieved with peridural anesthesia. In seven patients scheduled for major operations, they studied the separate and combined hemodynamic effects of segmental cervicothoracic (C5 to T4) peridural block and total peridural block (C5 to S5) achieved by supplementing the former with an extensive thoracolumbar peridural block. Total sympathetic block decreased CVP 26 per cent, TPR 6 per cent, HR 8 per cent, SV 13 per cent, CO 19 per cent and MABP 20 per cent. In contrast, in our subjects, CO remained slightly above normal, whereas TPR decreased 20 per cent, resulting in a 15 per cent decrease in MABP. The differences between the results of the two studies may be related to differences between the subjects' conditions, the agents used, and total doses injected. Their patients had a mean age of 49 years, had pathologic conditions requiring surgical

treatment, and were given mepivacaine (Carbocaine) in doses of 5 mg/kg, whereas our ten volunteers had a mean age of 32 years, were healthy, and received lidocaine in total doses averaging nearly 17 mg/kg.

In conclusion, because our data were obtained in young healthy subjects they should not be extrapolated to sick surgical patients. Furthermore, the cardiovascular responses of individuals to different levels of sympathetic blockade vary widely, depending upon degree of sympathetic tone prior to the block, whether blood volume is normal, the efficiency of neurogenic homeostatic mechanisms, and the condition of the heart. Patients with high sympathetic tone provoked by apprehension, injury, or disease, or with hypovolemia, have much greater hemodynamic changes. Patients in whom circulatory homeostatic mechanisms are dampened by narcotics, intravenous anesthetics, and other depressant drugs may not be able to mobilize compensatory mechanisms as effectively, and consequently will develop greater degrees of cardiovascular depression. Finally, with serious myocardial disease the reserves of both the extrinsic and intrinsic cardiac regulatory systems are decreased as the heart dilates and resting sympathetic tone increases.^{36, 37} Under these conditions, removal of sympathetic tone is poorly tolerated. This is especially true of patients with congestive heart failure in whom there is significant arteriolar and venular constriction modulated by increased sympathetic activity to complement the heart's reduced performance.³⁸ Since the increase in total peripheral vascular resistance in heart failure is an important compensatory mechanism for maintaining arterial pressure when cardiac output is low, extensive sympathetic blockade in these patients may prove disastrous.

References

1. Bromage PR: Physiology and pharmacology of epidural analgesia. A review. *ANESTHESIOLOGY* 28:592, 1967
2. Ward RJ, Bonica JJ, Freund FG, *et al.*: Epidural and subarachnoid anesthesia, cardiovascular and respiratory effects. *JAMA* 191: 275, 1965
3. Bonica JJ, Kennedy WF Jr, Ward RJ, *et al.*: A comparison of the effects of high subarachnoid and epidural anesthesia. *Acta Anaesth Scand suppl* 23:429, 1966

4. Akamatsu TJ, Ward RJ, Bonica JJ, *et al.*: Cardiovascular response to increased epidural pressure in the elderly surgical patient. *Pacif Med Surg* 75:161, 1967
5. Freund FG, Bonica JJ, Ward RJ, *et al.*: Ventilatory reserve and level of motor block during high spinal and epidural anesthesia. *ANESTHESIOLOGY* 28:834, 1967
6. Kennedy WF Jr, Sawyer TK, Gerbershagen HU, *et al.*: Systemic cardiovascular and renal hemodynamic alterations during peridural anesthesia in normal man. *ANESTHESIOLOGY* 31:414, 1969
7. Bonica JJ: Continuous peridural block. *ANESTHESIOLOGY* 17:626, 1956
8. Clarke RSJ, Hellon RF: Venous collection in forearm and hand measured by the strain-gauge and the volume plethysmograph. *Clin Sci* 16:103, 1957
9. Siggaard-Andersen O: Blood acid-base alignment nomogram. Scales for pH, pCO₂, base excess of whole blood of different hemoglobin concentrations, plasma bicarbonate, and plasma total CO₂. *Scand J Clin Lab Invest* 15:211, 1965
10. Beckett AH, Boyes RN, Parker JBR: Determination of lignocaine in blood and urine in human subjects undergoing local analgesic procedures. *Anaesthesia* 20:294, 1965
11. Rushmer RF: Initial ventricular impulse. A potential key to cardiac evaluation. *Circulation* 29:268, 1964
12. Lewis LW: Evaluation of sympathetic activity following chemical or surgical sympathectomy. *Curr Res Anesth Analg* 34:334, 1955
13. Goldstein A: Biostatistics: An Introductory Text. New York, The Macmillan Co., 1964
14. Morikawa S, Kaneko S, Miura J: The changes in the blood concentrations of Xylocaine administered through various routes in man. A study on the toxic reactions to local anesthetics. *Far East J Anaesth* 4:27, 1963
15. Braid DP, Scott DB: Dosage of lignocaine in epidural block in relation to toxicity. *Brit J Anaesth* 38:596, 1966
16. Mazze RI, Dunbar RV: Plasma lidocaine concentrations after caudal, lumbar epidural, axillary block, and intravenous regional anesthesia. *ANESTHESIOLOGY* 27:574, 1966
17. Lund PC, Covino BG: Distribution of local anesthetics in man following peridural anesthesia. *J Clin Pharmacol* 7:324, 1967
18. Bromage PR: Spinal Epidural Analgesia. Baltimore, The Williams & Wilkins Co., 1954
19. Lund PC: Peridural Analgesia and Anesthesia. Springfield, Illinois, Charles C Thomas, 1966
20. Wylie WD, Churchill-Davidson HC: A Practice of Anaesthesia. Second edition. London, Lloyd Luke, 1966
21. Greene NM: Physiology of Spinal Anesthesia. Second edition. Baltimore, The Williams & Wilkins Co., 1969
22. Shimosato S, Esten BE: The role of the venous system in cardiocirculatory dynamics during spinal and epidural anesthesia in man. *ANESTHESIOLOGY* 30:619, 1969
23. Kimmey JR, Steinhaus JE: Cardiovascular effects of procaine and lidocaine (Xylocaine) during general anesthesia. *Acta Anaesth Scand* 3:9, 1959
24. Foldes FF, Molloy R, McNall PG, *et al.*: Comparison of toxicity of intravenously given local anesthetic agents in man. *JAMA* 172: 1493, 1960
25. Jorfeldt L, Löfström B, Pernow B, *et al.*: The effect of local anesthetics on the central circulation and respiration in man and dog. *Acta Anaesth Scand* 12:153, 1968
26. Harrison DC, Sprouse JH, Morrow AG: The antiarrhythmic properties of lidocaine and procaine amide. Clinical and physiologic studies of their cardiovascular effects in man. *Circulation* 28:486, 1963
27. Binnion FF, Murtagh G, Pollock AM, *et al.*: Relation between plasma lignocaine levels and induced haemodynamic changes. *Brit Med J* 3:390, 1969
28. Kao FF, Jalar UH: The central action of lignocaine and its effect on cardiac output. *Brit J Pharmacol* 14:522, 1959
29. Van Dongen K: The action of cocaine, procaine, Xylocaine and their combination with adrenaline on the blood-pressure. *Arch Int Pharmacodyn* 96:49, 1953
30. D'Amato HE, Truant AP: Potentiation of epinephrine and norepinephrine blood pressure responses by local anesthetic agents. *Arch Int Pharmacodyn* 101:113, 1955
31. Austen WC, Moran JM: Cardiac and peripheral vascular effects of lidocaine and procainamide. *Amer J Cardiol* 16:701, 1965
32. Beckett AH, Boyes RN, Appleton PJ: The metabolism and excretion of lignocaine in man. *J Pharm Pharmacol* 18:76S, 1966
33. Åkerman B, Åström A, Ross S, *et al.*: Studies on the absorption, distribution and metabolism of labelled prilocaine and lidocaine in some animal species. *Acta Pharmacol Toxicol* 24:389, 1966
34. Sung CY, Truant AP: The physiological disposition of lidocaine and its comparison in some respects with procaine. *J Pharmacol Exp Ther* 112:432, 1954
35. McLean APH, Mulligan GW, Otton P, *et al.*: Hemodynamic alterations associated with epidural anesthesia. *Surgery* 62:79, 1967
36. Braunwald E, Ross J Jr, Sonnenblick EH: Mechanisms of contraction of the normal and failing heart. *New Eng J Med* 277: 910, 1967
37. Braunwald E, Chidsey CA, Pool PE, *et al.*: Congestive heart failure: Biochemical and physiological considerations. *Ann Intern Med* 64:904, 1966
38. Mason DT: The autonomic nervous system and regulation of cardiovascular performance. *ANESTHESIOLOGY* 29:670, 1968