

Dave Massa bought his own supplies and, in his home basement, made by hand a series of prototypes, which were gradually improved to produce the styletted catheter essentially as it is today. Early in his work, colleagues provided more derision than encouragement, but Dave continued to provide better models for trial in the operating rooms until the demand for "Massa" needles, as they were known locally, began to grow. It not being ethical for a physician to patent an instrument, the Rochester Products Company of Rochester, Minnesota, was engaged to produce the needles for commercial use and they were marketed as the "Rochester" needles, although the head of the

Department was inclined to want them to be given his name. This experience should lend encouragement to innovators with ideas who lack complicated and expensive laboratories. Advances just as important remain to be introduced in the future.

JOHN W. PENDER, M.D.
*Clinical Professor of Anesthesia
Medical School Affiliates
Stanford University School of Medicine
Stanford, California 94305*

(Accepted for publication January 22, 1979.)

Anesthesiology
51:183, 1979

Tracheal Constriction or Decreased Lung-Thorax Compliance from Opiates

To the Editor:—Using water-filled floppy cuffs on endotracheal tubes to measure changes in tracheal tone, Yasuda *et al.* concluded that morphine and fentanyl cause tracheal constriction, as evidenced by increased intracuff pressures.¹ Fentanyl and morphine may decrease chest-wall compliance ("tight chest" syndrome) by increasing thoracic and abdominal muscle rigidity^{2,3} and may also decrease pulmonary compliance by bronchoconstriction.⁴ Neuromuscular blocking agents will relieve muscle rigidity and improve chest-wall compliance. Crawley and Cross demonstrated that intracuff pressure reflects mean airway pressure.⁵ If tidal volume is maintained as lung and chest-wall compliance decreases, airway pressure and mean intracuff pressure will increase. Yasuda *et al.* do not report changes in airway pressure or tidal volume. Their use of pancuronium may prevent the decreased compliance from muscle rigidity, but not that due to bronchoconstriction. Employing a pressure-cycled ventilator (Bird Mark 4, 8[®]) will, by definition, prevent an increase in airway pressure over the set pressure. However, altering the tidal volume to maintain Pa_{CO₂} values within certain boundaries will necessitate changing the pressure limits of the

ventilator. One may thus question whether the increases in cuff pressure observed by Yasuda *et al.* were due to tracheal constriction, to decreases in lung or chest-wall compliance, or to a combination of all three.

BARRY J. SEGAL, M.D.
*Department of Anesthesiology
Tufts-New England Medical Center
Boston, Massachusetts 02111*

REFERENCES

1. Yasuda I, Hirano T, Yusa T, et al: Tracheal constriction by morphine and fentanyl in man. *ANESTHESIOLOGY* 49: 117-119, 1978
2. Sokoll MD, Hoyt JL, Gergis SD: Studies in muscle rigidity, nitrous oxide, and narcotic analgesic agents. *Anesth Analg (Cleve)* 51:16-20, 1972
3. Kallos, T, Wyche MQ, Garman JK: The effects of Innovar on functional residual capacity and total chest compliance in man. *ANESTHESIOLOGY* 39:558-561, 1973
4. Aviado DM: Regulation of bronchomotor tone during anesthesia. *ANESTHESIOLOGY* 42:68-80, 1975
5. Crawley BE, Cross DE: Tracheal cuffs: A review and dynamic pressure study. *Anaesthesia* 30:4-11, 1975

(Accepted for publication January 22, 1979.)

Anesthesiology
51:183-184, 1979

In reply:—In our study, tidal volume was held constant by using the mechanical stop on the Bird Mark 4[®] and by setting the pressure at more than 25 cm H₂O with the Mark 8[®]. Rigidity of the chest wall was avoided by the administration of pancuronium.

In measuring tracheal tone, the small waves produced by the ventilator reflect pressure changes in the airway, whereas amplified waves would reflect the increase of airway pressure by decreased lung or chest-wall compliance. In our study, amplified waves were