

Correspondence

Anesthesiology
51:567, 1979

Hazards of Edrophonium

To the Editor:—Youngberg¹ emphasized a serious and potentially fatal complication during the use of edrophonium chloride, 10 mg, for the conversion of paroxysmal atrial tachycardia. His caution concerning dosage was extended to the seemingly fit, elderly patient not taking any medication. We have recently had occasion to verify the concern expressed by Youngberg with an observation of an example of its use that caused a severe dysrhythmia.

A 62-year-old, previously healthy woman came to the Emergency Department complaining of pain in the right arm. In the past, she had had several episodes of short-lived palpitations, the nature of which had never been documented, the last episode being a year previously. She was not taking any medication. Physical examination was unremarkable, apart from a rapid heart rate and a blood pressure of 80/50 torr. A 12-lead electrocardiogram demonstrated supraventricular tachycardia at a rate of 170 beats/min, with no ischemic change. During monitoring of the electrocardiogram, edrophonium, 5 mg, was administered intravenously over 1 min. Within 2 min the supraventricular tachycardia was converted transiently to sinus rhythm at a rate of approximately 60 beats/min, followed by a 30-sec interval of atrial bradycardia at approximately 50 beats/min with no ventricular activity. Consciousness was lost within

15–20 sec. Sinus rhythm at a rate of approximately 60 beats/min was restored immediately by a precordial thump. The patient immediately regained consciousness, and full recovery occurred. We believe that when edrophonium is used to convert supraventricular tachycardias, the dosage for every patient should be 2 mg initially, as suggested by Youngberg. Furthermore, when ventricular asystole occurs as a result of the acetylcholine response, a precordial thump should be performed immediately.

S. LUDWIG, M.D.
Medical Resident
Health Sciences Centre
G. BRISTOW, M.D.
Associate Professor
Department of Anesthesia and
Director, Emergency Medical Service Unit
Health Sciences Centre
University of Manitoba
Winnipeg, Manitoba

REFERENCE

1. Youngberg JA: Cardiac arrest following treatment of paroxysmal atrial tachycardia with edrophonium. *ANESTHESIOLOGY* 50:234–235, 1979

(Accepted for publication June 28, 1979.)

Anesthesiology
51:567–568, 1979

Succinylcholine and Serum Calcium

To the Editor:—Bourke and Rosenberg¹ reported changes in plasma calcium during the administration of succinylcholine with or without pretreatment with a nonpolarizing muscle relaxant. Several months ago we undertook a similar study, which we have not submitted for publication. Our study differed from that of Bourke and Rosenberg in the following manner. We studied 18 patients. Arterial blood samples were drawn after premedication with atropine, 0.5 mg, intravenously and induction of anesthesia with thiopental (control); immediately after cessation of muscle fasciculations from succinylcholine 1 mg/kg

intravenously (sample 1); and one minute after cessation of fasciculations (sample 2). Ionized plasma Ca^{++} was measured from plasma separated anaerobically (Orion Biomedical® model SS 20). Simultaneous determinations of plasma pH (Radiometer® BMS 3 + PHM 73) were made.

When the slight respiratory acidosis (7.38) observed is corrected to pH 7.40 using Schaer and Bachmann's regression equation,² our results are: control, 1.96 ± 0.06 mEq/l; sample 1, 1.99 ± 0.06 mEq/l; sample 2, 1.99 ± 0.06 mEq/l. Statistical calculations using analysis of variance showed no significant dif-

ference among these values. The disparity between our findings and those of Bourke and Rosenberg probably arises from differences in protocol.

D. GLOMAUD, M.D.
Professeur Agrégé J. C. Salamagne
Département d'Anesthésiologie
Hôtel Dieu de Paris
1, Place du Parvis Notre-Dame
75 004 Paris, France

REFERENCES

1. Bourke AL, Rosenberg M: Changes in total serum Ca^{++} , Na^{+} and K^{+} with administration of succinylcholine. *ANESTHESIOLOGY* 49:361-363, 1978
2. Schaer H, Bachmann V: Ionized calcium in acidosis: Differential effect of hypercapnic and lactic acidosis. *Br J Anaesth* 46:842-848, 1974

(Accepted for publication June 28, 1979.)

Anesthesiology
51:568, 1979

A New Method for Nasogastric Tube Insertion

To the Editor:—It is often difficult to pass a nasogastric tube in an unconscious patient after a cuffed endotracheal tube is in place. We describe a reliable, relatively simple and safe method of nasogastric tube insertion. A conventional esophageal stethoscope is modified to accept 2–3 cm of a nasogastric tube tip by cementing a thin layer of rubber from the finger of a glove over the cuff, so the proximal cuff end has a pouch. A nasogastric tube is then passed through the nose until the distal tip can be taken out through the mouth. The nasogastric tube is placed into the cuff pouch (fig. 1), and the esophageal stethoscope, along with the nasogastric tube, is then passed orally into the esophagus until maximal-intensity heart sounds are heard. This locates the tube tip distal to the endotracheal tube cuff. The nasogastric tube is then held in place and separated from the esophageal stethoscope by advancing the stethoscope 3 cm and rotating 180 degrees. The nasogastric tube is then advanced easily into the stomach. The esophageal stethoscope can be left in place or removed.

KIE-CHUL OHN, M.D.
Resident
WEN-HSIEN WU, M.D.
Associate Professor
Department of Anesthesiology
New York University Medical Center




FIG. 1. The nasogastric tube tip placed in the pouch over the esophageal stethoscope cuff.

560 First Avenue
New York, New York 10016

(Accepted for publication June 28, 1979.)

Anesthesiology
51:568-569, 1979

Body Weight vs. Surface Area for Calculating Dose of Spinal Anesthetic

To the Editor:—I cannot agree with the conclusion of Dohi *et al.*¹ that the dosage of tetracaine for children is best calculated on the basis of body weight. Their

data clearly show that the time to recovery of motor function is directly related to the total doses of tetracaine. The 0.3-mg/kg dose administered to their