

RICHARD F. GARGIULO, M.D.  
Staff Anesthesiologist  
Memorial Hospital of Burlington County  
175 Madison Avenue  
Mount Holly, New Jersey 08060

## REFERENCE

1. Ford SR, Forrest WH, Eltherington L: The treatment of reflex sympathetic dystrophy with intravenous regional bretylium. *ANESTHESIOLOGY* 68:137-140, 1988

(Accepted for publication March 30, 1988.)

Anesthesiology  
69:148, 1988

## Detecting Cyanosis in Children

*To the Editor:*—The effect of the hemoglobin concentration on the ability to detect cyanosis was not mentioned by Coté *et al.*<sup>1</sup> in their article describing the efficacy of pulse oximetry in children. In that study, the majority of clinicians failed to detect cyanosis before it was detected by the pulse oximeter.

Detection of cyanosis depends on the absolute, rather than the relative, amount of reduced hemoglobin. Cyanosis becomes apparent when the mean capillary concentration of reduced hemoglobin exceeds 5 g/dl. Thus, in a patient with a hemoglobin concentration of 10 g/dl, there would have to be 50% reduced hemoglobin for the detection of cyanosis, whereas a patient with a hemoglobin concentration of 15 g/dl would have 33% reduced hemoglobin (oxygen saturation of 67%) before cyanosis could be detected.

Although Coté *et al.* do not list the hemoglobin concentrations of the children in the study, it is likely that they ranged from 10–15 g/dl. Therefore, it is not surprising that the clinicians participating in the study were

not able to detect cyanosis with the levels of oxygen saturation described.

Fortunately, except for atelectasis, the cited causes for the arterial oxygen desaturation (airway obstruction, laryngospasm, hypoventilation, and endobronchial intubation) are all likely to be detected by an astute clinician before hemoglobin oxygen saturation reaches the low levels needed to produce detectable cyanosis. The pulse oximeter, as proven by the study, is an invaluable adjunct to the skills of the clinician.

BETTY H. JOHNSON, M.D.  
Department of Anesthesiology  
New Hanover Memorial Hospital  
Wilmington, North Carolina 28403

## REFERENCE

1. Coté CJ, Goldstein EA, Coté MA, Hoaglin DC, Ryan JF: A single-blind study of pulse oximetry in children. *ANESTHESIOLOGY* 68:184-188, 1988

(Accepted for publication March 30, 1988.)

Anesthesiology  
69:148-149, 1988

*In Reply:*—Dr. Johnson's comments regarding hemoglobin concentration, oxygen saturation, and detection of cyanosis are appreciated and technically correct, but miss the point. Our study demonstrated the efficacy of a new technology (pulse oximetry) in diagnosing arterial desaturation prior to the *late* signs of desaturation (cyanosis and bradycardia). The early studies of cyanosis *versus* hemoglobin saturation found that 4–6 g/dl of desaturated hemoglobin are required to produce cyanosis. The amount of circulating hemoglobin is the major determinant of visible cyanosis, *i.e.*, one needs to have a hemoglobin above 5 g/dl.<sup>1-4</sup> There is no question, therefore, that the pulse oximeter is far superior and much more reliable than the human eye and, as Dr. Johnson correctly points out, able to detect desaturation prior to clinically apparent cyanosis.

The clinical ability to diagnose cyanosis is also modified by many factors, including thickness of the epi-

dermis, skin pigment, and pigment associated with jaundice or Addison's disease.<sup>1-5</sup> Additionally, it has been well established that other factors, including lighting conditions and variability among individual observers, affect clinical diagnosis of cyanosis.<sup>6</sup> Indeed, Comroe and Botelho, in a large, well-controlled study, found that "visual impressions of cyanosis are unreliable. Serious grades of arterial anoxemia may be unrecognized by many physicians."<sup>6</sup>

We did not report hemoglobin concentration because we did not regard it as a contributory factor; serious life-threatening desaturation was diagnosed sooner by oximetry than by the clinician on 17 occasions. The hemoglobin values in these children ranged from 10 to 20 g/dl. In eight patients, cyanosis was not diagnosed, despite oxygen saturations ranging from 44% to non-recordable. Several of these events occurred in newborn infants with hemoglobin values between 16 and 20