

may treat or prevent acute hypotonic hyponatremia and is more important in these patients than is the justification for seeing urine appear. In this regard, proposing a new classification in which ADH excess is listed as a prerenal cause of oliguria may lead to a further misconception that these normovolemic patients also need volume repletion. In patients with nonosmotic secretion of ADH, therapy should be directed toward the signs and symptoms of hypotonic hyponatremia rather than toward a transient reduction in urine output.

DAVID M. ROTHENBERG, M.D.
Assistant Professor
Rush-Presbyterian—St. Luke's Medical Center
Rush Medical College
1653 West Congress Parkway
Chicago, Illinois 60612

Anesthesiology
73:584, 1990

Excess ADH and Oliguria in Patients with Normal Renal Function: II.

To the Editor:—The article by Zaloga and Hughes provides interesting data from patients with normal renal function who had undergone surgery and were oliguric in the postoperative period.¹ In order to better understand this population of patients, it is important to know whether their lungs were being mechanically ventilated in the intensive care unit during the measurements of renal function, since this therapy may have had a significant effect on renal function.²

The authors conclude that antidiuretic hormone (ADH) excess was responsible for the oliguria in these critically ill patients, since plasma ADH concentrations were increased over normal in hydrated patients. Could it be possible that the ADH concentration was increased in these oliguric patients but had no relevance to the changes in renal function? In order to establish a role for ADH, it is necessary to measure plasma ADH concentrations in postoperative, critically ill patients who are not oliguric to compare to the values from the groups in Zaloga and Hughes's study. In addition, the classic effect of ADH is to produce a negative free water clearance, a parameter that was not calculated for the study patients.

ADH concentrations may be increased as a result of many stimuli, but this hormone may not always have an effect on renal function.

Anesthesiology
73:584–585, 1990

In Reply:—Most nephrology textbooks define oliguria as a urine output of less than 500 ml per 24-h period. Although this definition is satisfactory for outpatients, it is unsatisfactory for critically ill inpatients. Renal perfusion may change rapidly in critically ill patients, and generally it is believed that diminished renal perfusion (manifested by oliguria) in the setting of nephrotoxic drugs, sepsis, or other conditions common in these patients may lead to renal failure. In addition, urine output is frequently used as a monitor of other organ perfusion (*i.e.*, liver and gut). Thus, urine output is usually measured on an hourly basis in critically ill patients, and most textbooks of critical care and anesthesiology define oliguria as a urine output less than $0.5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ (less restrictive than the value we used). We believe that hourly monitoring of urine output is essential to the detection of renal

ARNOLD S. BERNIS, M.D.
Director of Nephrology
St. Francis Hospital
355 Ridge Avenue
Evanston, Illinois 60202

REFERENCES

1. Zaloga GP, Hughes SS: Oliguria in patients with normal renal function. *ANESTHESIOLOGY* 72:598–603, 1990
2. Arieff AI: Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *New Engl J Med* 314:1529–1535, 1986

(Accepted for publication June 25, 1990.)

Hemodynamic changes, renal blood flow, sympathetic stimulation, or other hormones may have been responsible for the oliguria noted in this sample of postoperative patients. Without the appropriate control group, interpretation of the ADH data is impossible.

ARNOLD J. BERRY, M.D.
Associate Professor of Anesthesiology
Department of Anesthesiology
Emory University School of Medicine
1364 Clifton Road, N.E.
Atlanta, Georgia 30322

REFERENCES

1. Zaloga GP, Hughes SS: Oliguria in patients with normal renal function. *ANESTHESIOLOGY* 72:598–603, 1990
2. Berry AJ: Respiratory support and renal function. *ANESTHESIOLOGY* 55:655–667, 1981

(Accepted for publication June 25, 1990.)

hypoperfusion before permanent renal damage occurs. Since hourly monitoring of urine output is the standard of practice in critical care and anesthesia, we chose this method of monitoring to define oliguria.

We agree that low urine output in a patient with clinical features of hypovolemia does not require urinary indices to aid in a diagnosis of volume depletion. On the other hand, in many patients clinical features are not conclusive, and in these patients, we believe, urinary indices may be helpful. A major point of our study¹ was the demonstration that patients with low urine output and a physical exam that does not indicate hypovolemia frequently have hormonal-excess-associated oliguria and do well without volume administration.

Normovolemic patients with oliguria and normal renal function do not require treatment with fluids. In fact, large volume infusion in