REFERENCES

- Kamaya H, Krishna P: New endotracheal tube (Univent Tube®) for selective blockade of one lung. ANESTHESIOLOGY 63:342– 343, 1985
- Karwande S: A new tube for single lung ventilation. Chest 92: 761-763, 1987
- 3. Steen J: Impact of tube design and materials on complications of
- tracheal intubation, Problems in Anesthesia, Volume 2. Edited by Kirby R, Brown D. Philadelphia, JB Lippincott, 1988, pp 211-224
- Jesseph J, Merendino K: The dimensional interrelationships of the major components of the human tracheobronchial tree. Surg Gynecol Obstet 105:210-214, 1957

(Accepted for publication April 8, 1991.)

Anesthesiology 75:166, 1991

Proper Needle Size for Performing Cervical Epidural Injections

To the Editor:—The case report by Jackson and Rauck¹ describing possible air embolism complicating a therapeutic cervical epidural steroid injection in a sitting patient deserves comment.

Anesthesiologists must clearly differentiate the needles and techniques required for single-shot epidural injections from those used with continuous catheter epidural injections. For therapy in our Pain Clinic, unless a catheter is used in an attempt at breaking up epidural scarring, we cannot justify using a needle larger than 0.7 mm (22-G). With a loaded syringe attached, this needle is inserted through an introducer to identify and enter the epidural space. In the case described, the authors used a spade-tipped, 16-G Tuohy needle. With that needle, the tissue area vulnerable to injury is almost six times as large as with our technique (2.26 vs. 0.385 mm²). Thus, sequelae such as backache, headache from inadvertent dural puncture, and injuries to nerves, arteries, veins and other tissues will be more frequent and severe.

Our personal experience with thousands of epidural injections of all types during the past 45 yr indicates that therapeutic epidural steroid injections properly performed and instrumented to match the requirements should be relatively benign procedures. *Primun non nocere!*

BERNARD S. GOFFEN, M.D. Director, Pain Clinic Department of Veterans Affairs Medical Center Salem, Virginia 24153

REFERENCE

 Jackson KE, Rauck RL: Suspected venous air embolism during epidural anesthesia. ANESTHESIOLOGY 74:190-191, 1991

(Accepted for publication April 10, 1991.)

Anesthesiology 75:166, 1991

In Reply:—We agree with Dr. Goffen that larger needles generally are more likely to cause greater injury to delicate structures such as arteries and nerves. However, for the following reasons we believe that 16-G Tuohy needles are appropriate for cervical epidural placement. Using the "hanging-drop technique" to identify the epidural space requires reliable transmission of the negative pressure through the needle to the hub. Increasing resistance through a needle's progressively smaller internal diameter will result in a loss of the sensitivity seen when the drop is drawn into the needle hub. It is our experience that using 18-G Tuohy needles results in less sensitivity than with 16-G needles, and thus it is our conclusion that using 22-G needles are even less useful.

Although it is certainly possible to identify the epidural space through the loss of resistance technique by using 22-G needles, this can be a source of complications. Also, to the best of our knowledge, a 22-G Tuohy needle is not available.

Certainly, an anesthetic procedure can result in acute complications. However, cervical epidural anesthesia is well documented in terms of overall safety, and we still believe that the technique of this procedure is easiest in the sitting position with at least a 16-18-G Tuohy needle.

KYLE JACKSON, M.D.
RICHARD L. RAUCK, M.D.
Director Pain Control Center
The Department of Anesthesia
The Bowman Gray School of Medicine
300 South Hawthorne Road
Winston-Salem, North Carolina 27103

REFERENCE

 Waldman SD: Complications of cervical epidural nerve blocks with steroids: A prospective study of 790 consecutive blocks. Reg Anesth 14:149-151, 1989

(Accepted for publication April 10, 1991.)