

CORRESPONDENCE

References

1. Sakura S, Sumi M, Sakaguchi Y, Saito Y, Kosaka Y, Drasner K: The addition of phenylephrine contributes to the development of transient neurologic symptoms after spinal anesthesia with 0.5% tetracaine. *ANESTHESIOLOGY* 1997; 87:771-8
2. Wang BC, Hillman DE, Spielholz NI, Turndorf H: Chronic neuro-

logical deficits and nesacaine-CE—an effect of the anesthetic, 2-chloroprocaine, or the antioxidant, sodium bisulfite? *Anesth Analg* 1984; 63:445-7

3. Hersh EV, Condouris GA, Havelin D: Actions of intrathecal chloroprocaine and sodium bisulfite on rat spinal reflex function utilizing a noninvasive technique. *ANESTHESIOLOGY* 1990; 72:1077-82

(Accepted for publication March 20, 1998.)

Anesthesiology
1998; 89:273

© 1998 American Society of Anesthesiologists, Inc.
Lippincott-Raven Publishers

Transient Neurologic Symptoms when Phenylephrine Is Added to Tetracaine Spinal Anesthesia—An Alternative

To the Editor:—The article by Sakura *et al.* regarding transient neurologic symptoms after tetracaine spinal anesthesia with phenylephrine was informative.¹ Why do anesthesiologists add a vasoconstrictor to prolong the duration of the anesthesia? As the authors indicate, "Prolongation is thought to result, at least in part, from a decrease in nerve blood flow resulting in reduced vascular uptake of the local anesthetic."² Thus, more local anesthetic is available for neuronal penetration, causing a more profound and prolonged block. It also is implied that this allows the anesthesiologist to limit the amount of local anesthetic injected, in turn decreasing cephalad spread. But if the vascular absorption of the local anesthetic decreases, is not that the equivalent of injecting more local anesthetic? Because there is an apparent adverse effect of adding phenylephrine to tetracaine spinal anesthesia,¹ why not just inject a little more tetracaine to prolong the anesthesia and omit the phenylephrine?

The Editorial View that accompanied the Sakura paper was confusing.³ Dr. Rowlingson agrees that Sakura *et al.*¹ have furthered our knowledge regarding transient neurologic symptoms after spinal anesthesia. He then goes a step further and states, "Perhaps tetracaine is a better (safer) drug for spinal anesthesia, because compared to lidocaine and bupivacaine it increases spinal cord blood flow." That conclusion, based on one dog study,² is a big leap. Tetracaine, like lidocaine, caused cauda equina syndrome after continuous spinal anesthesia,⁴⁻⁶ and it is neurotoxic to isolated nerve.⁷ Further, compared with bupivacaine, tetracaine is associated with a significantly higher incidence of tourniquet pain during lower extremity orthopedic operations.⁸ It is for these reasons that I do not share or understand Dr. Rowlingson's enthusiasm for tetracaine spinal anesthesia.

Donald H. Lambert, Ph.D., M.D.
Chief, Anesthesiology
Associate Chief, Surgical Services
Brockton-West Roxbury VAMC

West Roxbury, Massachusetts
Associate Professor of Anaesthesia
Harvard Medical School
Boston, Massachusetts

References

1. Sakura S, Sumi M, Sakaguchi Y, Kosaka Y, Drasner K: The addition of phenylephrine contributes to the development of transient neurologic symptoms after spinal anesthesia with 0.5% tetracaine. *ANESTHESIOLOGY* 1997; 87:771-8
2. Kozody R, Palahniuk RJ, Cumming MO: Spinal cord blood flow following subarachnoid tetracaine. *Can Anaesth Soc J* 1985; 32:23-9
3. Rowlingson JC: Transient neurologic symptoms. Now, with phenylephrine? *ANESTHESIOLOGY* 1997; 87:737-8
4. Nicholson M, Eversole U: Neurological complications of spinal anesthesia. *JAMA* 1946; 132:679-85
5. Rigler ML, Drasner K, Krejcie TC, Yelich SJ, Scholnick FT, De-Fontes J, Bohner D: Cauda equina syndrome after continuous spinal anesthesia. *Anesth Analg* 1991; 72:275-81
6. Schell RM, Brauer FS, Cole DJ, Applegate RL II: Persistent sacral nerve root deficits after continuous spinal anaesthesia. *Can J Anaesth* 1991; 38:908-11
7. Lambert LA, Lambert DH, Strichartz GR: Irreversible conduction block in isolated nerve by high concentrations of local anesthetics. *ANESTHESIOLOGY* 1994; 80:1082-93
8. Concepcion MA, Lambert DH, Welch KA, Covino BG: Tourniquet pain during spinal anesthesia: A comparison of plain solutions of tetracaine and bupivacaine. *Anesth Analg* 1988; 67:828-32

(Accepted for publication March 20, 1998.)