

- A-97** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Sevoflurane Anesthesia Does Not Induce Sister Chromatid Exchanges in Lymphocytes of Children Thorsten Krause, M.D.; Jens Scholz, M.D.; Lars Jansen, M.D.; Claudia Koch, M.D.; Jochen Schulte am Esch, M.D., *Anesthesiology, University-Hospital Eppendorf, Hamburg, Germany*. Degradation products of sevoflurane produce genotoxicity in vitro. This was not confirmed in vivo after sevoflurane anesthesia.
- A-98** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
The Role of Nitric Oxide in Isoflurane-Mediated Vasodilation in Humans Ali Mchaourab, MD; Thomas J. Ebert, MD, PhD, *Anesthesiology, Medical College of Wisconsin, Milwaukee, WI, United States*. Nitric oxide plays an important role in the vasodilation of vessels during isoflurane anesthesia, however whether it is active or permissive is not yet known.
- A-99** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Induction of Heme Oxygenase-1 Minimizes Venular Leukocyte-Endothelial Interaction Elicited by Halothane Anesthesia In Vivo Hiroshi Morisaki, MD; Tomihiro Katayama, MD; Yosbifumi Kotake, MD; Junzo Takeda, MD; Makoto Suematsu, MD, *Anesthesiology, Keio University School of Medicine, Tokyo, Japan*. Induction of heme oxygenase-1 modified halothane-induced leukocyte adhesion in vivo.
- A-100** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Intraoperative Morphine and Early Recovery after Remifentanyl-Sevoflurane-N₂O Anesthesia Hernan R. Munoz, MD; Mario E. Guerrero, MD; Verena Brandes, MD, *Department of Anesthesiology, Catholic University, Santiago, Chile*. The intraoperative administration of morphine during a remifentanyl-based anesthesia did not prolong the early recovery period compared to placebo.
- A-101** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Sevoflurane-Vaporized Cardioplegia Improves Myocardial Wall Motion after Coronary Revascularization Nader D. Nader, M.D., Ph.D.; Carlos M. Li, M.D.; Wiam Z. Kbadra, PhD; Anthony L. Panos, M.D.; Paul R. Knight, M.D., Ph.D., *Anesthesiology, University of Arkansas, Little Rock, AR, United States*. Vaporizing cardioplegia solution with sevoflurane decreased wall motion abnormality after coronary revascularization.
- A-102** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Minimum Alveolar Concentration (MAC) of Xenon with Sevoflurane in Humans Yosbinori Nakata, MD, MBA; Takahisa Goto, MD; Yosbiki Isbiguro, MD; Yosbinari Niimi, MD; Shigebo Morita, MD, *Anesthesia, Teikyo University Ichihara Hospital, Ichihara, Chiba, Japan*. The MAC values of xenon and sevoflurane are 63.1% and 1.74%, respectively.
- A-103** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Deletion in RyR1 Gene Associated with MH and with Abnormal Electrically Evoked Twitch Tension In Vitro Thomas E. Nelson, Ph.D.; Nyamkhisg Sambuughin, Ph.D., *Dept. of Anesthesiology, Wake Forest University School of Medicine, Winston-Salem, NC, United States*. A novel 3-base deletion in the RyR1 gene links to MH susceptibility and unusual contraction phenotype in 2 affected families.
- A-104** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Verapamil and Nifedipine Differentially Modulate the Effect of Sevoflurane on Cardiac L-Type Calcium Channel Hirotsugu Okamoto, M.D., Ph.D.; Anthony J. Mazzeo, M.D.; Zeljko J. Bosnjak, Ph.D.; Wai-Meng Kwok, Ph.D., *Anesthesiology, Medical College of Wisconsin, Milwaukee, WI, United States*. Verapamil and nifedipine may interact with sevoflurane at distinct sites on cardiac L-type calcium channel.
- A-105** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Effect of Halothane on IP₃ and Ryanodine Receptor Calcium Release Channels in Airway Smooth Muscle Christina M. Pabellick, M.D.; Y.S. Prakash, Ph.D.; Gary C. Steck, Ph.D., *Department of Anesthesiology, Mayo Clinic, Rochester, MN, United States*. Halothane depletes Ca²⁺ stores in airway smooth muscle by inducing leak through IP₃ and ryanodine receptor channels.
- A-106** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Intravenous Isoflurane Lipid Emulsion is More of a Myocardial Depressant Than Inhaled Isoflurane Christian Popa, MD; John L. Fontana, MD; Paul D. Mongan, MD, *Anesthesiology, Walter Reed Army Medical Center, Washington, DC, United States*. We compared the hemodynamic effects of IV and inhaled isoflurane and found similar HR and BP but greater myocardial depression using the IV route.
- A-107** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
"Paradoxical Arousal" during Bispectral Index Monitoring Ira J. Rampil, MS, MD; Hongmei Cai, PhD; Patricia B. Embree, RN; Scott D. Greenwald, PhD, *Anesthesia, University of California, San Francisco, CA, United States*. Paradoxical arousal (PA) slows EEG; we sought its prevalence. In 1901 anesthetized EEGs, we found 37 PA; 10x more frequent in vapor than iv anesthetics, 4x more common under 18 yr.
- A-108** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Optimization of Maintenance Concentrations of Volatile Anesthetics, Propofol, and Opioids: Costs and Awakening Times Heiko Roepcke, M.D.; Thomas Bouillon, M.D.; Joergen Bruhn, M.D., *Dep. of Anesthesiology, University of Bonn, Bonn, Germany*. Pharmacokinetic and -dynamic models were used to identify optimal anesthetic drug combinations with respect to minimal costs or awakening times.
- A-109** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Hemodynamic Responses to Isoflurane, Desflurane, and Sevoflurane in Rats Following Chronic Morphine or Cocaine Administration R. Sabar; A.D. Kaye; S. Vig; C. Ricaldi; J.E. Heavner, *Anesthesiology, Physiology, Pharmacology, Texas Tech University, Lubbock, TX, United States*. SBP response in rats exposed chronically to morphine or cocaine is dependent on anesthetic and its concentration.
- A-110** Room E, 10/17/2000 2:00 PM - 4:00 PM (PS)
Sevoflurane Does Not Affect Platelet Function in Coronary Artery Bypass Graft Hiroshi Sakamoto, M.D.; Takahisa Mayumi, M.D.; Osamu Kemmotsu, M.D., *Department of Anesthesia, Hokkaido OHNO Hospital, Sapporo, Hokkaido, Japan*. Sevoflurane does not affect platelet function and onset time and clot rate by Sono clot[®] compared with propofol in coronary artery bypass graft.