

## ASA ABSTRACTS

Anesthesiology  
V93, No 3A, Sep 2000

A-357 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Redistribution Hypothermia Correlates with the Threshold for Thermoregulatory Vasoconstriction in Patients Anesthetized with Isoflurane or Xenon but Not Nitrous Oxide** *Takahisa Goto, MD; Takashi Matsukawa, MD; Daniel I. Sessler, MD; Makoto Ozaki, MD; Shigeo Morita, MD, Teikyo University Ichibara Hospital, Ichibara-shi, Japan*

A-358 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Thrombin Reduces Fluid Leak Through Punctured Dura Mater in an In Vitro Model** *Dario A. Grisales, MD; Dan Paoli, MD; Conny Frostb, MD; Rebana Nawab, MD, Anesthesiology, University of South Florida College of Medicine, Tampa, FL, United States.* Thrombin patch significantly decreases a durotomy fluid leak at hydrostatic pressure up to 30 cmH<sub>2</sub>O.

A-359 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Increased Endothelial Cell Turnover in Human Cerebral Arteriovenous Malformations** *Tomoki Hashimoto, MD; Ricardo Mesa-Tejada, MD; Christopher M. Quick, PhD; Andrew W. Bollen, MD, DVM; William L. Young, MD, Departments of Anesthesia, Neurology, Neurosurgery, and Pathology, University of California, San Francisco, San Francisco, CA, United States.* Mitogenesis (Ki-67) is greater in AVMs than normal brain endothelium.

A-360 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Abnormal Expression of Angiopoietin-2 and Tie-2 in Human Cerebral Arteriovenous Malformations** *Tomoki Hashimoto, MD; Charles W. Emala, MD; Nancy J. Boudreau, PhD; Christopher M. Quick, PhD; William L. Young, MD, Dpts. of Anesthesia, Neurology, Neurosurgery and Surgery, University of California, San Francisco, CA, United States.* Angiopoietin-2 and Tie-2 expression were abnormal in cerebral AVMs.

A-361 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Propofol Versus Sevoflurane Anesthesia: Influence on Cerebral and Aortic Blood Flow Velocities** *Andrea Holzer, MD; Josef Stark, MD; Manfred Greber, MD; Andrew Donner, MD; Udo M. Illieich, MD, Anesthesiology and General Intensive Care, University of Vienna, Vienna, Austria.* We compared cerebral and systemic blood flow velocities in awake and anesthetized (propofol-sevoflurane) patients.

A-362 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**In Non-Human Primates Intracarotid Adenosine -Not Nitroprusside- Profoundly Increases Cerebral Blood Flow** *Shailendra Joshi, MD; Beverly Aagaard, MD; John Pile-Spellman, MD; Adam Libow, BA; Sulli J. Popilskis, DVM, Anesthesiology, Columbia University, New York, NY.* Intracarotid adenosine -not nitroprusside- in doses which lack systemic side-effects, profoundly increases CBF of baboons.

A-363 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**In Humans and Non-Human Primates Intracarotid Nitroprusside Does Not Augment Cerebral Blood Flow** *Shailendra Joshi, MD; William L. Young, MD; John Pile-Spellman, MD; Hoang Duong, MD; Beverly Aagaard, MD, Anesthesiology, Columbia University, New York, NY.* Intracarotid nitroprusside, an endothelium independent nitric-oxide donor, failed to augment cerebral blood flow in humans and baboons.

A-364 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Transcranial Doppler Ultrasonography with Induction of Anesthesia and Neuromuscular Blockade in Surgical Patients** *W. Andrew Kofke, MD; Julie McWhorter, BS; Natalie Shaheen, BS; Elizabeth Sinz, MD, Anesthesiology, West Virginia University, Morgantown, WV, United States*

A-365 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Pre-Incubation of Vascular Rings with Iohexol Does Not Affect cGMP Response to Nitroprusside** *Adam D. Libow, BA; Shailendra Joshi, MD; Lena S. Sun, MD; Carol A. Hirszman, MD; William Mack, BA, Anesthesiology, Columbia University, New York, NY.* Compared to equiosmotic concentrations of mannitol, iohexol did not impair cGMP generation in arterial rings after nitroprusside exposure.

A-366 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Influence of Early Factors on Long-Term Head Injury Outcome** *Colin Mackenzie; Shiu Ho; Nafeesa Jawed; Pat Dischinger; Michael Makely, National Study Center for Trauma & EMS and Shock Trauma Center, University of MD, Baltimore, MD, United States.* Traumatic brain injury patients requiring pre-hospital intubation had the same Glasgow Coma Scale (GCS) as non-intubated patients on discharge at 33 days despite their greater admission Injury Severity Score (ISS) and lower admission GCS. Postive admission blood alcohol had no effect on outcome status.

A-367 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Buspirone and Meperidine Synergistically Reduce the Shivering Threshold** *Masoud Mokhtarani, MD; Adel N. Mahgob, MD; Nobu Morioka, MD; Anthony Doufas, MD; Daniel I. Sessler, MD, Outcomes Research, Dept. of Anesthesia, UCSF, San Francisco, CA, United States.* Buspirone and meperidine synergistically reduce the shivering threshold to 33°C while causing little sedation or respiratory toxicity.

A-368 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Remifentanil Versus Alfentanil: Effect on Systemic Circulation and Cerebral Blood Flow in Neurosurgical Patients in the Sitting Position** *Bart M.B. Monteyne, MD; Jozef Van Aken, PhD; Michel M.R.F. Struys, PhD; Jacques Caemaert, PhD; Gaspard P.F. De Ley, PhD, Anesthesia, University Hospital, Gent, Belgium.* Remifentanil is an alternative to alfentanil to maintain CBF in the sitting position.

A-369 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**A Large Dose of Amino Acids Prevent Core Hypothermia Even during Abdominal Surgery** *Chibaru Negishi, MD; Daniel I. Sessler, MD; Kenji Atarashi, MD; Takashi Matsukawa, MD; Hidehiro Suzuki, MD, Anesthesiology, Tokyo Women's Medical University, Shinjuku, Tokyo, Japan.* Amino acids prevent most core hypothermia. Prevention was dose-dependent, with 0.4 and 0.6 g/kg/h providing most benefit.

A-370 Room C, 10/17/2000 9:00 AM - 11:00 AM (PS)

**Effect of Therapeutic Blood-Brain Barrier Disruption on Cerebrospinal Fluid Homeostasis** *Misha Perouansky, MD; Lev Ronin, MD; Ariele Eden-Openheim; Tal Siegal; Yoram G. Weiss, Anesthesiology, "Hadassah" Medical Center - Hebrew University School of Medicine, Jerusalem, Israel.* Mechanisms other than the BBB contribute to maintaining CSF/Blood gradients.