

Differential Effects of Volatile Anesthetics on Hepatic Heme Oxygenase-1 Expression in the Rat

Alexander Hoetzel, M.D.,* Sarah Geiger,† Torsten Loop, M.D.,* Armin Welle,† René Schmidt, M.D.,* Matjaz Humar, Ph.D.,* Heike L. Pahl, Ph.D.,‡ Klaus K. Geiger, M.D.,§ Benedikt H. J. Pannen, M.D.||

THE microsomal enzyme heme oxygenase (HO; EC 1.14.99.3) catalyzes the oxidation of heme to biliverdin-IXa, iron, and carbon monoxide.¹ So far, three isoforms of this enzyme have been cloned. While HO-2 and HO-3 are constitutively expressed,^{2,3} HO-1 is highly inducible in response to a variety of stimuli and has been identified as the major 32-kd heat shock (stress) protein (HSP) 32.⁴

Heme oxygenase-1 is essential for the function of the normal liver and plays a major protective role in the stress-exposed liver.⁵ It exerts antioxidant properties by cleavage of the prooxidant heme and by the production of biliverdin, which is subsequently reduced to the antioxidant bilirubin.⁶ Furthermore, its product carbon monoxide serves to maintain liver perfusion⁵ and has potent antiinflammatory effects.⁷

Many studies have shown differential effects of volatile anesthetics on the perfusion, function, and integrity of the liver. Moreover, preliminary evidence suggests that volatile anesthetics may interfere with stress gene expression and function.⁸ We therefore aimed to determine whether volatile anesthetics affect hepatic HO-1 gene expression.

Materials and Methods

Animals

The experimental protocol was approved by the local animal care and use committee, and all animals received humane care according to the criteria outlined in the *Guide for the Care and Use of Laboratory Animals*.⁹ Male Sprague-Dawley rats (Charles River, Sulzfeld, Germany) weighing 299 ± 16 g were subjected to arterial blood pressure monitoring *via* the carotid artery, and a tracheotomy was performed to maintain airway patency.¹⁰ The animals were randomized into one of the following groups ($n = 6$ for each group): group 1 (control), anesthesia with 60 mg/kg pentobarbital sodium

intraperitoneally (Alvetra, Neumuenster, Germany) followed by intravenous bolus injections of 6 mg/kg every 30 min; group 2, isoflurane (2.3 vol%; Baxter, Unterschleissheim, Germany); group 3, desflurane (12 vol%; Baxter, Unterschleissheim, Germany); or group 4, sevoflurane (4 vol%; Abbott, Wiesbaden, Germany).

These concentrations of volatile anesthetics correspond to a minimum alveolar concentration in the rat of 1.6–1.8.^{11–13} Volatile anesthetics were delivered in an atmosphere of 30% oxygen (Narcorex19; Draeger, Luebeck, Germany), continuously monitored (PM8050; Draeger, Luebeck, Germany), and spontaneously inhaled by the animals through a tracheotomy tube for the experiment's duration of 6 h. Body temperature was maintained at $38 \pm 4^\circ\text{C}$. This regimen resulted in an arterial oxygen tension of 60–100 mmHg and an arterial carbon dioxide tension of 30–50 mmHg in all groups. Two animals that served as positive controls for the detection of HSP27 and HSP70 were subjected to 1 h of hemorrhagic shock and 5 h of resuscitation as previously described.¹⁰ Additional experiments were performed to mimic volatile anesthetic-induced hypotension in rats anesthetized with pentobarbital using the vasodilator dihydralazine ($n = 4$; Novartis, Nuernberg, Germany). The dose of dihydralazine ranged from 4 to $8 \text{ mg} \cdot \text{kg}^{-1} \cdot 6 \text{ h}^{-1}$ and was adjusted to achieve a mean arterial pressure (MAP) of 76 ± 12 mmHg, which was the mean MAP of all animals subjected to volatile anesthetics.

RNA Extraction and Northern Blot Analysis

Total RNA was isolated from frozen liver tissue (TRIZOL; Gibco, Grand Island, NY) as previously described.¹⁰ Ten micrograms of RNA were loaded per lane, size fractionated, and transferred to a nylon membrane. Hybridization was performed with P³² γ -deoxycytosine triphosphate–radiolabeled HO-1,¹⁴ HSP27, and HSP70 complementary DNA.¹⁰ All blots were reprobated with 18S ribosomal RNA complementary DNA to verify equal loading. Autoradiographs were analyzed by laser scanning densitometry (Personal Densitometer; Molecular Dynamics, Krefeld, Germany).

Western Blot Analysis

Frozen liver tissue was homogenized and lysed in TOTEX buffer (20 mM HEPES [pH 7.9], 0.35 M NaCl, 20% glycerol, 1% Nonidet P-40, 1 mM MgCl₂, 0.5 mM EDTA, 0.1 mM EGTA) as previously described.¹⁰ Each lane of a

* Postdoctoral Research Fellow, § Chairman, Professor, || Associate Professor, Department of Anesthesiology and Critical Care Medicine, † Research Assistant, Student, ‡ Professor, Department of Experimental Anesthesiology, University of Freiburg.

Received from the Department of Anesthesiology and Critical Care Medicine, University of Freiburg, Germany. Submitted for publication January 14, 2002. Accepted for publication May 30, 2002. Supported by the Deutsche Forschungsgemeinschaft, Bonn, Germany (grants DFG PA 533/2-3 and 533/4-1; Heisenberg-Stipend PA 533/3-1 to B. H. J. Pannen).

Address reprint requests to Dr. Pannen: Anaesthesiologische Universitätsklinik, Hugstetterstrasse 55, D-79106 Freiburg, Germany. Address electronic mail to: pannen@nz.ukl.uni-freiburg.de. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.




Fig. 1. Expression of heme oxygenase-1 (HO-1) messenger RNA (mRNA) (A), 18S ribosomal RNA (B), or HO-1 protein (C) in the livers of two representative animals subjected to 6 h of anesthesia with pentobarbital sodium (C), isoflurane (I; 2.3 vol%), desflurane (D; 12 vol%), or sevoflurane (S; 4 vol%). (D) Relative densitometric units of HO-1 mRNA concentrations. Data are presented as median (box: 25th and 75th percentiles; error bars: 10th and 90th percentiles) for $n = 6$ animals per group. * $P < 0.05$ versus pentobarbital control (C). # $P < 0.05$ versus desflurane (D).

12% sodium dodecyl sulphate polyacrylamide gel contained 100 μ g of total protein. After protein separation and electroblotting, HO-1 was detected by a rabbit polyclonal anti-HO-1 antibody (1:1,000 dilution, SPA-895; StressGen Biotechnologies, Victoria, Canada) using the Enhanced Chemiluminescence (ECL) detection kit (Amersham Pharmacia, Freiburg, Germany) according to the manufacturer's instructions.

Data Analysis

Data are presented as median (box: 25th and 75th percentiles; error bars: 10th and 90th percentiles). Statistical differences between experimental groups were determined using a Kruskal-Wallis one-way analysis of

variance on ranks followed by a *post hoc* Student-Newman-Keuls test. A P value < 0.05 was considered to indicate a significant difference.

Results

Effect of Volatile Anesthetics on Heme Oxygenase-1 Gene Expression in the Liver

Hepatic HO-1 transcripts were barely detectable in the animals anesthetized with pentobarbital. In contrast, 6 h of anesthesia with isoflurane led to a substantial increase in HO-1 messenger RNA (mRNA) and protein (fig. 1). Densitometric analysis of all experiments revealed a more than 2.3-fold induction of HO-1 mRNA by isoflurane as compared to the pentobarbital control experiments (fig. 1).

Anesthesia with sevoflurane led to a 3.3-fold increase in HO-1 mRNA that tended to be even more pronounced as compared to isoflurane (fig. 1). In sharp contrast to the inducing effect of isoflurane and sevoflurane, desflurane did not upregulate HO-1—*i.e.*, the hepatic HO-1 mRNA content was comparable to that of pentobarbital controls (fig. 1). Western blot analyses revealed a similar pattern of HO-1 protein accumulation in response to the different anesthetic agents (fig. 1).

Effect of Volatile Anesthetics on Heat Shock Protein 27 and Heat Shock Protein 70 Gene Expression in the Liver

To test whether volatile anesthetics lead to a general induction of stress-inducible HSPs, the same RNA samples as shown in figure 1 were hybridized with radiolabeled HSP27 or HSP70 complementary DNA probes. No signal was detectable under control, isoflurane, desflurane, or sevoflurane anesthesia. To exclude the possibility that this was due to technical reasons, the RNA of two animals that had been subjected to hemorrhagic shock




Fig. 2. Expression of heat shock protein (HSP) 27 messenger RNA (mRNA) (A), HSP70 mRNA (B), and 18S ribosomal RNA (C) in the livers of two representative animals subjected to 6 h of anesthesia with pentobarbital sodium (C), isoflurane (I; 2.3 vol%), desflurane (D; 12 vol%), or sevoflurane (S; 4 vol%). + = positive control.




Fig. 3. (A) Mean arterial pressure of rats during anesthesia with pentobarbital sodium (C), isoflurane (I; 2.3 vol%), desflurane (D; 12 vol%), sevoflurane (S; 4 vol%), or pentobarbital + dihydralazine (Dihyd). Expression of heme oxygenase-1 (HO-1) messenger RNA (B) or 18S ribosomal RNA (C) in the livers of representative animals.

and resuscitation¹⁰ were used as positive controls and showed a strong signal (fig. 2).

Effect of Hypotension on Hepatic Heme Oxygenase-1 Gene Expression

The MAP was approximately 35% lower in the isoflurane, sevoflurane, and desflurane groups as compared to the pentobarbital controls (fig. 3). To evaluate the role of hypotension in hepatic HO-1 induction, additional pentobarbital control experiments were performed using the vasodilator dihydralazine to lower the MAP to the same extent (76 ± 12 mmHg) as observed in the animals treated with isoflurane, desflurane, or sevoflurane (78 ± 8 mmHg). However, none of these animals exhibited HO-1 mRNA accumulation in the liver (fig. 3).

Discussion

Our aim in this study was to evaluate the effect of volatile anesthetics on hepatic HO-1 expression. Our results indicate that different volatile anesthetics differ-

entially regulate HO-1 expression in the normal liver. Whereas isoflurane and sevoflurane induced HO-1 mRNA and protein, desflurane did not affect the expression of this gene.

It has been shown previously that halothane may cause hepatic expression of HO-1 in rats pretreated with phenobarbital if administered under hypoxic conditions.^{15,16} In contrast, neither halothane nor isoflurane induced HO-1 in the liver during hypoxia in the absence of phenobarbital pretreatment in these studies.¹⁶ This discrepancy between the historical data and the observations described here is most likely due to the fact that the total amount of volatile anesthetics administered was much lower in the former study. While Yamasaki *et al.*¹⁶ used a 0.3 minimum alveolar concentration of isoflurane, our experimental protocol included the administration of a 1.7 minimum alveolar concentration. Moreover, the time of exposure to the volatile anesthetics was limited to 2 h in the previous reports, while it was 6 h in the present study. Therefore, the sum of these data would be consistent with a dose and time dependency of the inducing effect of volatile anesthetics on hepatic HO-1 expression.

The reason for the differential induction of HO-1 by different volatile anesthetics is not known. However, it could be related to differences in the degree of hepatic cellular stress after exposure to these agents.¹⁷⁻¹⁹ Thus, we determined the effect of the different volatile anesthetics on the expression of other stress-inducible heat shock proteins, such as HSP27 and HSP70, under the same experimental conditions.^{10,15} However, neither HSP27 nor HSP70 mRNA could be detected after 6 h of anesthesia with the various agents. This is in accordance with previous reports that have demonstrated induction of HSP70 by volatile anesthetics only in combination with phenobarbital pretreatment and hypoxia but not by volatile anesthetics alone.^{15,16,20,21} These findings suggest that the signal transduction pathway responsible for volatile anesthetic-mediated HO-1 expression is independent of the one leading to the induction of the general heat shock response.

To exclude the possibility that the reduction in arterial blood pressure by volatile anesthetics might be involved in HO-1 induction, we lowered the arterial blood pressure in additional animals anesthetized with pentobarbital to a similar extent as in the volatile anesthetic groups. However, no induction of HO-1 could be observed under these conditions. Therefore, hypotension itself does not seem to be responsible for HO-1 induction. Moreover, since isoflurane, sevoflurane, and desflurane decreased the MAP to the same degree, this cannot explain their differential effects on HO-1 expression.

Taken together, our findings demonstrate that volatile anesthetics differentially induce HO-1 mRNA in the liver, and this effect does not extend to other HSPs. Therefore, volatile anesthetics can be considered as specific gene

regulators of hepatic HO-1, an enzyme that may exert potent organ protective functions.⁵

The authors are grateful to Shigeki Shibahara, M.D., Ph.D. (Chairman, Department of Applied Physiology and Molecular Biology, Tohoku University School of Medicine, Sendai, Japan), who kindly provided the HO-1 complementary DNA clone.

References

1. Tenhunen R, Marver HS, Schmid R: The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci U S A* 1968; 61:748-55
2. Cruse I, Maines MD: Evidence suggesting that the two forms of heme oxygenase are products of different genes. *J Biol Chem* 1988; 263:3348-53
3. McCoubrey WK Jr, Huang TJ, Maines MD: Isolation and characterization of a cDNA from the rat brain that encodes hemoprotein heme oxygenase-3. *Eur J Biochem* 1997; 247:725-32
4. Ewing JF, Maines MD: Rapid induction of heme oxygenase 1 mRNA and protein by hyperthermia in rat brain: heme oxygenase 2 is not a heat shock protein. *Proc Natl Acad Sci U S A* 1991; 88:5364-8
5. Pannen BHJ, Köhler N, Hole B, Bauer M, Clemens MG, Geiger KK: Protective role of endogenous carbon monoxide in hepatic microcirculatory dysfunction after hemorrhagic shock in rats. *J Clin Invest* 1998; 102:1220-8
6. Maines MD: The heme oxygenase system: A regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* 1997; 37:517-54
7. Otterbein LE, Bach FH, Alam J, Soares M, Lu HT, Wysk M, Davis RJ, Flavell RA, Choi AMK: Carbon monoxide has anti-inflammatory effects involving the mitogen-activated protein kinase pathway. *Nat Med* 2000; 6:422-8
8. Buzaleh AM, del Carmen Martinez M, del Carmen Batlle AM: Relevance of cytochrome P450 levels in the actions of enflurane and isoflurane in mice: studies on the haem pathway. *Clin Exp Pharmacol Physiol* 2000; 27:796-800
9. American Association for Laboratory Animal Science: Guide for the Care and Use of Laboratory Animals. National Institutes of Health Publication No. 86-23. Bethesda, MD, National Institutes of Health, 1985
10. Hoetzel A, Vagts DA, Loop T, Humar M, Bauer M, Pahl HL, Geiger KK, Pannen BHJ: Effect of nitric oxide on shock-induced hepatic heme oxygenase-1 expression in the rat. *Hepatology* 2001; 33:925-37
11. Antognini JF, Carstens E: Increasing isoflurane from 0.9 to 1.1 minimum alveolar concentration minimally affects dorsal horn cell responses to noxious stimulation. *ANESTHESIOLOGY* 1999; 90:208-14
12. Goren S, Kahveci N, Alkan T, Goren B, Korfali E: The effects of sevoflurane and isoflurane on intracranial pressure and cerebral perfusion pressure after diffuse brain injury in rats. *J Neurosurg Anesthesiol* 2001; 13:113-9
13. Mazze RI, Rice SA, Baden JM: Halothane, isoflurane, and enflurane MAC in pregnant and nonpregnant female and male mice and rats. *ANESTHESIOLOGY* 1985; 62:339-41
14. Shibahara S, Müller R, Taguchi H, Yoshida T: Cloning and expression of cDNA for rat heme oxygenase. *Proc Natl Acad Sci U S A* 1985; 82:7865-9
15. Odaka Y, Takahashi T, Yamasaki A, Suzuki T, Fujiwara T, Yamada T, Hirakawa M, Fujita H, Ohmori E, Akagi R: Prevention of halothane-induced hepatotoxicity by hemin pretreatment: Protective role of heme oxygenase-1 induction. *Biochem Pharmacol* 2000; 59:871-80
16. Yamasaki A, Takahashi T, Suzuki T, Fujiwara T, Hirakawa M, Ohmori E, Akagi R: Differential effects of isoflurane and halothane on the induction of heat shock proteins. *Biochem Pharmacol* 2001; 62:375-82
17. Kharasch ED, Thummel KE: Identification of cytochrome P450 2E1 as the predominant enzyme catalyzing human liver microsomal defluorination of sevoflurane, isoflurane, and methoxyflurane. *ANESTHESIOLOGY* 1993; 79:795-807
18. Ghantous HN, Fernando J, Gandolfi AJ, Brendel K: Minimal biotransformation and toxicity of desflurane in guinea pig liver slices. *Anesth Analg* 1991; 72:796-800
19. Suttner SW, Schmidt CC, Boldt J, Hüttner I, Kumle B, Piper SN: Low-flow desflurane and sevoflurane anesthesia minimally affect hepatic integrity and function in elderly patients. *Anesth Analg* 2000; 91:206-12
20. Lin WQ, Van Dyke RA, Marsh HM, Trudell JR: Time course of 72-kilodalton heat shock protein induction and appearance of trifluoroacetyl adducts in livers of halothane-exposed rats. *Mol Pharmacol* 1994; 46:639-43
21. Van Dyke RA, Mostafapour S, Marsh HM, Li Y, Chopp M: Immunocytochemical detection of the 72-kDa heat shock protein in halothane-induced hepatotoxicity in rats. *Life Sci* 1992; 50:PL41-5