

α_2 Adrenoceptor-mediated Presynaptic Inhibition of Primary Afferent Glutamatergic Transmission in Rat Substantia Gelatinosa Neurons

Yasuhiko Kawasaki, M.D.,* Eiichi Kumamoto, Ph.D.,† Hidemasa Furue, Ph.D.,‡ Megumu Yoshimura, M.D., Ph.D.§

Background: Although intrathecal administration of norepinephrine is known to produce analgesia, cellular mechanisms for this action have not yet been fully understood.

Methods: The actions of norepinephrine (50 μ M) on glutamatergic transmission were examined by using the whole cell patch clamp technique in substantia gelatinosa neurons of an adult rat spinal cord slice with an attached dorsal root.

Results: Norepinephrine inhibited the amplitude of monosynaptically evoked A δ -fiber and C-fiber excitatory postsynaptic currents in a reversible manner. When compared in magnitude between the A δ -fiber and C-fiber excitatory postsynaptic currents, the former inhibition ($50 \pm 4\%$, $n = 20$) was significantly larger than the latter one ($28 \pm 4\%$, $n = 8$). Both actions of norepinephrine were mimicked by an α_2 adrenoceptor agonist, clonidine (10 μ M), and an α_{2A} agonist, oxymetazoline (10 μ M), but not by an α_1 agonist, phenylephrine (10 μ M), and a β agonist, isoproterenol (40 μ M). The inhibitory actions were antagonized by an α_2 antagonist, yohimbine (1 μ M), all of the results of which indicate an involvement of α_2 adrenoceptors. Norepinephrine did not affect the amplitude of miniature excitatory postsynaptic current and of a response of substantia gelatinosa neurons to AMPA, indicating that its action on evoked excitatory postsynaptic currents is presynaptic in origin.

Conclusions: Norepinephrine inhibits A δ -fiber- and C-fiber-mediated sensory transmission to substantia gelatinosa neurons through the activation of the α_2 adrenoceptor (possibly α_{2A} type, based on the current, published behavioral and anatomical data) existing in primary afferent terminals; this action of norepinephrine is more effective in A δ -fiber than C-fiber transmission. This could contribute to at least a part of inhibitory modulation of pain sensation in the substantia gelatinosa by intrathecally administered norepinephrine.

CATECHOLAMINES, such as epinephrine and phenylephrine, have been used in spinal anesthesia together with local anesthetics with an expectation of a contraction of local vessels by the monoamines resulting in a decrease in the clearance of the anesthetics from the subarachnoid space and thus antinociception,^{1,2} while intrathecal administration of norepinephrine itself is

known to have an antinociceptive effect when assessed by the tail-flick and hot-plate tests.^{3,4} There is much evidence supporting the latter idea. Nociceptive information is transmitted through thinly myelinated A δ -afferent and unmyelinated C-afferent fibers from the periphery to the spinal cord, especially substantia gelatinosa (SG) neurons,^{5,6} where the information is modulated. Among this modulatory system, there is a descending norepinephrine-containing fiber pathway from cell groups designated A5, A6 (nucleus locus ceruleus), and A7 (subceruleus) in the pons,⁷⁻¹⁰ electrical stimulation of which results in behavioral analgesia.¹¹⁻¹⁴ This norepinephrine pathway is known to be also activated by systemically administered opioids^{15,16} or electrical stimulation of the midbrain periaqueductal gray region.^{17,18} *In situ* hybridization and immunohistochemical studies have demonstrated the presence of adrenoceptors in dorsal root ganglion and dorsal horn neurons in the rat,^{19,20} suggesting a role of norepinephrine at presynaptic and postsynaptic sites in the modulation. Previous studies have reported in SG neurons of the spinal cord and also spinal trigeminal nucleus that norepinephrine hyperpolarizes membrane,^{21,22} potentiates inhibitory (γ -aminobutyric acid-mediated and glycinergic) transmission,²³ and inhibits glutamatergic excitatory transmission.²⁴ The former two actions have been examined in detail, whereas the last one remains to be examined in detail. Although there is known to be a difference between A δ -fiber and C-fiber excitatory transmission to SG neurons in the action of neuromodulators, such as a γ -aminobutyric acid type B receptor agonist, baclofen,²⁵ serotonin,²⁶ nociceptin,²⁷ and anandamide,²⁸ and also of capsaicin,²⁹ this has not yet been revealed for the norepinephrine action because Travagli and Williams²⁴ examined the action of norepinephrine on excitatory transmission evoked in spinal trigeminal SG neurons by stimulating the trigeminal tract where the stimulation could not be separately given to each of the fibers. It has not yet been unveiled which types of adrenoceptors engage in modulation by norepinephrine of each of the A δ -fiber and C-fiber transmissions, although the norepinephrine-induced hyperpolarization and presynaptic facilitation of inhibitory transmission are reported to be due to the activation of α_2 and α_1 adrenoceptors, respectively.²¹⁻²³ In the current study, we examined the effect of norepinephrine on monosynaptic A δ -fiber and C-fiber excitatory postsynaptic currents (EPSCs) and its pharmacological property in the adult rat SG using the whole

* Graduate Student, † Professor, Department of Physiology, Saga Medical School. ‡ Lecturer, Department of Physiology, Saga Medical School. Current position: Department of Physiology, Kyusyu University School of Medicine, Kyusyu, Japan. § Professor, Department of Physiology, Saga Medical School. Current position: Department of Physiology, Kyusyu University School of Medicine, Kyusyu, Japan.

Received from the Department of Physiology, Saga Medical School, Saga, Japan. Submitted for publication May 15, 2002. Accepted for publication October 25, 2002. Support was provided solely from institutional and/or departmental sources. Presented as abstracts at the 30th annual meeting of the Society for Neuroscience, New Orleans, Louisiana, November 6, 2000, and at the 31st annual meeting of the Society for Neuroscience, San Diego, California, November 15, 2001.

Address reprint requests to Dr. Kumamoto: Department of Physiology, Saga Medical School, 5-1-1 Nabeshima, Saga 849-8501, Japan. Address electronic mail to: kumamoto@post.saga-med.ac.jp. Individual article reprints may be purchased through the Journal Web site, www.anesthesiology.org.

cell patch clamp technique under the condition of a blockade of the hyperpolarizing action.

Materials and Methods

Preparation of Spinal Cord Slices

This study was approved by the institutional Animal Use and Care Committee at Saga Medical School (Saga, Japan). The technique used for obtaining slice preparations from the rat spinal cord was the same as that described elsewhere.^{27,30} Briefly, adult male Sprague-Dawley rats (7–8 weeks old) were deeply anesthetized with urethane (1.2 g/kg). The lumbosacral spinal cord (L1–S3) was then removed and placed in preoxygenated (95% O₂ and 5% CO₂) Krebs solution (117 mM NaCl, 3.6 mM KCl, 2.5 mM CaCl₂, 1.2 mM MgCl₂, 1.2 mM NaH₂PO₄, 25 mM NaHCO₃, and 11 mM glucose) at 1–3°C; the rats were then immediately sacrificed by exsanguination. After cutting all of ventral and dorsal roots near the root entry zone, except for the L4 or L5 dorsal root on one side, the pia-arachnoid membrane was removed. The spinal cord was mounted on a Vibratome Series 1000® (Technical Products International, O'Fallon, MO), and then a 600- to 650- μ m-thick transverse slice was cut with an attached dorsal root having a length of 6–14 mm. The slice was placed on a nylon mesh in the recording chamber and then perfused at a rate of 15–20 ml/min with Krebs solution maintained at 36 \pm 1°C. Before the start of the experiment, the slice was preincubated for at least 1 h with Krebs solution.

Whole Cell Recordings from Substantia Gelatinosa Neurons and Stimulation of the Dorsal Root

Substantia gelatinosa neurons were identified by their location under a binocular microscope with light transmitted from below as reported previously^{25,27,31}; the SG was discernible as a relatively translucent band. Blind whole cell voltage clamp recordings were made from neurons that are located at the center of SG, with a patch pipette that was made up of a thin-walled fiberglass (1.5-mm OD) using a single-stage horizontal puller (P-97; Sutter Instrument, Novato, CA). The patch pipettes had a tip resistance of 5–10 M Ω when filled with a solution having the following composition: 110 mM Cs₂SO₄, 0.5 mM CaCl₂, 2 mM MgCl₂, 5 mM EGTA, 5 mM HEPES, 5 mM Mg-ATP, 5 mM tetraethylammonium, and 1 mM guanosine 5'-O-(2-thiodiphosphate), where guanosine 5'-O-(2-thiodiphosphate) and K⁺-channel blockers (Cs⁺ and tetraethylammonium) were added to inhibit a hyperpolarizing effect of norepinephrine through the action of G proteins and to block an activation of K⁺ channels, respectively. After making a rigid seal (resistances: 5–20 G Ω) in the cell-attached mode by a gentle suction given into the patch pipette, the membrane patch was ruptured by a brief period of more powerful suction, resulting in the

whole cell configuration. Signals were amplified with a patch clamp amplifier (Axopatch 200B; Axon Instruments, Foster City, CA) in the voltage clamp mode. Data were low-pass filtered at 5 kHz, digitized at 333 kHz with an A/D converter, and stored and analyzed with a personal computer using the pCLAMP data acquisition program (version 6.0; Axon Instruments). The program used for analyzing miniature EPSCs (mEPSCs) detects spontaneous events if the difference between the baseline and a following current value exceeds a given threshold of 6 pA and separating valleys are less than 50% of adjacent peaks; the validity of the method was confirmed by measuring visually individual mEPSCs on a fast time scale in several cases.³⁰

Orthodromic stimulation of the dorsal root to elicit EPSCs was performed with a suction electrode at 0.2 Hz unless otherwise mentioned. The strength of stimuli used was 1.2 times the threshold to elicit EPSCs, fearing a conduction block of action potentials in the dorsal root. The holding potential (V_H) used was –70 mV, at which glycine and γ -aminobutyric acid type A receptor-mediated synaptic currents were invisible. The duration of stimuli used was 0.1 ms throughout the experiments, and conduction velocities were calculated from the latency of monosynaptic EPSC and the length of dorsal root. A δ -fiber and C-fiber evoked EPSCs were distinguished from each other on the basis of the conduction velocity of afferent fibers and stimulus threshold; they were considered as monosynaptic in origin when the latency remained constant and there was no failure during stimulation at 20 Hz for 1 s or when failures did not occur during repetitive stimulation at 1 Hz for 20 s, respectively, as reported previously.^{25,26,32,33} During these stimulations, a conduction block of action potentials did not occur when examined by using rat dorsal root ganglion neurons.^{25,33} Although in some neurons monosynaptic A δ -fiber or C-fiber EPSCs were accompanied by polysynaptic A δ -fiber or C-fiber EPSCs, such neurons were not used for analysis if the peak of the monosynaptic EPSCs was contaminated by the polysynaptic EPSCs.

Application of Drugs

Drugs were dissolved in Krebs solution and then applied to the SG by an exchange of superfusing solution *via* a three-way stopcock with one containing them at known concentrations without a change in superfusion rate and thus in temperature. Solutions in the recording chamber completely altered within 20 s. The drugs used in this work were (\pm)-norepinephrine (Aldrich Chemical Co., Milwaukee, WI), yohimbine hydrochloride, (–)-phenylephrine (Wako, Osaka, Japan), α -amino-3-hydroxy-5-methyl-4-isoxazolepropionate (AMPA), (–)-isoproterenol hydrochloride, clonidine hydrochloride, oxymetazoline hydrochloride, guanosine 5'-O-(2-thiodiphosphate) (Sigma, St Louis, MO), and 6-cyano-7-nitroquinoxaline-2,3-dione (Tocris Cookson, St. Louis, MO).




Fig. 1. Monosynaptic excitatory postsynaptic currents (EPSCs) evoked in substantia gelatinosa neurons by stimulating the dorsal root. (A and B) Two types of monosynaptic EPSCs evoked at higher frequencies (*left*: 20 Hz in A and 1 Hz in B) and 0.2 Hz (*right*), which are different in latency from an artifact of the stimulus. A δ -fiber and C-fiber EPSCs were distinguished from each other on the basis of the conduction velocity of afferent fibers and stimulus threshold (9.6 m/s and 62 μ A and 0.5 m/s and 560 μ A in A and B, respectively). The A δ or C responses, respectively, were considered as monosynaptic in origin when there were no failures and the latency remained constant during stimulation at 20 Hz or when failures did not occur during stimulation at 1 Hz. Both of the monosynaptic A δ -fiber and C-fiber EPSCs were observed in 31% of neurons examined (C; stimulus strength, 600 μ A). Each of the records is a superimposition of seven or eight traces of EPSCs; $V_H = -70$ mV.

Statistical Analysis

Data are presented as mean \pm SEM. Statistical significance was determined as $P < 0.05$ using either the paired or unpaired Student t test unless otherwise mentioned. The Kolmogorov-Smirnov test was also used to compare mEPSCs in the absence and presence of norepinephrine in the distributions of their amplitudes and interevent intervals. In all cases, n refers to the number of neurons studied.

Results

Whole cell patch clamp recordings were made from a total of 158 SG neurons. Stable recordings could be obtained from neurons in spinal cord slices maintained *in vitro* for more than 12 h, and recordings were made from single neurons for up to 4 h. All experiments were performed at least 10 min later, enough time for guanosine 5'-O-(2-thiodiphosphate) and K^+ -channel blockers in patch pipette solutions to diffuse into SG neurons, after the establishment of whole cell configuration; norepinephrine (50 μ M) did not change holding currents at -70 mV. In all SG neurons tested, mEPSCs were downward at -70 mV and were blocked by 6-cyano-7-nitroquinoxaline-2,3-dione (10 μ M; $n = 2$), as reported previously,^{27,30} indicating the activation of AMPA receptors.

More Inhibition by Norepinephrine of Monosynaptic A δ -fiber Than C-fiber EPSCs Elicited in Substantia Gelatinosa Neurons

When stimulating the dorsal root with a strength of less than 200 μ A, 82% ($n = 75$; this percentage was almost comparable to a value [79%] obtained previously²⁷) of 92 neurons examined could elicit monosynaptic EPSCs that had no failure and no change in latency when elicited at 20 Hz (fig. 1A, left). Conduction velocity values (3.8–15 m/s), estimated from the latency of EPSC and the length of dorsal root, were almost within a range of those of A δ -fibers obtained from the experiment in dorsal root ganglion neurons, as reported previously.^{25,32} Monosynaptic A δ -afferent EPSCs evoked at 0.2 Hz had a mean amplitude of 279 ± 23 pA (range, 59–884 pA; $V_H = -70$ mV; fig. 1A, right). On the other hand, when stimulated with a strength of more than 200 μ A, 83% ($n = 53$; this percentage was also similar to a value [80%] obtained previously²⁷) of 64 neurons tested exhibited monosynaptic EPSCs that had no failures in response to stimuli at 1 Hz (fig. 1B, left), albeit a variability in EPSC latency was observed in some neurons.²⁵ The EPSCs originated from afferent fibers whose conduction velocities were 0.3–0.8 m/s, a range of those of C fibers.^{25,32} The amplitude of monosynaptic C-afferent EPSCs evoked at 0.2 Hz averaged to be 253 ± 26 pA (range, 96–714 pA; $V_H = -70$ mV; fig. 1B, right). Thirty-one percent ($n = 20$) of the neurons tested exhibited both of the monosynaptic




Fig. 2. Actions of norepinephrine ($50 \mu\text{M}$) on monosynaptic excitatory postsynaptic currents (EPSCs) evoked in substantia gelatinosa neurons. (*A* and *B*) A δ -fiber and C-fiber EPSCs, respectively, in the absence (*left* and *right*) and presence (*middle*; where control EPSC is superimposed for comparison) of norepinephrine. Note that the A δ -fiber EPSC was more sensitive to norepinephrine than the C-fiber EPSC. This was so in a neuron in which both of the A δ -fiber and C-fiber EPSCs were observed (*C*). Each record in *A*, *B*, and *C* (stimulus strength: 52, 250, and $790 \mu\text{A}$, respectively) shows an average of 12 traces of EPSCs; $V_H = -70 \text{ mV}$.

A δ -fiber and C-fiber EPSCs, as seen in figure 1C. These EPSCs were completely inhibited by 6-cyano-7-nitroquinoxaline-2,3-dione ($10 \mu\text{M}$; $n = 2$), indicative of an involvement of AMPA receptors, as reported previously.^{27,32}


Effects of norepinephrine ($50 \mu\text{M}$) on the monosynaptic A δ -fiber and/or C-fiber EPSCs were examined in a total of 24 SG neurons. As seen from figures 2A–C, each of the EPSCs was inhibited in amplitude by norepinephrine in a reversible manner. These effects on A δ -fiber and C-fiber EPSCs were, respectively, maximal at approximately 1 and 2 min following the application of norepinephrine (fig. 3A). When examined in many neurons, the inhibitions (measured at approximately 2 min following norepinephrine superfusion) of the peak amplitudes of A δ -fiber and C-fiber EPSCs were, respectively, $50 \pm 4\%$ ($n = 20$) and $28 \pm 4\%$ ($n = 8$), with the values being significantly different from each other ($P < 0.01$), as

seen in figure 3B. This greater sensitivity of A δ -fiber EPSC than C-fiber EPSC to norepinephrine was also observed in a neuron in which both of the EPSCs were elicited ($n = 5$; for example, see fig. 2C).

Involvement of α_2 Adrenoceptors in Norepinephrine-induced Inhibition of Monosynaptic A δ -fiber and C-fiber EPSCs Evoked in Substantia Gelatinosa Neurons

We next examined which subtypes of adrenoceptors are involved in the norepinephrine-induced inhibition of A δ -fiber and C-fiber EPSCs by use of their agonists and antagonists. When superfused for 2 min, an α_2 adrenoceptor agonist, clonidine ($10 \mu\text{M}$), as well as norepinephrine, inhibited the peak amplitude of monosynaptically evoked A δ -fiber and C-fiber EPSCs (by $24 \pm 2\%$ [$n = 13$] and $19 \pm 3\%$ [$n = 11$], respectively), as seen in figures

Fig. 3. Comparison of the action of norepinephrine ($50 \mu\text{M}$) between A δ -fiber and C-fiber excitatory postsynaptic currents (EPSCs) in substantia gelatinosa neurons. (*A*) Time courses of changes in the peak amplitude of A δ -fiber EPSC (\circ) and C-fiber EPSC (\bullet) under the action of norepinephrine, relative to that in the control. There was a time delay of 1–2 min before the norepinephrine action shows a maximal effect. Each of them was obtained from a different neuron. Each point is an average of the amplitudes of three consecutive EPSCs. (*B*) Relative peak amplitude of A δ -fiber and C-fiber EPSC under the action of norepinephrine to that in the control. The number of neurons examined is shown in parentheses; $*P < 0.01$; $V_H = -70 \text{ mV}$.






Fig. 4. A δ -fiber excitatory postsynaptic currents (EPSCs) (A) and C-fiber EPSCs (B) evoked in substantia gelatinosa neurons in the absence (left) and presence (right; where control EPSC is superimposed for comparison) of an α_2 agonist, clonidine (10 μ M; a), an α_{2A} agonist, oxymetazoline (10 μ M; b), an α_1 agonist, phenylephrine (10 μ M; c), and a β agonist, isoproterenol (40 μ M; d). Note that both of the EPSCs were depressed by clonidine and oxymetazoline but not phenylephrine and isoproterenol. Each of the records is an average of 12 traces of EPSCs; $V_H = -70$ mV.

4Aa and Ba. An agonist of α_{2A} adrenoceptor, oxymetazoline (10 μ M),³⁴ exhibited a similar action on A δ -fiber and C-fiber EPSCs (inhibition: $34 \pm 9\%$ [$n = 4$] and $16 \pm 2\%$ [$n = 4$], respectively; these were not different from each other; $P > 0.05$; figs. 4Ab and Bb). On the other hand, an α_1 adrenoceptor agonist, phenylephrine (10 μ M), and a β adrenoceptor agonist, isoproterenol (40 μ M), were without the inhibition on A δ -fiber and C-fiber EPSCs, as seen in figures 4Ac, Bc, Ad, and Bd. Figure 5 summarizes the effects of the adrenoceptor agonists on A δ -fiber and C-fiber EPSCs. With respect to antagonists, an α_2 adrenoceptor antagonist, yohimbine (1 μ M), superfused for 5 min prior to the application of norepinephrine (50

μ M), greatly depressed its inhibitory effect on A δ -fiber or C-fiber EPSCs (peak amplitude: $95 \pm 3\%$ [$n = 12$] and $97 \pm 6\%$ [$n = 4$] of control, respectively, in the presence of the antagonist), as seen in figures 6A and B. Altogether, these results indicate that the suppressive actions of norepinephrine on A δ -fiber and C-fiber EPSCs are mediated by α_2 adrenoceptors.

Lack of the Effect of Norepinephrine on AMPA Responses in Substantia Gelatinosa Neurons

Figures 7A and B demonstrate the effect of norepinephrine (50 μ M) superfused for 2 min on mEPSCs recorded from SG neurons. Both mEPSC amplitude and





Fig. 5. Effects of various α adrenoceptor agonists and β adrenoceptor agonist on A δ -fiber and C-fiber excitatory postsynaptic currents (EPSCs) in substantia gelatinosa neurons. (A and B) Relative peak amplitudes of A δ -fiber EPSC (A) and C-fiber EPSC (B) in the presence of an α_1 agonist, phenylephrine (10 μ M; α_1), an α_2 agonist, clonidine (10 μ M; α_2), an α_{2A} agonist, oxymetazoline (10 μ M; α_{2A}), or a β agonist, isoproterenol (40 μ M; β) to that in the control (Ctl). Note that the α_2 and α_{2A} but not α_1 and β agonists significantly inhibited both EPSCs. The number of neurons examined is shown in parentheses; * $P < 0.01$.

Fig. 6. Effect of norepinephrine ($50 \mu\text{M}$) on A δ -fiber and C-fiber excitatory postsynaptic currents (EPSCs) evoked in substantia gelatinosa neurons in the presence of an α_2 adrenoceptor antagonist, yohimbine ($1 \mu\text{M}$). (A and B) A δ -fiber and C-fiber EPSCs, respectively, in the control (left) and in the presence of norepinephrine together with yohimbine (right; where control EPSC is superimposed for comparison). Note that norepinephrine had no effects on both EPSCs in the presence of yohimbine. Each of the records is an average of 12 traces of EPSCs; $V_H = -70 \text{ mV}$.



frequency were unaffected by norepinephrine; they were $98 \pm 7\%$ ($P > 0.05$; $n = 10$) and $102 \pm 1\%$ ($P > 0.05$; $n = 10$) of control ($17 \pm 3 \text{ pA}$ and $9 \pm 2 \text{ Hz}$), respectively, as reported by Baba *et al.*²³ When examined for cumulative distributions of the amplitude and interevent interval of mEPSC, they were also unaffected by norepinephrine, as seen from figure 7C. To determine any effect of norepinephrine on the sensitivity of SG neurons to L-glutamate, we examined whether an AMPA

($10 \mu\text{M}$) response is affected by norepinephrine ($50 \mu\text{M}$). As seen in figure 7D, norepinephrine did not affect the peak amplitude of the AMPA response ($88 \pm 5\%$ of control, $n = 12$; $P > 0.05$).

Discussion

The current study demonstrated that norepinephrine inhibits glutamatergic excitatory transmission to SG neu-




Fig. 7. Norepinephrine ($50 \mu\text{M}$) did not affect a sensitivity of substantia gelatinosa neurons to AMPA. (A) Chart recording of miniature excitatory postsynaptic currents (mEPSCs) in the absence and presence of norepinephrine. (B) mEPSCs, shown in an expanded time scale, in the absence (left) and presence (right) of norepinephrine for a period indicated by a mark shown below the chart recording in A. (C) Cumulative distributions of the amplitude (left) and interevent interval (right) of mEPSCs before (continuous line) and during (dotted line) 1 min after the application of norepinephrine (NE), which were obtained by analyzing 645 and 702 mEPSC events, respectively. Norepinephrine had no effect on their distributions ($P > 0.05$; Kolmogorov-Smirnov test). (D) AMPA ($10 \mu\text{M}$) responses, obtained by its superfusion for 30 s, in the absence and presence of norepinephrine. $V_H = -70 \text{ mV}$.

rons in the spinal dorsal horn from the periphery. It was revealed here for the first time that both of A δ and C primary afferent transmission are depressed by norepinephrine, although Travagli and Williams²⁴ had not examined the action of norepinephrine on each of their transmissions to SG neurons of the spinal trigeminal nucleus. Both actions of norepinephrine were due to the activation of α_2 adrenoceptors, as reported for evoked EPSCs in spinal trigeminal SG neurons.²⁴ This result may be consistent with the observations that an antinociceptive effect of either intrathecally administered norepinephrine or electrical stimulation of sites near the A7 cell group and also of the periaqueductal gray is due to the activation of α_2 adrenoceptors.^{4,14,16,17} Although the α_2 adrenoceptors are subdivided into α_{2A} , α_{2B} , and α_{2C} receptors,³⁴ the α_2 action in the current study appears to be due to the activation of the α_{2A} type because an α_{2A} adrenoceptor agonist, oxymetazoline,³⁴ reduced the A δ and C afferent transmission. This idea is supported by a report of Stone *et al.*³⁵ which demonstrated that a mouse expressing a point mutation in the α_{2A} receptor was without α_2 agonist-mediated spinal analgesia in the tail-flick test. Although the activation of β adrenoceptors is known to enhance neurotransmitter release in many types of neurons, including sympathetic ganglion neurons,³⁶ this was not the case in SG neurons because isoproterenol did not affect the transmission.

Since norepinephrine did not affect mEPSC amplitude and the sensitivity of SG neurons to AMPA, its actions on evoked EPSCs were presynaptic in origin. Since these actions were examined for monosynaptic transmission, the norepinephrine actions are suggested to be due to the activation of α_2 , possibly α_{2A} , adrenoceptors existing in primary afferent central terminals. Consistent with this idea, an inhibition by norepinephrine of the release of L-glutamate from rat spinal cord synaptosomes is mediated by α_2 adrenoceptors,³⁷ and the α_{2A} adrenoceptor is expressed in primary afferent terminals in the rat.²⁰ The action on C-fiber transmission in the current study may be consistent with the observation that capsaicin-induced release of L-glutamate from spinal cord synaptosomes is reduced by the activation of the α_{2A} adrenoceptor³⁸ because capsaicin is known to excite C fibers. Although a cellular mechanism for the norepinephrine action was not examined here, this would be due to an inhibition of voltage-gated Ca²⁺ channels existing in nerve terminals because norepinephrine reduces Ca²⁺-channel currents in many types of neurons, including dorsal root ganglion neurons.³⁹ There was a dissociation between the actions of norepinephrine on evoked EPSC and mEPSC in that the former was presynaptically depressed, while mEPSC frequency was unaffected, an observation different from that in spinal trigeminal SG neurons.²⁴ One explanation for this discrepancy may be that norepinephrine acts differentially on evoked and spontaneous transmitter release mechanisms in the spi-

nal dorsal horn, resulting in a distinct action on their EPSCs, because each of the releases in the SG is suggested to be mediated by different types of Ca²⁺ channels.⁴⁰ Alternatively, mEPSCs in spinal cord SG neurons may be produced by inputs not only from primary afferent fiber axon but also from interneuron axon, the former terminals having more α_2 adrenoceptors than the latter ones, where mEPSCs mainly originate from interneuron terminals. This idea may be consistent with the observation that many α_{2A} adrenoceptors are colocalized in the spinal dorsal horn with neuropeptides, which are contained in primary afferent fibers.²⁰ A similar discrepancy between spontaneous and evoked transmission has been seen in the actions of a μ -opioid receptor agonist³¹ and anandamide²⁸ in spinal cord SG neurons. Although Pan *et al.*⁴¹ have very recently reported a clonidine-induced decrease in mEPSC frequency in SG neurons, a discrepancy between this and the current study may be due to the fact that different SG neurons were tested because they examined neurons in the outer layer of SG, while we investigated neurons located at the center of SG. The possibility cannot be ruled out that SG neurons exhibiting no effect of norepinephrine on mEPSCs in the current study (where the blind patch clamp technique was used) had located in the inner layer of SG because visually identified neurons in the inner layer of SG appeared to be without actions of clonidine on mEPSCs.⁴¹

The current study revealed for the first time that norepinephrine inhibits A δ -fiber transmission more effectively than C-fiber transmission. There are two possible explanations for this result. One is that α_2 adrenoceptors are more densely expressed in A δ -fiber than C-fiber terminals in the spinal dorsal horn. The other is that there is a different type of α_2 adrenoceptors in each of the A δ -fiber and C-fiber terminals, although oxymetazoline could not discriminate between them in the extent of inhibition. This remains to be examined by using other agents regarding subtypes of α_2 adrenoceptors. The action of norepinephrine was the same as that of anandamide²⁸ but different from those of baclofen²⁵ and nociceptin²⁷ in that A δ -fiber transmission was more sensitive than C-fiber transmission. It is suggested that norepinephrine as well as anandamide may inhibit fast-conducting transmission more potently than slow-conducting pain transmission.

In conclusion, the current study provides a cellular basis for the antinociceptive action of norepinephrine through a mechanism in primary afferent terminals at the spinal cord level. Although norepinephrine may contribute to a prolongation of analgesia through its vasoconstrictive action in spinal anesthesia, the current finding of the inhibition of excitatory transmission supports its role as an important negative modulator of pain transmission to SG neurons together with a norepinephrine-induced hyperpolarization^{21,22} and enhancement of inhibitory transmission.²³

The authors thank Tadahide Totoki, M.D., Ph.D. (Professor of Anesthesiology, Saga Medical School, Saga, Japan), for his encouragement during this study.

References

1. Concepcion M, Maddi R, Francis D, Rocco AG, Murray E, Convino BG: Vasoconstrictors in spinal anesthesia with tetracaine: A comparison of epinephrine and phenylephrine. *Anesth Analg* 1984; 63:134-8
2. Vaida GT, Moss P, Capan LM, Turndorf H: Prolongation of lidocaine spinal anesthesia with phenylephrine. *Anesth Analg* 1986; 65:781-5
3. Reddy SVR, Maderdrut JL, Yaksh TL: Spinal cord pharmacology of adrenergic agonist-mediated antinociception. *J Pharm Exp Ther* 1980; 213:525-33
4. Howe JR, Wang J-Y, Yaksh TL: Selective antagonism of the antinociceptive effect of intrathecally applied alpha adrenergic agonists by intrathecal prazosin and intrathecal yohimbine. *J Pharm Exp Ther* 1983; 224:552-8
5. Kumazawa T, Perl ER: Excitation of marginal and substantia gelatinosa neurons in the primate spinal cord: Indications of their place in dorsal horn functional organization. *J Comp Neurol* 1978; 177:417-34
6. Willis Jr WD, Coggeshall RE: *Sensory Mechanisms of the Spinal Cord*, 2nd edition. Edited by Willis WD Jr, Coggeshall RE. New York, Plenum Press, 1991, pp 94-115
7. Satoh K, Kashiba A, Kimura H, Maeda T: Noradrenergic axon terminals in the substantia gelatinosa of the rat spinal cord: An electron-microscopic study using glyoxylic acid-potassium permanganate fixation. *Cell Tissue Res* 1982; 222:359-78
8. Westlund KN, Bowker RM, Ziegler MG, Coulter JD: Noradrenergic projections to the spinal cord of the rat. *Brain Res* 1983; 263:15-31
9. Hagiwara S, Senba E, Yoshida S, Tohyama M, Yoshiya I: Fine structure of noradrenergic terminals and their synapses in the rat spinal dorsal horn: An immunohistochemical study. *Brain Res* 1990; 526:73-80
10. Rajaofetra N, Ridet J-L, Poulat P, Marlier L, Sandillon F, Geffard M, Privat A: Immunocytochemical mapping of noradrenergic projections to the rat spinal cord with an antiserum against noradrenaline. *J Neurocytol* 1992; 21:481-94
11. Jones SL, Gebhart GF: Quantitative characterization of ceruleospinal inhibition of nociceptive transmission in the rat. *J Neurophysiol* 1986; 56:1397-410
12. Proudfit HK: Pharmacologic evidence for the modulation of nociception by noradrenergic neurons. *Prog Brain Res* 1988; 77:357-70
13. Jones SL: Descending noradrenergic influences on pain. *Prog Brain Res* 1991; 88:381-94
14. Yeomans DC, Clark FM, Paice JA, Proudfit HK: Antinociception induced by electrical stimulation of spinally projecting noradrenergic neurons in the A7 catecholamine cell group of the rat. *Pain* 1992; 48:449-61
15. Wilson PR, Yaksh TL: Pharmacology of pain and analgesia. *Anaesth Intensive Care* 1980; 8:248-56
16. Yaksh TL: Pharmacology of spinal adrenergic systems which modulate spinal nociceptive processing. *Pharmacol Biochem Behav* 1985; 22:845-58
17. Aimone LD, Jones SL, Gebhart GF: Stimulation-produced descending inhibition from the periaqueductal gray and nucleus raphe magnus in the rat: Mediation by spinal monoamines but not opioids. *Pain* 1987; 31:123-36
18. Bajic D, Proudfit HK: Projections of neurons in the periaqueductal gray to pontine and medullary catecholamine cell groups involved in the modulation of nociception. *J Comp Neurol* 1999; 405:359-79
19. Nicholas AP, Pieribone V, Hökfelt T: Distributions of mRNAs for alpha-2 adrenergic receptor subtypes in rat brain: An in situ hybridization study. *J Comp Neurol* 1993; 328:575-94
20. Stone LS, Broberger C, Vulchanova L, Wilcox GL, Hökfelt T, Riedl MS, Elde R: Differential distribution of α_{2A} and α_{2C} adrenergic receptor immunoreactivity in the rat spinal cord. *J Neurosci* 1998; 18:5928-37
21. North RA, Yoshimura M: The actions of noradrenaline on neurones of the rat substantia gelatinosa *in vitro*. *J Physiol* 1984; 349:43-55
22. Grudt TJ, Williams JT, Travagli RA: Inhibition by 5-hydroxytryptamine and noradrenaline in substantia gelatinosa of guinea-pig spinal trigeminal nucleus. *J Physiol* 1995; 485:113-20
23. Baba H, Shimoji K, Yoshimura M: Norepinephrine facilitates inhibitory transmission in substantia gelatinosa of adult rat spinal cord: I. Effects on axon terminals of GABAergic and glycinergic neurons. *ANESTHESIOLOGY* 2000; 92:473-84
24. Travagli RA, Williams JT: Endogenous monoamines inhibit glutamate transmission in the spinal trigeminal nucleus of the guinea-pig. *J Physiol* 1996; 491:177-85
25. Ataka T, Kumamoto E, Shimoji K, Yoshimura M: Baclofen inhibits more effectively C-afferent than A δ -afferent glutamatergic transmission in substantia gelatinosa neurons of adult rat spinal cord slices. *Pain* 2000; 86:273-82
26. Ito A, Kumamoto E, Takeda M, Takeda M, Shibata K, Sagai H, Yoshimura M: Mechanisms for ovariectomy-induced hyperalgesia and its relief by calcitonin: Participation of 5-HT $_{1A}$ -like receptor on C-afferent terminals in substantia gelatinosa of the rat spinal cord. *J Neurosci* 2000; 20:6302-8
27. Luo C, Kumamoto E, Furue H, Chen J, Yoshimura M: Nociceptin inhibits excitatory but not inhibitory transmission to substantia gelatinosa neurones of adult rat spinal cord. *Neuroscience* 2002; 109:349-58
28. Luo C, Kumamoto E, Furue H, Chen J, Yoshimura M: Anandamide inhibits excitatory transmission to rat substantia gelatinosa neurones in a manner different from that of capsaicin. *Neurosci Lett* 2002; 321:17-20
29. Yang K, Kumamoto E, Furue H, Li Y-Q, Yoshimura M: Action of capsaicin on dorsal root-evoked synaptic transmission to substantia gelatinosa neurons in adult rat spinal cord slices. *Brain Res* 1999; 830:268-73
30. Lao L-J, Kumamoto E, Luo C, Furue H, Yoshimura M: Adenosine inhibits excitatory transmission to substantia gelatinosa neurons of the adult rat spinal cord through the activation of presynaptic A1 adenosine receptor. *Pain* 2001; 94:315-24
31. Kohno T, Kumamoto E, Higashi H, Shimoji K, Yoshimura M: Actions of opioids on excitatory and inhibitory transmission in substantia gelatinosa of adult rat spinal cord. *J Physiol* 1999; 518:803-13
32. Nakatsuka T, Park J-S, Kumamoto E, Tamaki T, Yoshimura M: Plastic changes in sensory inputs to rat substantia gelatinosa neurons following peripheral inflammation. *Pain* 1999; 82:39-47
33. Nakatsuka T, Ataka T, Kumamoto E, Tamaki T, Yoshimura M: Alteration in synaptic inputs through C-afferent fibers to substantia gelatinosa neurons of the rat spinal dorsal horn during postnatal development. *Neuroscience* 2000; 99:549-56
34. Bylund DB, Eikenberg DC, Hieble JP, Langer SZ, Lefkowitz RJ, Minneman KP, Molinoff PB, Ruffolo RR, Trendelenburg U: International union of pharmacology nomenclature of adrenoceptors. *Pharmacol Rev* 1994; 46:121-36
35. Stone LS, MacMillan LB, Kitto KF, Limbird LE, Wilcox GL: The α_{2A} adrenergic receptor subtype mediates spinal analgesia evoked by α_2 agonists and is necessary for spinal adrenergic-opioid synergy. *J Neurosci* 1997; 17:7157-65
36. Kumamoto E, Kuba K: Mechanism of long-term potentiation of transmitter release induced by adrenaline in bullfrog sympathetic ganglia. *J Gen Physiol* 1986; 87:775-93
37. Kamisaki Y, Hamada T, Maeda K, Ishimura M, Itoh T: Presynaptic α_2 adrenoceptors inhibit glutamate release from rat spinal cord synaptosomes. *J Neurochem* 1993; 60:522-6
38. Li X, Eisenach JC: α_2A -Adrenoceptor stimulation reduces capsaicin-induced glutamate release from spinal cord synaptosomes. *J Pharm Exp Ther* 2001; 299:939-44
39. Holz IV GG, Rane SG, Dunlap K: GTP-binding proteins mediate transmitter inhibition of voltage-dependent calcium channels. *Nature* 1986; 319:670-2
40. Bao J, Li JJ, Perl ER: Differences in Ca $^{2+}$ channels governing generation of miniature and evoked excitatory synaptic currents in spinal laminae I and II. *J Neurosci* 1998; 18:8740-50
41. Pan Y-Z, Li D-P, Pan H-L: Inhibition of glutamatergic synaptic input to spinal lamina II $_o$ neurons by presynaptic α_2 -adrenergic receptors. *J Neurophysiol* 2002; 87:1938-47